

Atmospheric ice-nucleating particles in the Eastern Mediterranean and the contribution of mineral and biological aerosol ~~potential influence of fertile soils~~

5 Mark D. Tarn¹, Bethany V. Wyld¹, Naama Reicher², Matan Alayof², Daniella Gat², Alberto Sanchez-Marroquin¹, Sebastien N. F. Sikora¹, Alexander D. Harrison¹, Yinon Rudich², Benjamin J. Murray¹

¹School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, United Kingdom

²Department of Earth and Planetary Sciences, Weizmann Institute of Science, Rehovot 76100, Israel

Correspondence to: Mark D. Tarn (m.d.tarn@leeds.ac.uk), Benjamin J. Murray (b.j.murray@leeds.ac.uk)

Abstract. While the atmosphere in the Eastern Mediterranean is part of the dust belt, it encounters air masses from Europe, the Mediterranean Sea, and the Sahara and Arabian Deserts that bring with them a whole host of potential dust and bioaerosol compositions and concentrations via long-range transport. The consequential changes in the populations of ice-nucleating particles (INPs), aerosols that influence weather and climate by the triggering of freezing in supercooled cloud water droplets, including in the convective cloud systems in the region, are not so well understood beyond the influence of desert dust storms in increasing INP concentrations. Here, we undertook an intensive INP measurement campaign in Israel to monitor changes in concentrations and activity from four major air masses, including the potential for activity from biological INPs. Our findings show that the INP activity in the region is likely dominated by the K-feldspar mineral content, with southwesterly air masses from the Sahara Desert and easterly air masses from the Arabian Deserts markedly increasing both aerosol and INP concentrations. Most intriguingly, a handful of air masses that passed over the Nile Delta and the northern Fertile Crescent, regions containing fertile agricultural soils and wetlands ~~a crescent shaped region of fertile soils, agriculture, wetlands, and marshes stretching from the Nile Delta and over northern Syria to eastern Iraq~~, brought high INP concentrations with strong indicators of biological activity. These results suggest that the Fertile Crescent could be a sporadic source of high-temperature biological ice-nucleating activity across the region that could periodically dominate the otherwise K-feldspar-controlled INP environment. We propose that these findings ~~and warrants~~ further exploration in future studies in the region, which ~~This may be~~ particularly pertinent ~~due~~ given the ongoing desertification of the Fertile Crescent that could reveal further sources of dust and fertile soil-based INPs in the future Eastern Mediterranean region.

1 Introduction

Clouds play an important role in weather and climate due to precipitation and their radiative properties, but their representation in cloud-feedback and climate models retains a high degree of uncertainty due ~~to poor representation of a number of~~ a deficiency in our understanding of a number of dynamic and microphysical processes, including

30 cloud-aerosol interactions (Ceppi et al., 2017; Storelvmo, 2017). A rare class of aerosol particles known as ice-nucleating particles (INPs) are able to catalyse the formation of ice in supercooled cloud water droplets at temperatures higher than around $-35\text{ }^{\circ}\text{C}$ (Murray et al., 2012; Hoose and Möhler, 2012; Kanji et al., 2017), below which cloud droplets can freeze homogeneously (Herbert et al., 2015; Rosenfeld and Woodley, 2000).

The presence of INPs can thus alter the ratio of water to ice in mixed-phase clouds and influence their albedo
35 and lifetime (Lohmann and Feichter, 2005). Therefore, an accurate representation of INPs in global models is important, particularly in a changing climate (Murray et al., 2021), but this is currently lacking (Kanji et al., 2017; Vergara-Temprado et al., 2017) since INPs are not only rare but also highly variable globally in terms of their activity, concentrations, transport, processing, sources and sinks.

To better characterise atmospheric INPs, it is necessary to undertake field campaigns that collect essential
40 and seasonal data on INP properties from around the globe and to determine the origins of those INPs and their effects on clouds. The Eastern Mediterranean is a fascinating location in terms of the varying air mass trajectories that reach it and the potential for vastly different INP ~~populationseommunities~~ associated with each, which could influence the clouds in the region, e.g. deep convective cloud systems (Funatsu et al., 2007). A large proportion of air flowing into deep convective clouds can come directly from the boundary layer, hence boundary layer aerosol are of direct
45 relevance for the development of deep convective clouds (Hoffmann et al., 2015; Hawker et al., 2021). Dayan (1986) studied five-day back trajectories from Israel over a five-year period and identified five major source regions of air masses: (i) northwest Europe, crossing the Mediterranean Sea, throughout the whole year, (ii) north-easterly continental flow originating in Eastern Europe during the summer season, (iii) infrequent south-easterly flow from the Syrian and Arabian Deserts that occurred mainly during Autumn (fall), (iv) south-westerly flow along the North
50 African coast most frequent during late winter and spring, and (v) south-southwesterly flow from inland North Africa that mostly occurred during winter and spring.

One of the major features of the region are the dust events associated with the ~~western-Sahara, Desert Syrian and Arabian Deserts~~ (Dayan et al., 1991; Ganor, 1994, 1991; Ganor and Mamane, 1982; Yaalon and Ganor, 1979; Mamane et al., 1980; Athanasopoulou et al., 2016; Reicher et al., 2018; Reicher et al., 2019)
55 ~~and the Syrian and Arabian Deserts~~. Transported dust can have residence times of hours to weeks (Ganor, 1991, 1994; Kubilay et al., 2000), and can cover heights of up to 6 km above mean sea level (a. m. s. l.) during transport (Israelevich et al., 2002). Dayan et al. (1991) found that Saharan events usually have high dust loadings and last for 2-4 days. Conversely, those from the Arabian Desert last for shorter periods (~ 1 -2 days) and are restricted to a more shallow atmospheric transport layer (up to ~ 2000 m a. m. s. l.) that may prevent their passing much beyond the Eastern Mediterranean.

60 Saharan dust events are important in terms of INP population and sources in the Eastern Mediterranean (Ganor, 1994), particularly as the number of dust days caused by African dust each year increased decade-by-decade until around 2010 (Ganor et al., 2010), although this trend has recently reversed (Nissenbaum et al., 2023). In addition to dust, westerly air flows can bring high levels of sea salt from the Mediterranean Sea (Levin and Lindberg, 1979;

Foner and Ganor, 1992) and biogenic sulphate aerosols (Ganor et al., 2000). Dust storms that pass over the Mediterranean Sea can contain internally mixed dust and sea salt (Levin et al., 2005).

There have been a handful of INP measurement campaigns in the Eastern Mediterranean region, specifically around Israel, and it is unsurprising given the importance of Saharan dust that several studies have either focussed on the effect of Saharan dust events or have drawn conclusions from them. Studies have shown that INP activity doubled during “dust days” compared to “clean days” (a relative term in the region), with south-westerly air masses from the Sahara Desert –contributing to the “dust days” while the “clean days” comprised north-westerly air masses from Europe (Levi and Rosenfeld, 1996) ~~performed ground level filter sampling of aerosols with INP analysis using a thermal diffusion chamber, finding that INP counts more than doubled during “dust days” compared to “clean days”, and further that a major fraction of the dust that they collected in rainwater samples originated from the Sahara Desert.~~

~~This conclusion was backed up by the more recent work of Ardon Dryer and Levin (Ardon-Dryer and Levin, 2014), who performed ground level (60 m a. m. s. l.) aerosol filter sampling with immersion mode INP analysis achieved using the FRIDGE-TAU (Frankfurt Ice nuclei Deposition freezing Experiment, the Tel Aviv University version) droplet assay. Throughout the campaign the researchers saw southwesterly (Saharan dust) or northwesterly (European) air masses, with the INP data again showing a doubling of INP activity during the “dust days” of the former case and the “clean days” of the latter. Aircraft analysis of winter cumulus clouds found that INP concentrations roughly approximated measured ice crystal concentrations within about one order of magnitude (Gagin, 1975).~~

Size-resolved INP analysis has demonstrated that ice-nucleating activity during dust events increased with increasing aerosol particle size and concentration, and that the activity of supermicron particles was similar for different dust events, suggesting that common mineral species were controlling ice nucleation (Reicher et al., 2018; Reicher et al., 2019). This latter hypothesis was similarly proposed during a study of INPs from three locations on the Saudi Arabian peninsula (Roesch et al., 2021), with the ice-nucleating activity being comparable to that of Saharan dust. However, a study of Saharan and non-Saharan transported dust samples from various locations showed that sieved and milled surface samples from Israel demonstrated some of the lowest ice-nucleating activities amongst all samples (Boose et al., 2016). Further INP analysis in Israel during ~~Further, analyses during~~ the annual Lag BaOmer bonfire festival event showed that, despite an increase in total aerosol concentrations there was no associated increase in INPs (Ardon-Dryer and Levin, 2014), which is consistent with studies during similar bonfire ~~and firework~~ events in the United Kingdom (Adams et al., 2020).

A number of campaigns have taken place in Cyprus in the Eastern Mediterranean, northwest of Israel. Ground-based filter samples collected from air masses passing over land and oceans sectors found no significant differences between their INP concentrations at temperatures lower than -15°C , implying that the INP population was dominated by long range transported aerosol (Gong et al., 2019). However, a few samples showed elevated activity at temperatures greater than -15°C , which suggested a population of biological INPs speculated to originate

100 in Cyprus. As a result, the researchers proposed that the standard methods of parameterising ice-nucleating activity (e.g. for desert dusts), such as the ice-active surface site density ($n_s(T)$), are unsuitable if the aerosol particle composition is unknown. During the INUIT-BACCHUS-ACTRIS campaign, vertical profiles over Cyprus using unmanned aircraft systems (UASs) suggested that ice-nucleating activity was dominated by Saharan dust in the lower and middle troposphere, with concentrations of an order of magnitude higher than at ground level which may have comprised weak marine and terrestrial sources in Cyprus (Schrod et al., 2017). This data was later used to validate a new method for vertical profiling of INPs (Marinou et al., 2019) using ground-based and spaceborne lidar observations from the campaign combined with INP parameterisations (DeMott et al., 2010; DeMott et al., 2015).

Several ship campaigns have taken place in the seas around Israel.

Boose et al. (Boose et al., 2016) performed INP analysis of a series of Saharan and non-Saharan transported airborne dust samples from a number of locations, and surface samples from other locations, including sieved and milled surface samples from Israel. Surprisingly, the authors found that the surface collected Israel dust demonstrated some of the lowest ice-nucleating activity amongst all of the other samples.

Reicher et al. (Reicher et al., 2018; Reicher et al., 2019) studied the ice-nucleating ability of size-resolved aerosol collected during dust events in Israel, analysed using the Weizmann Supercooled Droplets Observation on a Microarray (WISDOM) microfluidic instrument. Their results showed that ice-nucleating ability increased with particle size and concentration. The activity of supermicron particles was similar for different dust events, suggesting common mineral species controlling the nucleation.

Gagin (1975) performed aircraft and ground-based filter sampling of aerosols with INP analysis via a thermal diffusion ice nucleus counter to study winter cumulus clouds over Israel, finding that INP concentrations roughly approximated the measured ice crystal concentrations within about one order of magnitude.

Campaigns have taken place in Cyprus in the Eastern Mediterranean, northwest of Israel. Schrod et al. (2017), during the INUIT-BACCHUS-ACTRIS campaign, studied vertical profiles over Cyprus using unmanned aircraft systems (UASs) with electrostatic precipitation-based sampling to collect INPs for analysis by FRIDGE. Ice-nucleating activity was dominated by Saharan dust in the lower and middle troposphere, with concentrations of an order of magnitude higher than at ground level that may have comprised weak marine and terrestrial sources in Cyprus.

As part of the same INUIT-BACCHUS-ACTRIS campaign, Marinou et al. (2019) used ground-based and spaceborne lidar observations combined with INP parameterisations (DeMott et al., 2010; DeMott et al., 2015) to develop a new method for the vertical profiling of INPs. Validation was achieved using the vertical profile data of Schrod et al. (2017) obtained using UASs and FRIDGE.

(Gong et al., 2019) undertook a ground-based filter sampling campaign in Cyprus with INP analysis achieved using the Leipzig Ice Nucleation Array (LINA) and Ice Nucleation Spectrometer of the Karlsruhe Institute of Technology (INSEKT) instruments. Air masses were divided into “land” and “ocean” sectors, with no significant

differences found between their INP concentrations at temperatures lower than $-15\text{ }^{\circ}\text{C}$, implying that this INP population was dominated by long range transported aerosol. However, a few samples showed elevated activity at temperatures greater than $-15\text{ }^{\circ}\text{C}$, which suggested a population of biological INPs speculated to originate in Cyprus. As a result, the authors proposed that the standard methods of parameterising the INP activities (e.g. for desert dusts), such as the ice active surface site density ($n_s(T)$), are unsuitable if the aerosol particle composition is unknown.

During a ship cruise in the INPs measured in sea-spray aerosol (SSA) and sea surface microlayer (SML) in the western and central regions of the Mediterranean Sea showed lower than expected concentrations and was presumed to be due to the oligotrophic nature of the Mediterranean Sea (Trueblood et al., 2021). Interestingly, a performed sampled sea spray aerosol (SSA) and the sea surface microlayer (SML) marine and analysed their INP content using a Dynamic Filter Processing Chamber (DFPC). SML results showed lower INP concentrations than expected from the literature and was presumed to be due to the oligotrophic nature of the Mediterranean Sea. A dust wet deposition event demonstrated that, while INP concentrations in the SML increased by an order of magnitude almost immediately, the concentrations in the SSA took 3 days to increase. A

(Prodi et al., 1983) cruise performed aerosol sampling and INP analysis via a static diffusion chamber at $-16\text{ }^{\circ}\text{C}$ during a ship campaign in 1979 that crossed the Mediterranean, Red, and Arabian Seas, showed that the highest INP concentrations were measured in the southern section of the Red Sea (Prodi et al., 1983), and the authors noted a correlating correlation with the mass concentration of mineral aerosol apart from then SSA production was high. However, we note several other instances in that their dataset where the INP data increased much higher, relatively speaking, than the mineral content did, suggesting that other non-mineral INPs, e.g. biological INPs, may have been present.

undertook a more recent cruise around the Arabian Peninsula included INP measurements in the in which they measured INP concentrations in aerosol and seawater using the Scripps Institution of Oceanography automated ice spectrometer (SIO-AIS) droplet assay. Several days' of data were collected in the Red Sea (Beall et al., 2022), with air masses from the Mediterranean, the Nile Delta and the Sinai Peninsula that each contained some degree of dust loading. Following heating and peroxide treatment of the samples, the with the authors researchers noting that agricultural soil dust from the fertile Nile Delta region exhibited greater heat sensitivity that is indicative of biological INP content. Thus, the authors suggested that INP activity at temperatures below around $-15\text{ }^{\circ}\text{C}$, while the activity at warmer temperatures was likely due to fertile soils from the agriculturally important Nile Delta region, which comprises part of the Fertile Crescent region that arcs from the Nile Delta through Israel, Lebanon, and northern Syria and southern Turkey, then down through northern and eastern Iraq and western Iran. As a consequence of this and in noting that aerosol aging can have an effect on INP activity, the researchers, similar to Gong et al. (2019), proposed that mineral dust parameterisations alone may not be suitable for the representation of regional ice-nucleating INP activity when the INP-INP population contains unknown INP populations that are not is not dominated by the mineral dust content.

~~Roesch et al. (2021) studied cloud condensation nuclei (CCN) and INPs from three locations on the Saudi Arabian peninsula, finding that surface samples from the east coast were ice nucleation active at warmer temperatures than those in the northeastern and central regions of the country, highlighting that INPs from a relatively small region could have very different properties. The authors also found that INP activity was similar to that of Saharan dust.~~

170 To better understand INP concentrations and activity in the Eastern Mediterranean region beyond dust events, we undertook an intensive two-week field campaign in Rehovot, Israel, in October-November 2018 where we collected ground-based filter samples for INP analysis by immersion mode droplet freezing assays. Given the variety of air masses received by this location, we hoped to determine whether some sources could potentially influence local INP populations over the typical dust events.

175 During this campaign we experienced four major air masses that we were able to study: (i) a southwesterly dust event from the Sahara Desert that passed over the fertile Nile Delta region, (ii) a northwesterly air mass from Europe, (iii) a prolonged easterly event from the Arabian Desert that contained air with both high- and low-loadings of dust, and (iv) an air mass that passed over the agriculturally important northern Fertile Crescent region before sweeping eastward to the sampling site.

180 2 Experimental

2.1 Aerosol sampling and preparation of particle suspensions

185 Atmospheric aerosol samples were collected over 10 days, from 25/10/18 to 03/11/18 (DD/MM/YY), from the roof of the Department of Earth and Planetary Sciences building at the Weizmann Institute of Science in Rehovot, Israel (~~31°54'26.1"N, 34°48'37.7"E~~~~31.9° N, 34.8° E~~, ~80 m a. m. s. l.). Aerosol sampling was performed using two different methodologies: (1) the collection of aerosol particles onto filters, and (2) the sampling of aerosol particles into water using an impinger. Aqueous suspensions of particles were prepared in both instances using purified water (18.2 MΩ cm at 25 °C, 0.22 μm filtered) ~~that was collected every morning~~ from a Thermo Scientific™ Barnstead™ GenPure™ water purification system ~~and stored in 50 mL polypropylene centrifuge tubes (Starstedt Ltd.).~~

190 Filter-based sampling for INP analysis was performed using our previously described technique (O'Sullivan et al., 2018). Two BGI PQ100 Air Sampling Systems (Mesa Laboratories, Inc., Lakewood, CO, USA) with PM₁₀ (particulate matter of ≤10 μm diameter) inlet heads were purchased from Air Monitors (Tewkesbury, UK) and were employed in order to draw air through porous filters onto which aerosol particles were adsorbed. The BGI PQ100 is used as an Environmental Protection Agency (EPA) Reference Method for PM₁₀ (designation no. RFPS-1298-124). PM₁₀ aerosol were sampled onto Whatman® Nuclepore™ track-etched membrane polycarbonate filters (1.0 μm pore size, 47 mm diameter filters, purchased from Sigma-Aldrich, Dorset, UK) at a flow rate of 16.67 L min⁻¹ (1 m³ h⁻¹).
195 While the filters had 1.0 μm pore sizes, studies have demonstrated that membrane filters are able to capture particles much smaller than the given pore size (Soo et al., 2016; Lindsley, 2016); 1.0 μm pore size polycarbonate filters have

been shown to have collection efficiencies of 50-90 % for particle sizes of 0.01-0.2 μm , and 90-100 % for particle sizes larger than 0.2 μm (Burton et al., 2006; Lindsley, 2016).

200 Filter-based sampling for INP analysis was performed using two main strategies: (i) the collection of samples for 3 hours every morning and 3 hours every afternoon (yielding 3,000 L of sampled air for each sample), and (ii) the collection of samples for 24 h from midday to midday (24,000 L of sampled air). The 3 h samples were collected using one PQ100 sampler and the 24 h samples using a second PQ100 sampler. Collection of the 3 h samples was intended to provide a relatively short time resolution of the INP measurements, while the 24 h samples were originally
205 collected to achieve a large volume of sampled air since it was not immediately clear how much air would need to be sampled to obtain an INP signal. However, the sampler used for the 24 h samples failed during collection of the fourth sample between 28th-29th October after 17.5 h, and no further 24 h samples were collected from that point.

In one experiment during the afternoon of 29th October 2018, the two samplers were run side-by-side for 3 h, with one collecting PM_{10} aerosol onto a filter as usual and the other collecting PM_1 aerosol simultaneously onto a
210 second filter via the use of a PM_1 adaptor (SCC 2.229 Cyclone, Mesa Labs) inserted into the sampler. Two handling blanks were also performed for the filter-based sampling by attaching a HEPA (high-efficiency particulate air) filter to the BGI PQ100 sampler and pulling filtered air through a track-etched membrane filter. One handling blank filter was sampled for 1 h during the early stages of the campaign, and another filter sampled for 3 h during the latter stages.

Details of the sampling times and volumes of air sampled for each filter collected are provided in Table S1
215 in the Supplementary Information (SI). Filters collected for INP analysis were removed from the BGI PQ100 sampling system and immediately inserted into a 50 mL centrifuge tube (Sarstedt Ltd., Leicester, UK) using tweezers, then 4 mL of purified water added via pipette. The tube was shaken vigorously for several seconds before being vortexed on a vortex mixer (Labnet VX100) for 5 min, allowing the adsorbed aerosol particles to be washed off the filter and into suspension ready for analysis via a droplet freezing assay.

220 Impinger samples were collected during the afternoon for 7 days of the campaign using a Bertin Technologies Coriolis® Micro air sampler, purchased from Air Monitors. This impinger measures particles larger than 0.5 μm , with a collection efficiency, D_{50} (i.e. the particle size at which the collection efficiency is 50 %), of $<0.5 \mu\text{m}$ (information; Carvalho et al., 2008). Impingers have previously been successfully employed by several research groups for INP sampling and analysis, (Šantl-Temkiv et al., 2017; Garcia et al., 2012) including the use of a Coriolis
225 Micro for the detection of ice-nucleating proteins by genetic sequencing (Els et al., 2019). Here, ambient air was aspirated into a cone filled with 10 mL of purified water, with aerosol particles that entered the sampler inlet being centrifuged to the wall of the cone and extracted from the air into the water. Sampling was performed in multiple 10 min intervals (the maximum runtime of the sampler), at a flow rate of either 100 L min^{-1} or 300 L min^{-1} , until 6,000 L of air had been sampled into the water-filled cone. A lid was screwed onto the cone once it had been removed from
230 the sampler to prevent contamination. Purified water was added to the cone following each sampling interval due to

evaporational losses during sampling, and the final volume of water in the cone (~5-10 mL) was determined by weight at the end of the collection period.

235 A handling blank was performed for the impinger samples by filling a sampling cone with purified water, attaching it to the impinger, and then removing it and screwing a lid onto it ready for droplet freezing assay analysis of the water. It was not possible at the time to pull HEPA-filtered air through the sampler for the handling blank due to the awkward shape of the inlet. Details of the sampling times and final water volumes are provided in Table S1 in the ESI. The water from the cone was analysed directly via a droplet freezing assay.

2.2 Heat treatment of aerosol suspensions

240 An aliquot of every aqueous suspension from the filter samples and the impinger samples was subjected to a heat treatment test. Heat treatments are used to test for the presence of heat-labile ice-nucleating materials (O'Sullivan et al., 2018; Christner et al., 2008b; Christner et al., 2008a; Daily et al., 2022), which are generally equated to ice-nucleating proteins that suggest the presence of biogenic INPs (e.g. bacteria or fungi). Such tests can only be used to indicate of the possibility of biogenic INPs being present but cannot alone provide definitive evidence. Since some minerals can also lose their activity upon heating, the assumption that a loss of activity may be due to ice-nucleating
245 proteins is considered to only be viable if the main mineral dust component is microcline K-feldspar, which does not lose activity following heat treatment (Daily et al., 2022).

The aerosol compositions of dust events in Israel mainly compromise calcite (59 ± 15 %), quartz (23 ± 7 %), dolomite (11 ± 8 %), feldspars (5 ± 3 %), halite (2 ± 1 %) and traces of clay minerals (Ganor and Mamane, 1982; Foner and Ganor, 1992; Ganor, 1975). While the carbonates (calcite and dolomite) and quartz dominate the
250 composition, it is the feldspars that are thought to be the most important factor for the nucleation of ice in mixed-phase clouds (Atkinson et al., 2013; Kiselev et al., 2017). While different feldspars have vastly different ice-nucleating activities (Harrison et al., 2016), microcline potassium (K)-rich feldspar exhibits much greater activity than quartz or Na-Ca feldspars (plagioclase series) (Harrison et al., 2019).

As a consequence, the presence of K-feldspar, even at concentrations of around only 1 % (Harrison et al.,
255 2022), is expected to dominate the ice-nucleating ability of airborne desert dust. Boose et al. (2016) analysed sieved and milled surface dust from Israel and found K-feldspar contents of 1.7 ± 0.3 % and 1.4 ± 0.3 %, respectively, and agreed with Ganor et al. (Ganor and Mamane, 1982; Foner and Ganor, 1992; Ganor, 1975) that the samples comprised mostly of carbonates. Based on this, we assume here that microcline K-feldspar is the dominant component of the mineral dust in Israel in terms of ice-nucleating activity, and so the heat test for biogenic INPs remains viable.

260 Heat treatment was performed by adding 1 mL of aqueous suspension to a sealed glass vial and heating at 95 °C for 30 min in a bath of boiling water, consistent with the recommendations of Daily et al. (2022). The sample was then allowed to cool to room temperature prior to analysis via a droplet freezing assay for comparison to the untreated

sample. A control test was also performed by performing droplet freezing assays on an aliquot of purified water before and after being subjected to heat treatment.

265 2.3 Droplet freezing assays for INP analysis using the $\mu\text{L-NIPI}$

All aqueous suspensions from the filter-based samples (3 h and 24 h collection periods) and the impinger-based samples were analysed using the Microlitre Nucleation by Immersed Particle Instrument ($\mu\text{L-NIPI}$) (Whale et al., 2015) immediately following their collection and preparation. This involved setting the droplet freezing apparatus up in the laboratories of the Weizmann Institute of Science. The advantage of analysing the samples immediately following collection was that we reduced the possibility of changes in the INP population that have been observed on storage and transport of filter samples (Beall et al., 2020).

Hydrophobic glass slides (22 mm \O \times 0.22 mm siliconised glass, Hampton Research, Aliso Viejo, CA, USA) were washed with ethanol and purified water, dried with dry nitrogen gas, and placed atop the cold plate of a Stirling engine-based cold stage (EF600, Grant Asymptote Ltd., Cambridge, UK).

275 Sample tubes/cones of aqueous suspension were shaken vigorously and vortexed on a vortex mixer for 5 min, then \sim 40 droplets (1 μL volume each) were pipetted onto the hydrophobic glass slide. A Perspex shield was placed over the glass slide to form a chamber, which was purged with dry N_2 gas to prevent the formation of condensation on the cold stage. The cold stage was cooled at a rate of 1 $^\circ\text{C min}^{-1}$ to -40°C and videos were recorded via a webcam as the array of droplets froze, with the droplet freezing temperatures being determined by synchronisation of the videos with the cold plate temperature measurements. The temperature error of the $\mu\text{L-NIPI}$ was $\pm 0.4^\circ\text{C}$ (Whale et al., 2015).

2.4 Processing of droplet freezing assay data

Droplet freezing assay analysis of the aqueous suspensions of collected ambient aerosol allowed the fraction of the droplets frozen as a function of temperature, $f_{\text{ice}}(T)$, to be determined using Equation (1) (Vali, 1971, 1994):

285

$$f_{\text{ice}}(T) = \frac{n_{\text{ice}}(T)}{n_{\text{tot}}}, \quad (1)$$

where $n_{\text{ice}}(T)$ is the number of droplets frozen at temperature T , and n_{tot} is the total number of droplets. The fraction frozen values represent the raw data for the droplet freezing assays, and these are shown for all of the campaign results in Figures S1-S7 in the SI.

290

Early in the campaign (from 25/10/18 to 30/10/18 (DD/MM/YY)), the bulk of the droplets in the purified water blanks froze in the region of around -23°C to -29°C . However, on 31/10/18, a change occurred in the blanks, and from that point until the end of the campaign the majority of droplets froze from around to -21°C to -26°C ,

thus exhibiting a noticeable increase in the background signal. The cause of this change in the quality of the blanks
295 could not be identified, despite several possibilities being tested. To compare sample data against the appropriate set
of blanks, the blanks were separated into two categories: (i) “early” blanks analysed between 25/10/18 and 30/10/18,
and (ii) the poorer “later” blanks analysed from 31/10/18 to 04/11/18. The fraction frozen data for these two sets of
blanks are shown in Figure S1 in the SI.

Handling blanks were performed in order to assess the extent of contamination that could occur as part of the
300 handling of the samples, using the methods described earlier for the filter and impinger samples. The results are
shown in Figure S2 in the SI and show little difference between the handling blanks and the purified water,
demonstrating that there was little, if any, contamination introduced into the samples as part of the handling
procedure.

Controls for the heat test were also performed by heating purified water and testing it before and after the
305 treatment. Two such tests were performed: one using an “early” blank and one using a “later” blank, and the results
are shown in Figure S3 in the SI. The control test using water from the “early” part of the campaign demonstrated a
small increase in ice-nucleating activity following the heat treatment. However, the control test for the “later” blank
demonstrated that the heated water had a similar activity to the unheated water, and the heated “early” water showed
similar results to both the heated and unheated “later” water. This suggests very clean blanks (e.g. the “early”
310 untreated water here) are prone to a small amount of contamination upon heating, whereas blanks already containing
some notable contamination are unaffected by further heating. Given this, and the fact that no decrease in activity of
the purified water blanks was observed following heating, the heat test was determined to be viable for the ambient
aerosol samples.

Fraction frozen curves were generated for the aerosol samples, and can be found in the SI: results for the 3h
315 filter samples are shown in SI Figure S4, the 24 h filter samples are shown in SI Figure S5, the impinger samples are
shown in SI Figure S6, and the heat tests for all of the samples are shown alongside the unheated results in SI Figure
S7. The plots also show the blanks, and the data for both the blanks and the samples are separated into the “early”
and “later” categories so that the samples can be compared against their appropriate blanks.

2.5 Background subtraction of INP data

320 The data shown in the fraction frozen curves were used to perform background subtraction of the blanks from the
sample data in order to remove the influence of impurities from the INP signal. This procedure is described in detail
by Vali (2019) and Sanchez-Marroquin et al. (2021). The first step was to bin the data from all of the individual
droplet freezing assays, both the blanks and samples, into 0.5 °C temperature intervals, and then calculate the
differential freezing nucleus spectrum, $k(T)$ ($\text{cm}^{-3} \text{ } ^\circ\text{C}^{-1}$), using Equation (2) (Vali, 2019; Vali, 1971):

325

$$k(T) = -\frac{1}{v \cdot \Delta T} \cdot \ln\left(1 - \frac{\Delta N}{N(T)}\right), \quad (2)$$

where v is the droplet volume (1 μL , i.e. 0.001 cm^3), ΔT is the temperature interval (0.5 $^\circ\text{C}$), N is the number of unfrozen droplets in the temperature interval, and ΔN is the number of droplets frozen in the temperature interval.

330 The $k(T)$ values for the background data (i.e. the purified water blanks) were obtained by combining all of the droplet freezing temperatures within the respective “early blanks” and “later blanks” populations. The average and standard deviation of the $k(T)$ values in each temperature bin were determined and these are shown in Figure S8 in the SI.

335 These average blank $k(T)$ values for the “early” and “later” populations were subtracted from the sample $k(T)$ values for each temperature bin, as shown in Figure S9 in the SI. Uncertainty values were determined by calculating the randomness of active site distribution within the droplet population during the droplet freezing experiments via Monte Carlo simulations (Harrison et al., 2019), the Poisson uncertainties in the data, and the standard deviation of the background $k(T)$ values, with these uncertainties being combined in quadrature to give the final error values for the sample data. Following the background subtraction of the data in terms of $k(T)$, the final values were converted
340 into the cumulative integrated ice-active site volume density, $K(T)$, a singular approximation, by summing the background-subtracted sample $k(T)$ values for temperatures warmer than T , as per Equation (3):

$$K(T) = \sum_0^T k(T) \cdot \Delta T, \quad (3)$$

345 The singular approximation assumes that ice nucleation is temperature-dependent and time-independent, that each droplet freezes due to a single nucleation event, droplet freezing occurs at a characteristic temperature depending on the nature of the INPs, and that each droplet contains the same average surface area of ice-nucleating particles. The nucleation of many materials is actually both site- and time-dependent (Knopf et al., 2020; Vali, 2008), but for simplicity we make the assumption here that time-dependence is second order (Vali, 2008; Holden et al., 2019;
350 Holden et al., 2021) and so apply the singular approximation.

The $K(T)$ values for the samples calculated using Equation (3), for both the “early” and “later” populations, are shown in Figure S10 in the SI. From the background-subtracted $K(T)$ values, the INP concentrations, N_{INP} (L^{-1} of sampled air), were calculated using Equation (4):

$$355 \quad N_{\text{INP}} = K(T) \cdot \frac{V_{\text{wash}}}{V_{\text{air}}}, \quad (4)$$

where V_{wash} is the volume of water used to wash the particles off a collection filter and into suspension or the volume of water in the cone of the impinger (see Table S1), and V_{air} is the volume of sampled air onto the filter or into the

impinger cone (see Table S1). INP concentrations for both the “early” and “later” samples populations for all samples
360 are shown in Figure S11 in the SI.

The ice-nucleating activity of the aerosol throughout the campaign was quantified via calculation of the ice-
active site density per surface area, $n_s(T)$ (cm^{-2}), from N_{INP} using Equation (5):

$$n_s(T) = \frac{N_{\text{INP}}}{dS}, \quad (5)$$

365

where dS is the surface area concentration of particles in the sampled air ($\text{cm}^2 \text{L}^{-1}$), assuming that the total surface
area was equal to the surface area of mineral dust. dS was calculated for each filter and impinger sampling time based
on the OPC and SMPS data ([described below](#)) and is provided for each sample in Table S32 in the SI in terms of μm^2
 cm^{-3} .

370 2.6 Meteorological data

Local meteorological data was obtained from the Israel Meteorological Service (IMS; <https://ims.gov.il/en>) station #44, Bet Dagan: 32°00'26.3"N-, 34°48'49.6"E, 31 m elevation, -approximately 11 km north of the campaign site. This data includes temperature, atmospheric pressure, rainfall, wind direction, wind speed, and relative humidity.

375 2.6.7 Aerosol monitoring

PM₁₀ mass concentrations were obtained from the Israeli Ministry of Environment (IME; <https://air.sviva.gov.il/>), and were measured in the Rehovot station (station #56, 31°53'N 34°48'E) located ~1 km from the sampling site, and the concentrations for each sampling period are provided in Table S2 in the SI. The 5 min mean data was used, alongside air mass back trajectories and wind directions, to help categorise the air masses each day, as described in Section 3.1 and by Gat et al. (2021).

380

Aerosol sizes and concentrations were also monitored during the campaign using several instruments. An optical particle counter (OPC; GRIMM Technologies Model 1.109) monitoring in the range of 0.25 – 32 μm was used continuously throughout the campaign and provided data every 6 s. A Scanning Mobility Particle Sizer (SMPS) spectrometer (Model 3938, TSI Inc.) was used to take measurements in the range of 14.1 – 710.5 nm on a daily basis
385 during the filter and impinger sampling periods.

Since the SMPS was not run continuously throughout the campaign, illustrated time series data for comparison with meteorological conditions shows only the OPC measurements. However, for processing of aerosol sample data, ~~The OPC data was used to determine particle concentration (dN), and PM₁₀, PM_{2.5} and PM₄ concentrations. The~~the OPC and SMPS data were combined to determine the particle concentration (dN), PM₁₀ mass

390 concentrations, and the surface area concentrations, dS , throughout the campaign in the size range of 14.1 nm to 10
 μm (due to the use of a PM_{10} filter sampler). PM_{10} , $\text{PM}_{2.5}$ and PM_1 mass concentrations were calculated using a
particle density, ρ , value of 1.0 g cm^{-3} from the OPC data as this provided the best comparison with the IME station
data. However, given that mineral dust has a density of 2.65 g cm^{-3} , the PM_{10} mass concentrations were also
395 were converted from the mobility-equivalent particle diameter, d_{me} , to the volume-equivalent sphere diameter, d_p ,
using Equation (6), as per Möhler et al. (2008):

$$d_p = \frac{1}{\chi} d_{\text{me}}, \quad \text{Equation (6)}$$

400 where χ is the dynamic shape factor that accounts for the shapes of non-spherical particles. Möhler et al. (2008)
described how secondary organic aerosols (SOA) had a χ of 1.0, while dry minerals and dusts could have a χ of 1.3,
with mixtures of minerals/dusts and SOA having χ values of 1.1–1.2. The optical particle diameter measurements
from the OPC were assumed to be equivalent to the volume-equivalent diameter already, as per Peters et al. (2006),
and so were used as is.

405 A χ value of 1.1 was used when handling the SMPS data from this campaign as it offered the best agreement
between the OPC and SMPS data in $dN/d\log D_p$ and $dS/d\log D_p$, while the uncertainties when processing the data
included the effect of varying the χ value between 1.0 and 1.2 to account for variability in the types of the aerosols
and since the $dN/d\log D_p$ and $dS/d\log D_p$ occasionally matched well between the SMPS and OPC when using these
values. Following the conversion of the SMPS data to d_p , the overlap in the SMPS and OPC particle size ranges was
410 addressed by restricting the SMPS size range to 0.0128–0.315 μm (D_p) and the OPC size range to 0.325–10 μm for
 PM_{10} as this offered the best agreement in $dN/d\log D_p$ and $dS/d\log D_p$.

2.87 Size-resolved biological analysis

Size-segregated ambient dust particles were analysed for their bacterial and fungal content as described by Gat et al.
415 (2021). Aerosol particles were collected onto Cyclopore polycarbonate filters every day of the campaign using a
micro-orifice uniform deposit impactor (MOUDI; MSP Corporation model 110-R) (Marple et al., 1991), which
operated at 30 L min^{-1} for 8 h, similarly to Huffman et al. (2013) and Mason et al. (2015). The collection efficiencies
for the stages of this MOUDI are provided in Reicher et al. (2019). DNA was extracted from the filters using the
DNeasy® PowerWater® kit (Qiagen) with slight modifications to the manufacturer's protocol: elution was divided
420 into two steps, adding 50 μL of elution buffer (EB) solution and centrifuging at 13,000 g for 1 min each time. The
filters were cut in half, and half of each filter was stored as a backup at $-20 \text{ }^\circ\text{C}$.

425 ~~The amount of DNA that was collected on each filter was very small, nearly below the minimal threshold that can allow amplification and detection by polymerase chain reaction (PCR). Therefore, two filter halves from sequential MOUDI sampler stages were combined in each PowerWater® bead-tube, which enabled the extraction of a sufficient amount of DNA, as well as keeping an identical back-up. Two halves from two sequential MOUDI sampler stages were combined in each PowerWater® bead-tube.~~ Stages 2 ($D_{50} = 5.6 \mu\text{m}$) and 3 ($D_{50} = 3.2 \mu\text{m}$) of the MOUDI were combined to give the “coarse” particle size range ($3 \mu\text{m} - 18 \mu\text{m}$), stages 4 ($D_{50} = 1.8 \mu\text{m}$) and 5 ($D_{50} = 1.0 \mu\text{m}$) were combined to give the “intermediate” particle size range ($0.3 \mu\text{m} - 3 \mu\text{m}$), and stage 6 ($D_{50} = 0.6 \mu\text{m}$) and 7 ($D_{50} = 0.3 \mu\text{m}$) refer to the “fine” particle size range ($<1.3 \mu\text{m}$). After extraction, the DNA samples were stored at $-20 \text{ }^\circ\text{C}$ for further analysis. Blank filters were extracted similarly.

430 Total bacterial and fungal concentrations were determined by quantitative polymerase chain reaction (qPCR), using primers listed in Table S43 in the ESI. Bacterial DNA was analysed via amplicon sequencing of the 16S rRNA gene, while fungal activity was analysed using the ribosomal internal transcribed spacer (ITS) region. qPCR reactions were performed as follows: $10 \mu\text{L}$ of SensiFAST® SYBR mix (Bioline), $2 \mu\text{L}$ each of forward and reverse primers (initial concentration of 10 mM each, with a final concentration of 1 mM each) ~~mM of each primer~~, $2 \mu\text{L}$ of extracted DNA (template), and $4 \mu\text{L}$ of PCR-grade deionised water (DI) to ~~reach~~ a total reaction volume of $20 \mu\text{L}$, with triplicates for each DNA sample and a negative control (i.e. a blank filter). qPCR was conducted separately for ITS and 16S. Amplification was performed in 96-well plates using a StepOnePlus® Real-Time PCR instrument (Thermo Fisher Scientific), as follows: 5 min at $95 \text{ }^\circ\text{C}$, followed by 40 cycles of 5 s at $95 \text{ }^\circ\text{C}$, and 30 s at $60 \text{ }^\circ\text{C}$ or $52 \text{ }^\circ\text{C}$ for 440 16S or ITS annealing temperatures, respectively. A melting curve stage at $60\text{--}95 \text{ }^\circ\text{C}$ was included, confirming substrate specific amplification. Standard curves were constructed using known concentrations of plasmids containing the relevant genes along with a non-template control.

2.8 Meteorological data

445 ~~Local meteorological data was obtained from the Israel Meteorological Service (IMS, <https://ims.gov.il/en>) station #42, Bet Dagan: 32.0073 latitude, 34.8138 longitude, 31 m elevation, approximately 14 km north of the campaign site. This data includes PM_{10} concentrations, temperature, atmospheric pressure, rainfall, wind direction, wind speed, and relative humidity.~~

2.9 Air mass back trajectories

450 The National Oceanic and Atmospheric Administration (NOAA) Air Resources Laboratory’s HYSPLIT (Hybrid Single Particle Lagrangian Integrated Trajectory) model (<https://www.arl.noaa.gov/hysplit/>) (Stein et al., 2016; Fleming et al., 2012; Draxler and Hess, 1998) was used to generate 72 h back trajectories of the movements of air masses to the sampling region during the field campaign. Back trajectories were obtained for $10:00$ local time ($8:00$ UTC) and $18:00$ local time ($16:00$ UTC) every day of the campaign.

The 72 h duration of the back trajectories was chosen as a consensus in order to assess the long range transport of air masses while limiting the trajectory error (Kahl and Samson, 1986; Karaca et al., 2009; Sari et al., 2020). The trajectories were calculated for three arrival heights (500 m, 1,500 m, and 2,500 m) because mineral dust is transported below this height 3 km in this season during the season in which the campaign took place (Gobbi et al., 2004). In most cases, all three trajectories travelled in the same path and therefore air mass classifications (see Table 1) were based on the three of them. However, even if only one trajectory originated in a dust source area, and high PM_{10} concentrations (from the IME site) were measured, this would be defined as the dust origin.

3 Results and Discussion

3.1 Ice-nucleating particle concentrations

Of the different types of sampling performed (3 h filters, 24 h filters, 20 min impinger samples), the 3 h filter samples provide a more complete and consistent time series for the campaign; hence they are the main results presented here. The final INP concentrations for the 3 h filter samples, collected every morning and afternoon throughout the campaign, are presented in Figure 1a, and demonstrate a range of variability in INP concentrations during the two-week campaign, from $\sim 0.2 - 14$ INP L^{-1} of air in the temperature range of -10 to -25 °C. Concentrations were sufficiently high, and even with the relatively short sampling times and volumes employed here they were largely above the baseline of the cold stage instrument, enough to be well above the lowest freezing temperature of the instrument (see Figure S9 in the SI), varying between $\sim 0.1 - 2$ INP L^{-1} at -20 °C.

Some INP concentrations also exhibited a “hump” of activity at the warmer range of freezing temperatures (in particular the “181029 afternoon” sample), which could be indicative of biological INP activity (Schnell, 1977; Conen et al., 2011; O’Sullivan et al., 2014; O’Sullivan et al., 2018), i.e. a potential “biological hump” in the N_{INP} data. This phenomenon is discussed in further detail later in the manuscript.

The 24 h filter data, shown in Figures S11a and S12 in the SI, shows consistent INP concentrations that reside within the median of the 3 h filter data. This is likely a consequence of averaging the INP data over such a long period of time during changes in N_{INP} between high and low values, for example if there were rapid changes in N_{INP} with diurnal cycles or changes in air masses. This highlights the need for shorter time resolution measurements.

The N_{INP} values from the Coriolis® impinger, measuring particles > 0.5 μm , are shown in Figures S11 and S13 in the SI, were collected on time scale of 20 min (60 min in one case) and showed a large degree of variability depending on the air mass, and this will be described further when discussing the time series of the campaign. Further, as for the 3 h filter results, the “181029 afternoon” exhibited a notable potential “biological hump” on N_{INP} at the warmer temperatures, which will be explored alongside the 3 h filter data later in the manuscript.

Figure 1b illustrates the 3 h filter data shown in Figure 1a but now separated into two populations: “dust events” and “clean days”. The categorisation of these populations follows that is prescribed by Gat et al. (2021), who


performed an analysis of airborne bacterial and fungal communities during the same campaign. Air masses were categorised based on the back trajectories for a given day, as assigned by Gat et al. (2021) previously, and these are shown in Table 1 (see Table S2 in the SI): southwesterly = SW, northwesterly = NW, and easterly = E. Days during which the PM₁₀ concentrations (using data from the IME site) for the entire day were greater than 44 µg m⁻³ were classified as dust events as per Krasnov et al. (2016), and were designated with a lower case “d”. Thus, INP measurements taken on days designated as dust events are shown as brown lines in Figure 1b, while those on clean days are shown as blue lines. However, it must be noted that the region is typically dusty regardless, hence the term “clean days” is relative; most “clean days” would still be dusty, but below the 44 µg m⁻³ threshold. Also shown in Figure 1b are the boundaries of the global INP data compiled from precipitation sample data from between 1971-2014 by Petters and Wright (2015) to provide context for where this campaign data sits.

Table 1: Classification of air mass trajectory categories as per Gat et al. (2021), which was part of the same field campaign. Days with dust events are identified with a subscript “D”, and were identified as dust events if the PM₁₀ loading for the day exceeded 44 µg m⁻³ (Krasnov et al., 2016; Gat et al. 2021). Dates are in the DD/MM/YY format.

<u>Air mass trajectory</u>	<u>ID</u>	<u>Dates (DD/MM/YY)</u>
<u>Southwesterly with Saharan dust event</u>	<u>SW_D</u>	<u>25/10/18</u>
<u>Northwesterly</u>	<u>NW</u>	<u>26/10/18 – 27/10/18</u>
<u>Easterly with Syrian and/or Arabian dust event</u>	<u>E_D</u>	<u>28/10/18 – 31/10/18</u>
<u>Easterly</u>	<u>E</u>	<u>01/11/18 – 03/11/18</u>

3.2 Comparison to literature *N*_{INP} data

INP concentrations from this campaign are plotted against relevant literature data in Figure 1c. The results demonstrate that, with only a few exceptions at temperatures warmer than ~-19 °C, the dust days had consistently higher INP concentrations than the clean days. The data fits largely well within the boundaries of the global INP from Petters and Wright (2015), with the Eastern Mediterranean data sitting in the middle of the compiled data above ~-24 °C.



510 **Figure 1:** Plots showing ice-nucleating particle (INP) concentrations (N_{INP}) in Israel. (a) Background-subtracted INP concentration spectra for samples collected for 3 h onto filters from 25th October–4th November 2018. The temperature error is estimated to be 0.4 °C (Whale et al., 2015). Dates are given in the YYMMDD format. (b) The same data shown represented as being from a dust event (brown) or a clean day (blue), overlaid onto the INP concentration range of precipitation samples compiled by Petters and Wright (2015) shown in grey. (c) Comparison of INP concentration data from this study to literature data obtained in Israel and in nearby regions of the Eastern Mediterranean. Literature data are represented as discrete data points (Gagin, 1975; Levi and Rosenfeld, 1996; Schrod et al., 2017; Tarn et al., 2020) or as lines (Beall et al., 2022; Gong et al., 2019) as appropriate.

515

When compared to the INP concentrations measured by Gong et al. (2019) in Cyprus from both land and oceanic sources at an earlier time in the year (2-30 April 2017), our clean day data is consistent with much of the land sector data and much higher than the ocean sector data. However, data obtained in the Red Sea (5-7 July 2017), with air masses flowing over the Nile Delta and the Sinai Peninsula, by Beall et al. (2022) during a ship campaign, compares well with much of the dust event data from our campaign. Only a portion of the data of Schrod et al. (2017), taken at altitudes of 717-2,550 m a. s. l. over Cyprus using UAVs/UASs, is in the same temperature regime as our INP measurements, but in the ranges of $-17.5\text{ }^{\circ}\text{C}$ to $-22.5\text{ }^{\circ}\text{C}$ the data sets compare well despite the altitude difference, although the Schrod et al. (2017) data typically covers a wider INP concentration range.

Aircraft and ground-based measurements by Gagin (1975) in Israel showed INP concentrations similar to ours at $-20\text{ }^{\circ}\text{C}$, and while there is some comparison at $-15\text{ }^{\circ}\text{C}$ and $-12\text{ }^{\circ}\text{C}$, the concentrations tend to trend higher than all but our most active samples. Levi and Rosenfeld (1996) collected filter samples at ground level for INP analysis at $-15\text{ }^{\circ}\text{C}$, using a static thermal diffusion chamber, during dust events and clean days. Their results showed that INP concentrations more than doubled during dust storms.

The data from Tarn et al. (2020) was obtained using three of the same samples collected during the October-November 2018 campaign: a 3 h filter sample (“181031 afternoon”), a 24 h filter sample (“181025 to 181026 (24 h)”), and an impinger sample (“181103 afternoon (impinger)”). These samples were analysed using the Lab-on-a-Chip Nucleation by Immersed Particle Instrument (LOC-NIPI). The LOC-NIPI is a continuous flow cold stage instrument that analyses $100\text{ }\mu\text{m}$ diameter droplets ($\sim 520\text{ pL}$). Although this data does not overlap in temperature with the μL -NIPI data (using $1\text{ }\mu\text{L}$ droplet arrays), it extends the INP data to colder temperatures and is complementary to the 3h filter results shown here.

The “dust event” and “clean day” data of Ardon-Dryer and Levin (2014) from ground-based measurements is much higher in concentration compared to all other available literature data from the region, but showed similar trends to the those illustrated here. The INP parameterisation of DeMott et al. (2015) for mineral dust particles larger than $0.5\text{ }\mu\text{m}$ was calculated based on the OPC data and is shown in Figure 1c, but is much higher than the bulk of the INP data from the region.

3.3 Time series of the Eastern Mediterranean campaign


A time series of multiple data sets for the duration of the campaign in Rehovot, Israel, from 25th October to 3rd November 2018, is provided in Figure 2. Meteorological data obtained from the IMS station in Bet Dagan (11 km away) is shown in Figures 2a-e. Aerosol data measured using the GRIMM OPC is provided for particle concentrations (dN) (Figure 2f), $\text{PM}_{10/2.5/1}$ mass concentrations (Figure 2g), and particle size distributions in terms of D_p vs. $dN/d\log D_p$ (Figure 2h). The SMPS data is not represented here as the SMPS was not used continuously, unlike the OPC, though the SMPS data was used when calculating $n_s(T)$. The raw particle count data for both the OPC and

550 ~~SMPS are illustrated in Figures S14 and S15 in the SI, respectively. Raw data from the GRIMM OPC and the SMPS~~
~~are provided in Figures S14 and S15 in the SI, respectively.~~ A higher resolution version of the D_p vs. $dN/d\log D_p$ plot
comprising both the OPC and SMPS data, and for the full size range up to $32 \mu\text{m}$, is shown in Figure S16a in the SI
~~I, while T~~ the surface area distribution (dS) in terms of D_p vs. $dS/d\log D_p$ ~~for the OPC and SMPS data, for the full size~~
555 ~~range, is also~~ shown in Figure S16b in the SI. Figure 2g shows the PM_{10} data from both the OPC and the IME station,
and shows good agreement when a particle density of 1 g cm^{-3} was assumed when processing the OPC data. However,
since mineral dust can have a density of 2.65 g cm^{-3} , the OPC data was also processed using this density value to
demonstrate the upper limit of PM_{10} mass concentration, highlighting an approximately 2-3 \times increase compared to
the values processed using 1 g cm^{-3} during periods when more coarse particles were measured in higher
concentrations (see Figure 2h).

560 INP concentrations at selected temperatures ($-14 \text{ }^\circ\text{C}$, $-16 \text{ }^\circ\text{C}$, $-18 \text{ }^\circ\text{C}$, $-20 \text{ }^\circ\text{C}$, and $-22 \text{ }^\circ\text{C}$) were calculated
from the 3 h filter data in Figure 1a and are illustrated in Figure 2ig, while the sampling times are shown as purple
bars throughout the figure. The air mass category and ID developed by Gat et al. (2021) for the campaign is provided
for each day at the top of the image based on wind directions and air mass back trajectories. HYSPLIT air mass back
trajectories for 10:00 and 18:00 local time are provided in Figure S17 in the SI, and are in good agreement with those
565 of Gat et al. (2021), who used a Lagrangian analysis tool, LAGRANTO 2.0 (Sprenger and Wernli, 2015) for analysis
of the same time periods.

The campaign began with the tail-end of a Saharan dust storm event from the southwest (SW_D) (Figure 2e)
on the 25th October, with the air mass also passing over the fertile land of the Nile Delta (Figure S17 in the SI). This
air mass yielded the highest aerosol concentrations throughout the campaign, with particle surface area concentration
570 modes for particles of $\sim 0.2 \mu\text{m}$ and $\sim 4 \mu\text{m}$ diameter, and some of the highest INP concentrations throughout the
campaign. This was immediately followed by an intense rainstorm later that same night (Figure 2d), which saw
aerosol concentrations plummet for the following two days with the air masses coming from the northwest through
Central and Eastern Europe and over the Mediterranean Sea. INP concentrations at the colder temperatures ($-20 \text{ }^\circ\text{C}$
and $-22 \text{ }^\circ\text{C}$) dropped significantly on the 26th October compared to the SW_D event, while they remained somewhat
575 comparable at warmer temperatures ($-16 \text{ }^\circ\text{C}$ and $-18 \text{ }^\circ\text{C}$). Concentrations at all of these temperatures dropped further
on the 27th of October during the same NW trajectory period, although there were multiple changes in wind direction
throughout those sampling times as shown in Figure 2e which may have influenced those results. The 20-60 min
impinger samples from the 26th and 27th of October exhibited the lowest INP concentrations in the impinger data
(Figure S13 in the SI), consistent with the data from the 3 h filter samples.

580



585 **Figure 2: Time series of meteorological and aerosol data for the field campaign in Rehovot, Israel, with dates in the DD/MM/YYYY format. The air mass category for each day is provided at the top (e.g. SW_D, NW1...). 3 h filter sampling times are shown as vertical purple bars. (a-e) Meteorological data from the Israel Meteorological Service (IMS) station at Bet Dagan: (a) Relative humidity. (b) Atmospheric pressure. (c) Temperature, (d) Rainfall, (e) Wind speed, with wind direction represented by coloured arrows: westerly winds are shown in shades of red and easterly winds in shades of blue. (f) Total aerosol particle concentration, dN , determined using the OPC. (g) PM_{10} , (for $\rho = 1 \text{ g cm}^{-3}$ and 2.65 g cm^{-3}), $PM_{2.5}$ and PM_1 mass concentrations from the OPC data (hourly moving average), alongside PM_{10} concentrations from the Israeli Ministry of Environment (IME)IMS station near the sampling site. (h) Particle size (D_p) distributions ($dN/d\log D_p$) obtained using the OPC. (i) Background-subtracted ice-nucleating particle concentrations (N_{INP}) at $-14 \text{ }^\circ\text{C}$, $-16 \text{ }^\circ\text{C}$, $-18 \text{ }^\circ\text{C}$, $-20 \text{ }^\circ\text{C}$, and $-22 \text{ }^\circ\text{C}$ during each sampling period.**

590

595 The 28th to the 31st of October saw the air masses coming from the east and carrying dust from Syria (the E_D category). This saw an increase in particle concentrations and surface area concentrations for larger particle sizes, in addition to higher INP concentrations, similar to those in the SW_D event. INP concentrations (at -18 °C to -22 °C) remained consistently high throughout this 4-day E_D period. Interestingly, while the 72 h air masses back trajectories on the 30th and 31st October came almost entirely over land from the east, those on the 28th largely came from Europe, 600 passing over a portion of the Mediterranean Sea, and only spent a relatively short time over the land to the East of the sampling location. Impinger samples were collected for 20 min during the afternoons of the 28th to 30th October and these showed high INP concentrations (Figure S13 in the SI), again consistent with the 3 h filter samples.

The afternoon of the 29th October was of particular interest, with notably high INP concentrations at -14 °C to -16 °C, with the 20 min impinger sample from that time yielding the highest INP concentrations of the impinger 605 dataset. Here, the air mass came from the North ([see the HYSPLIT air mass back trajectory in Figure 3](#)), passing over the eastern side of the Black Sea and over the region known as the Fertile Crescent, which spans the northern regions of Israel, Syria, and Iraq. The Fertile Crescent is the historical origin of agriculture and animal herding that started around 12,000 years ago (Salamini et al., 2002), though in modern times has suffered from long-term drying and is increasingly prone to drought (Kelley et al., 2015; Zittis et al., 2022), but still remains an active location for agriculture and vegetation (Zaitchik et al., 2007). Given this unique air mass trajectory amongst the rest of the dataset, the high 610 INP concentrations at warmer temperatures, and the “hump” in the INP spectrum that may indicate biological INP content as mentioned earlier, this data is hence discussed in further detail later in the manuscript.

Following the E_D dust event, the remainder of the campaign from 1st to 4th November was characterised by easterly winds and air mass back trajectories that were similar to the preceding two days, but with lower aerosol 615 particle concentrations. With this decrease in aerosol concentration came a steady decrease in INP concentrations at the colder temperatures, and a general trend of INP concentrations in the afternoons being higher than those in the morning. The 20 min impinger samples from the 2nd and 3rd November also showed decreasing activity following the E_D event (see Figure S13 in the SI).





Figure 3: Air mass back trajectory (72 h duration) for the 29th October 2018 (E_{D2}), obtained using NOAA’s HYSPLIT tool, showing three arrival heights at the sampling site: 500 m (red), 1,500 m (blue), and 2,500 m (green) above ground level (AGL).

625 3.4 Potential biological INP content

As discussed earlier, some INP concentrations in Figure 1a showed a “hump” of higher activity in the spectra at warmer temperatures, which may be indicative of biological INP activity that could be associated with dust or soil particles (O'Sullivan et al., 2014; Conen et al., 2011; Schnell, 1977; Schnell and Vali, 1976). The most notable sample spectra exhibiting this potential “biological hump” was from the afternoon of 29th October (Figure 1a). That specific
630 air mass had passed from Eastern Europe and over the Fertile Crescent to the north of the sampling site (Figure 3) during the easterly dust event (E_{D2}). This was corroborated by a similar hump in the 20 min impinger data during that same timeframe (see Figure S18 in the SI).

By coincidence, the sampling on the afternoon of the 29th October was performed simultaneously with two BGI PQ100 filter samplers: one set up for PM₁₀ and one set up for PM₁. The results for these can be seen in Figure
635 1a, with a more direct comparison in Figure S18 in the SI alongside the impinger data from the same timeframe. The N_{INP} of the PM₁ aerosols was relatively high, though also notably lower than for PM₁₀, suggesting that while much of the ice-nucleating activity was due to particles in the 1-10 μm size range, there was still a lot of activity caused by particles $<1 \mu\text{m}$. The PM₁ data also showed the same “biological” hump as in the PM₁₀ data.



640

Figure 34: Box-and-whisker pPlot showing the effect of heat treatment (95 °C for 30 min, as per Daily et al. (2022)) on the ice-nucleating activity of aqueous particle suspensions obtained from 3 h filter samples. This heat treatment is used as an indicator for the potential presence of proteinaceous INPs should there be a notable loss of activity at warmer temperatures following the heat treatment. Boxes represent 1 standard deviation from the mean (1σ; 68 %) while whiskers represent 2 standard deviations (2σ; 95 %). Dates are given in the YYMMDD format.

645

As a further means to test for biological ice-nucleating activity, we performed heat treatments (95 °C for 30 min) (Christner et al., 2008a; Daily et al., 2022) on nearly every sample collected and compared the ice-nucleating activities of the original samples to the heated samples in Figure 3-4 for the 3 h filter samples. The heat test results for the 24 h filter samples and the impinger samples are provided in Figure S19 and Figure S20 in the SI, respectively.

Heat treatment of the sample from the afternoon of 29th October demonstrated a significant loss in activity at the warmer parts of the PM₁₀ and PM₁ curves (Figure 34), i.e. a loss of the “hump” in each case (also see Figure S18 in the SI). The heat treatment of an impinger sample also showed a flattening of the hump (see Figure S18 in the SI), though not a similar dramatic loss in activity as for the filter samples. The heat treatment is not a definitive test for biological material, but since we expect the mineral ice-nucleating activity to be controlled by K-feldspar in this region (Boose et al., 2016) we believe it to be a reasonable indicator that the “humps” in the N_{INP} data were due to biological INP activity given the caveats of the procedure described by Daily et al. (2022). Thus, it is possible that biological INPs may have been entrained into the air mass as it passed over the Fertile Crescent to the north of the sampling site (see Figure 3). While speculative, this suggests that the Fertile Crescent could be a potential source of high temperature, heat-labile INPs for the Eastern Mediterranean and Middle Eastern regions.

While not as pronounced, the N_{INP} data from the morning of the 29th October also exhibited a notable hump that was lost following heat treatment, with the air mass for that sample having passed over a small region of the Fertile Crescent to the north of the sampling site. No other sampled air masses (for 500 m AGL) passed over the same area throughout the campaign, making these measurements unique, but the hypotheses therefore somewhat speculative. However, Gong et al. (2019) noted a potential biological signal in their INP concentrations at temperatures warmer than -15 °C during their campaign in Cyprus and suggested it originated locally, but we note that a handful of their air mass back trajectories appeared to pass over the Fertile Crescent and the Nile Delta regions.

A handful of other N_{INP} spectra potentially displayed smaller potential biological humps and were found to be sensitive to heat treatment (Figure 34). The sample for the afternoon of 25th October also comprised from a unique air mass for the campaign, -originating from the west of Italy before travelling over the north of the Sahara Desert and passing over the Nile Delta, the southeasternmost part of the Fertile Crescent, which may have contributed biological INPs to the sample. This would also agree with the findings of Beall et al. (2022) during the Red Sea leg of their cruise when they encountered air masses from the Nile Delta that showed heat-labile, likely biological, INP content. The sample from the afternoon of 26th October also demonstrated a small degree of heat sensitivity following the rain event that washed a lot of dust from the air, with the following air mass passing over Central/Eastern Europe and the Mediterranean Sea prior to reaching the sampling site.

Samples for the 1st (morning) and 3rd (afternoon) of November, obtained during the easterly air mass event (E1-E3) may have shown a minor decrease in N_{INP} following heat treatment in the 3 h filter and/or the impinger samples, and the air masses did not appreciably differ from those throughout the rest of the easterly event. We do note, however, that the Fertile Crescent extends to eastern Iraq and western Iran, and so could potentially have had

an influence on air masses encountered by the sampling site, although the contribution of biological INPs here was relatively minor, particularly compared to the results from the air masses that passed over the northern Fertile Crescent (E_{D2}).

685 Israel has varying airborne microbiomes that depend on air mass, and Gat et al. (Gat et al., 2017; Gat et al., 2021; Erkorkmaz et al., 2023; Peng et al., 2021) have measured a variety of desert- and soil-associated bacteria and fungi in air masses that passed over the Nile Delta and the Arabian Peninsula, with some of the latter cases also containing agricultural soil bacteria. Samples from days with no dust events mostly contained bacteria associated with human activity. Interestingly, dust storms were found to lower the proportion of fungi in the airborne microbiome of the region (Peng et al., 2021). Further, biological components found in soil dust associated with wind events in the
690 region have been found to exhibit reduced biological activity after sterilisation, which further supports the heat treatment results and the hypothesis presented here (Pardo et al., 2017).

Throughout the October-November 2018 measurement campaign, Gat et al. (2021) performed an in-depth analysis of the microbiome of the air masses via amplicon sequencing of the 16S rRNA gene and ribosomal internal transcribed spacer (ITS) region to study size-resolved bacteria and fungi, respectively. Their results showed that
695 bioaerosol communities varied with air mass, and that there was significant transport of bacteria as aggregates or attached to dust particles during the SW_D and E_D dust events (Erkorkmaz et al., 2023; Gat et al., 2021). Bacterial communities, particularly those of single cells smaller than 0.6 µm, were comprised of mixtures from local and transported sources. Further analysis of that data, presented here in Figure 5, demonstrates the highest concentration of airborne fungal (ITS) gene copy numbers (GCNs) across the three particle size ranges (fine, intermediate, coarse)
700 during the SW_D event on the 25th of October, where the air mass had travelled over the Nile Delta, while the bacterial (16S) concentrations were the highest in both the coarse and intermediate modes during that event. This finding is supported by recent study by Erkorkmaz et al. (2023), who confirmed that more bacteria are transported in the large size fraction in dust plumes across the Eastern Mediterranean.

705 The GCN data for bacteria and fungi during the 29th of October (E_{D2}) appear unspectacular for the coarse mode of particles, but closer inspection reveals that the fine and intermediate fungal concentrations were higher than all other days except for the 25th of October (SW_D). The bacterial concentrations in the intermediate size range were the second highest of the campaign during E_{D2}, while the fine particle concentrations were relatively high. The biological content from the Easterly (E) direction air masses (1st and 2nd ~~October~~ November) was also relatively high, particularly in the coarse mode.




Figure 5: Plots showing size-resolved biological aerosol particle concentrations, in terms of gene copy number (GCN) per litre of air, for (a) bacteria, analysed using 16S ribosomal RNA, and (b) fungi, using the ribosomal internal transcribed spacer (ITS) region. Particle size ranges were stratified into: coarse (3 μm – 18 μm), intermediate (0.8 μm – 3 μm) and fine (<1.3 μm).

While these values are for total populations rather than populations known to have ice-nucleating activity, together with the heat treatment data and the findings of Gat et al. (2021), this data further supports the hypothesis that biological INPs may have been entrained in air masses that passed over the fertile soils Fertile Crescent, particularly the Nile Delta region and the northern Syria region. This remains speculative given the lack of data, but suggests that these regions may warrant further exploration of their potential contributions to the regional ice-nucleating particle populations.

3.5 Ice-active site surface density, $n_s(T)$

725 The ice-nucleating activity of the aerosol in the region throughout the campaign was quantified via calculation of the
ice-active site density per surface area, $n_s(T)$ (cm^{-2}), from the INP concentrations (L^{-1}) and the surface area
concentrations obtained from the combined OPC and SMPS ($\chi = 1.1 \pm 0.1$) data, using Equation (5). The resultant
spectra for $n_s(T)$ are shown in Figure ~~5a~~6a, while Figure ~~5b~~6b shows the same data separated into “dust events” and
“clean days”. Figure ~~5e~~6c provides a comparison of these $n_s(T)$ values to the relevant literature, here with the
campaign data separated into “dust events” (SW_D and E_D air masses) and “clean days” (NW and E air masses). Most
of the data for the campaign lie on top of each other at the colder temperatures (< -18 °C), with some of the samples
730 from the E air mass (clean days) being lower than the bulk of the data.




Figure 56: Plots showing the ice-active site surface density, $n_s(T)$ for INPs in Israel. (a) Background-subtracted $n_s(T)$ spectra for 3 h filter samples. Dates are given in the YYMMDD format. (b) The 3 h filter sample data separated into dust events (brown) and clean days (blue). (c) Comparison of $n_s(T)$ values to relevant literature, including $n_s(T)$ data from Israel and the East Mediterranean region (Ardon-Dryer and Levin, 2014; Beall et al., 2022; Gong et al., 2019; Reicher et al., 2019; Reicher et al., 2018; Tarn et al., 2020). Several parameterisations for mineral dust are provided, including those for surface samples from arid locations around the world (Ullrich et al., 2017), sieved and milled surface samples from Israel (Boose et al., 2016), supermicron and submicron particles in desert dust storms (Reicher et al., 2019), the Harrison parameterisation for K-feldspar assuming K-feldspar comprised around 1 wt% of airborne mineral dust (Harrison et al., 2019), and O'Sullivan parameterisation for fertile soils (O'Sullivan et al., 2014).

Our data was around 2 orders of magnitude smaller than the parameterisation of Ullrich et al. (2017) derived from soil samples and precipitated dust samples collected from a number of arid locations, including Israel and the Nile
745 Delta. However, the bulk of the data below $-18\text{ }^{\circ}\text{C}$ match very well with the K-feldspar parameterisation developed
by Harrison et al. (2019), here being scaled to 1 wt% (noting that Boose et al. (2016), also plotted here, found $1.7 \pm$
 $0.4\text{ }\%$ wt% and $1.3 \pm 0.4\text{ }\%$ wt% K-feldspar in sieved and milled surface dust from Israel, respectively). The data at
temperatures warmer than about $-20\text{ }^{\circ}\text{C}$ lies increasingly higher than Harrison et al. (2019) and within the lower
uncertainties of the two parameterisations of Reicher et al. (2019) for supermicron and submicron dust during dust
750 events. Some samples overlap even more with the Reicher et al. (2019) regimes, such as “181029 afternoon” which
likely had more influence from biological INPs. Sample “181026 afternoon” demonstrated a high $n_s(T)$ value since
particle concentrations had dropped following a rain event, and thereafter may have been influenced by bioaerosols
in the air mass passing over Europe and the Mediterranean Sea.

Our data overlapped with some of the higher $n_s(T)$ data of Gong et al. (2019) from Cyprus, which followed
755 the Harrison et al. (2019) K-feldspar into the warmer temperatures. However, the data of Beall et al. (2022) from the
Red Sea matched ours very well. As for the N_{INP} comparison, the Israel data of Ardon-Dryer and Levin (2014) was
largely an order of magnitude or so higher in $n_s(T)$ than our observations, with only one of their clean day datasets
being comparable though their data does compare well with the Ullrich et al. (2017) parameterisation for desert dust.
The fertile soil parameterisation from O'Sullivan et al. (2014) is plotted and also sits much higher than our data, being
760 somewhat more similar to the Ullrich et al. (2017) parameterisation.

Some of the collected data compared well at the colder temperatures with that of Reicher et al. (2018) for
dust storms. The data shown from Tarn et al. (2020) represents a 3 h filter (“181031 afternoon”), as analysed using
the LOC-NIPI microfluidic platform, and is complementary to the datasets obtained using the $\mu\text{L-NIPI}$ for this
campaign.

765 That the $n_s(T)$ values for the clean days and the dust event days largely collapsed on top of each other,
particularly at temperatures $< -18\text{ }^{\circ}\text{C}$, is not surprising given the atmosphere always contains dust particles and so
the term “clean” is relative only to the high loadings measured during elevated dust events. Likewise, the activities
below $-18\text{ }^{\circ}\text{C}$ fall onto the Harrison et al. (2019) parameterisation for 1 wt% K-feldspar is consistent with the INP
population being dominated by K-feldspar mineral dust, with samples exhibiting increased $n_s(T)$ at warmer
770 temperatures potentially being due to biological content.

However, the “cleanest” day was the 26th October following an intense rain event the previous night (see
Figure 2d) that resulted in low aerosol concentrations, yet surprisingly high INP concentrations considering.
Interestingly, the midday INP concentrations from a more westerly wind direction were notably higher than those
from the afternoon more northwesterly direction (see Figure 2e). The surprisingly high INP concentrations following

775 the rain event resulted in high $n_s(T)$ activities given the low total aerosol concentrations, the reasons for which are unclear.

Overall, these results bring us to a similar conclusion as Gong et al. (2019) and Beall et al. (2022), that, while $n_s(T)$ parameterisations for mineral dust may be relevant for field samples below around -15 to -18 °C when the population is expected to be dominated by K-feldspar, they are not representative of regional INP activity at
780 temperatures much warmer than -18 °C when the INP population may contain substantial biological content. Other methods of representing INPs at warmer temperatures may therefore be warranted to account for multiple temperature-dependent INP populations that vary with location. This could, for example, be achieved via the application of region-and/or season-specific or $n_s(T)$ parameterisations, such as those of Reicher et al. (2019) for dust events in Israel that captured both the mineral and potential biological INP populations measured during the campaign described here. Alternatively, ~~Therefore, as suggested by Gong et al. (2019) demonstrated the use of and Beall et al. (2022), it may be that other methods of representation of INP activity are warranted. This could include the use of frequency distributions (or probability density functions, (PDFs))~~ ~~Welti et al. (2018) to determine that their measured INP populations in the Eastern Mediterranean were from long-range transport rather than local sources. These use a lognormal distribution to approximate the variability in INP concentration at each temperature, and assumes random~~
785 ~~dilution upon transport in the atmosphere; a skewed frequency distribution indicates an INP population from a local source, while a regular bell shape would represent long-range transport (Welti et al., 2018) from observed INP concentrations, as suggested by Gong et al. (2019) or $n_s(T)$ parameterisations for specific regions and events/seasons, such as those of Reicher et al. (2019) for dust events in Israel that here captured both the K-feldspar INP population but also the potential biological INP population indicated in these results.~~ This method was recently employed by
790 Frostenberg et al. (2023) in the development of a new temperature-dependent INP parameterisation based on lognormally-distributed INP concentration frequencies, without requiring aerosol information, that could be implemented into models using regional and seasonal INP frequency distributions.

4 Conclusions

The Eastern Mediterranean is an interesting location for aerosol and INP analysis, encountering air masses from
800 Europe, sea-spray aerosol from the Mediterranean Sea, and dust storms from the Sahara and Arabian Deserts, with the local atmosphere always containing some degree of dust loading. Given such variation in sources and transport of INPs, it is important to try to understand their potential and comparative effects on the region. Here, we undertook an intensive two-week INP sampling campaign in Israel to monitor concentrations and activity every morning and afternoon, with coincident measurements of local aerosol concentrations. Four main air masses were encountered
805 during the campaign, each offering some insights into the INP populations in the region. A Saharan dust event from the south-west yielded high INP concentrations that notably featured a potential biological component, possibly as a

result of passing over the fertile Nile Delta shortly prior to reaching the sampling site. Northwesterly winds that brought an air mass from central/eastern Europe and crossed a section of the Mediterranean Sea brought very low concentrations of aerosols and INPs, though this also followed a rain event that reduced the overall aerosol loadings in the region. Easterly winds brought air masses from Syria and the Arabian Desert with both high (dust events) and low aerosol loadings. The high loadings yielded high INP concentrations that tailed off as the dust loading of the air masses decreased over several days. Some of these samples from the Arabian Desert region may have had a slight biological INP component ~~in some cases~~ but this was inconclusive.

The final and most ~~remarkable-unique~~ air mass travelled from the north, in particular Eastern Europe, before turning west towards the sampling site, notably passing over fertile soils located in the area known as the Fertile Crescent. The Fertile Crescent has undergone substantial changes in the last century that have resulted in considerable drying and droughts, particularly since the 1950s when irrigation of the marsh region diverted waters from the Tigris and Euphrates rivers (“Fertile Crescent” (History, 2017; National-Geographic, 2022)). This was further exacerbated in Iraq in 1991 when ~~Saddam Hussein’s~~ the government built dams and dikes to drain the marshes. However, although many dams were later broken following the regime change in 2003 that allowed waters to flood back into parts of the region (Reuters, 2016). While the region is far less fertile than it once was, it still comprises a vast region of lands and dried soils that contain bacteria and fungi, in addition to farmlands, wetlands, rivers and marshes. It is possible that this change in land use can cause the soils to be more mobile and sensitive to winds. For example, the presence of dams, water transfer policies and extensive droughts throughout the region have been linked to an increase in dust aerosol optical depth in the western and southern regions of Iran (Ghasem et al., 2012; Hamzeh et al., 2021), while- Iraq saw an increased frequency of dust events from 1980-1993, a decreased frequency from 1993-2001, and an increase again after 2001 (Attiya and Jones, 2020).

The INP concentrations measured during the two-week campaign in the Eastern Mediterranean in 2018 were not only high, but heating analysis pointed towards a large biological component of INPs active at warmer temperatures in some samples. Given the broad territory that the Fertile Crescent covers, including Israel, Lebanon, Syria, and Iraq in the east, it is possible that some small amount of biological ice-nucleating activity encountered in some of the Easterly air masses also originated from these fertile lands, while signals that passed over the Nile Delta and northern regions containing fertile soils exhibited a much stronger signal.

In all, INP activity in the region throughout the campaign was largely consistent with being controlled by ~1 % K-feldspar mineral dust at temperatures below about -15 to -18 °C. However, the measurements associated with each of the four air masses suggested that the INP population in Israel can be influenced by biological INPs activated at warmer temperatures, whose origins are associated with fertile lands around the Eastern Mediterranean and the Middle East, most notably the Fertile Crescent and including the Nile Delta. To our knowledge, this is the first time that the Fertile Crescent, apart from a discussion of the Nile Delta by Beall et al. (2022), has been discussed as a

840 potential source of high-temperature biological INPs for the region around it, likely due to a lack of measurements associated with that particular location.

While this hypothesis is ~~still-very~~ speculative given the ~~limited~~~~small~~ number of relevant sampling events, the capacity for these fertile lands periodically influencing INP populations in the surrounding areas is compelling, particularly should the current long-term drying trends continue and so alter the INP populations. Thus, we believe
845 that the Fertile Crescent regions warrant more dedicated analyses of their associated INP populations in order to understand ~~whether they could~~~~their potential have an~~ influence on the vast surrounding regions, and to what extent if that is the case.-

Our results also bring us to similar conclusions to those of Gong et al. (2019) and Beall et al. (2022), in that $n_s(T)$ parameterisations for mineral dust are not necessarily appropriate for estimating INP activity in models when INP populations
850 active at warmer temperatures may dominate the mineral population and skew $n_s(T)$ values based on particle surface area. Other methods of INP representation may be required, or a database of regional and seasonal $n_s(T)$ or N_{INP} parameterisations based on observations.

Conflicts of interest

One of the authors is a member of the Editorial Board of *Aerosol Research*.

855 Acknowledgements

The authors thank the Weizmann–UK Making Connections Program, the European Research Council (ERC; grant no. 648661 MarineIce) and the Engineering and Physical Sciences Research Council (EPSRC; grant no. EP/R513258/1) for funding. Grace C. E. Porter is thanked for assistance with the INP background subtraction calculations. Burak Adnan Erkorkmaz is thanked for discussions.

860 Data availability

The data sets for this paper ~~will be made~~~~are~~ publicly available in the University of Leeds Data Repository (<https://doi.org/10.5518/1487>~~<https://archive.researchdata.leeds.ac.uk/>~~)(Tarn et al., 2024).

Author contributions

MDT, NR, YR and BJM designed the study. MDT, YR and BJM obtained funding for the study. MDT, NR, MA, SNFS and
865 ADH organised the logistics of the campaign, and prepared and tested equipment for the study. MDT, BVW, MA, NR and DG

performed the experiments. MDT, BVW, MA, NR, DG and AS-M analysed the data from the study. YR and BJM supervised the study. MDT, NR, YR and BJM wrote the paper, and all authors contributed to editing.

References

- 870 Adams, M. P., Tarn, M. D., Sanchez-Marroquin, A., Porter, G. C. E., O'Sullivan, D., Harrison, A. D., Cui, Z., Vergara-
Temprado, J., Carotenuto, F., Holden, M. A., Daily, M. I., Whale, T. F., Sikora, S. N. F., Burke, I. T., Shim, J. U., McQuaid,
J. B., and Murray, B. J.: A Major Combustion Aerosol Event Had a Negligible Impact on the Atmospheric Ice-Nucleating
Particle Population, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032938,
<https://doi.org/10.1029/2020JD032938>, 2020.
- 875 Ardon-Dryer, K., and Levin, Z.: Ground-based measurements of immersion freezing in the eastern Mediterranean, *Atmos.
Chem. Phys.*, 14, 5217-5231, 10.5194/acp-14-5217-2014, 2014.
- Athanasopoulou, E., Protonotariou, A., Papangelis, G., Tombrou, M., Mihalopoulos, N., and Gerasopoulos, E.: Long-range
transport of Saharan dust and chemical transformations over the Eastern Mediterranean, *Atmos. Environ.*, 140, 592-604,
<https://doi.org/10.1016/j.atmosenv.2016.06.041>, 2016.
- 880 Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., Dobbie, S., O'Sullivan, D.,
and Malkin, T. L.: The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds, *Nature*, 498, 355-
358, 10.1038/nature12278, 2013.
- Attiya, A. A., and Jones, B. G.: Climatology of Iraqi dust events during 1980–2015, *SN Applied Sciences*, 2, 845,
10.1007/s42452-020-2669-4, 2020.
- 885 Beall, C. M., Lucero, D., Hill, T. C., DeMott, P. J., Stokes, M. D., and Prather, K. A.: Best practices for precipitation sample
storage for offline studies of ice nucleation in marine and coastal environments, *Atmos. Meas. Tech.*, 13, 6473-6486,
10.5194/amt-13-6473-2020, 2020.
- Beall, C. M., Hill, T. C. J., DeMott, P. J., Köneman, T., Pikridas, M., Drewnick, F., Harder, H., Pöhlker, C., Lelieveld, J.,
Weber, B., Iakovides, M., Prokeš, R., Sciare, J., Andreae, M. O., Stokes, M. D., and Prather, K. A.: Ice-nucleating particles
near two major dust source regions, *Atmos. Chem. Phys.*, 22, 12607-12627, 10.5194/acp-22-12607-2022, 2022.
- 890 Boose, Y., Welti, A., Atkinson, J., Ramelli, F., Danielczok, A., Bingemer, H. G., Plötze, M., Sierau, B., Kanji, Z. A., and
Lohmann, U.: Heterogeneous ice nucleation on dust particles sourced from nine deserts worldwide – Part 1: Immersion
freezing, *Atmos. Chem. Phys.*, 16, 15075-15095, 10.5194/acp-16-15075-2016, 2016.
- Burton, N. C., Grinshpun, S. A., and Reponen, T.: Physical Collection Efficiency of Filter Materials for Bacteria and Viruses,
The Annals of Occupational Hygiene, 51, 143-151, 10.1093/annhyg/mel073, 2006.
- 895 Carvalho, E., Sindt, C., Verdier, A., Galan, C., O'Donoghue, L., Parks, S., and Thibaudon, M.: Performance of the Coriolis
air sampler, a high-volume aerosol-collection system for quantification of airborne spores and pollen grains, *Aerobiologia*,
24, 191-201, 10.1007/s10453-008-9098-y, 2008.
- Ceppi, P., Brient, F., Zelinka, M. D., and Hartmann, D. L.: Cloud feedback mechanisms and their representation in global
climate models, *Wiley Interdisciplinary Reviews: Climate Change*, 8, e465, 10.1002/wcc.465, 2017.
- 900 Christner, B. C., Cai, R., Morris, C. E., McCarter, K. S., Foreman, C. M., Skidmore, M. L., Montross, S. N., and Sands, D. C.:
Geographic, seasonal, and precipitation chemistry influence on the abundance and activity of biological ice nucleators in
rain and snow, *Proc. Natl. Acad. Sci. U.S.A.*, 105, 18854-18859, 10.1073/pnas.0809816105, 2008a.
- Christner, B. C., Morris, C. E., Foreman, C. M., Cai, R., and Sands, D. C.: Ubiquity of Biological Ice Nucleators in Snowfall,
Science, 319, 1214-1214, 10.1126/science.1149757, 2008b.

- 905 Conen, F., Morris, C. E., Leifeld, J., Yakutin, M. V., and Alewell, C.: Biological residues define the ice nucleation properties of soil dust, *Atmos. Chem. Phys.*, 11, 9643-9648, 10.5194/acp-11-9643-2011, 2011.
- Daily, M. I., Tarn, M. D., Whale, T. F., and Murray, B. J.: An evaluation of the heat test for the ice-nucleating ability of minerals and biological material, *Atmos. Meas. Tech.*, 15, 2635-2665, 10.5194/amt-15-2635-2022, 2022.
- Dayan, U.: Climatology of Back Trajectories from Israel Based on Synoptic Analysis, *J. Clim. Appl. Meteorol.*, 25, 591-595, 10.1175/1520-0450(1986)025<0591:COBTFI>2.0.CO;2, 1986.
- 910 Dayan, U., Heffter, J., Miller, J., and Gutman, G.: Dust Intrusion Events into the Mediterranean Basin, *J. Appl. Meteorol.*, 30, 1185-1199, 1991.
- DeMott, P. J., Prenni, A. J., Liu, X., Kreidenweis, S. M., Petters, M. D., Twohy, C. H., Richardson, M. S., Eidhammer, T., and Rogers, D. C.: Predicting global atmospheric ice nuclei distributions and their impacts on climate, *Proceedings of the National Academy of Sciences*, 107, 11217-11222, 10.1073/pnas.0910818107, 2010.
- 915 DeMott, P. J., Prenni, A. J., McMeeking, G. R., Sullivan, R. C., Petters, M. D., Tobo, Y., Niemand, M., Möhler, O., Snider, J. R., Wang, Z., and Kreidenweis, S. M.: Integrating laboratory and field data to quantify the immersion freezing ice nucleation activity of mineral dust particles, *Atmos. Chem. Phys.*, 15, 393-409, 10.5194/acp-15-393-2015, 2015.
- Draxler, R. R., and Hess, G. D.: An overview of the HYSPLIT_4 modelling system for trajectories, dispersion and deposition, *Australian Meteorological Magazine*, 47, 295-308, 1998.
- 920 Els, N., Larose, C., Baumann-Stanzer, K., Tignat-Perrier, R., Keuschnig, C., Vogel, T. M., and Sattler, B.: Microbial composition in seasonal time series of free tropospheric air and precipitation reveals community separation, *Aerobiologia*, 35, 671-701, 10.1007/s10453-019-09606-x, 2019.
- Erkorkmaz, B. A., Gat, D., and Rudich, Y.: Aerial transport of bacteria by dust plumes in the Eastern Mediterranean revealed by complementary rRNA/rRNA-gene sequencing, *Commun. Earth Environ.*, 4, 24, 10.1038/s43247-023-00679-8, 2023.
- 925 Fleming, Z. L., Monks, P. S., and Manning, A. J.: Review: Untangling the influence of air-mass history in interpreting observed atmospheric composition, *Atmospheric Research*, 104-105, 1-39, <https://doi.org/10.1016/j.atmosres.2011.09.009>, 2012.
- Foner, H. A., and Ganor, E.: The chemical and mineralogical composition of some urban atmospheric aerosols in Israel, *Atmospheric Environment. Part B. Urban Atmosphere*, 26, 125-133, [https://doi.org/10.1016/0957-1272\(92\)90045-T](https://doi.org/10.1016/0957-1272(92)90045-T), 1992.
- 930 Frostenberg, H. C., Welti, A., Luhr, M., Savre, J., Thomson, E. S., and Ickes, L.: The chance of freezing – a conceptual study to parameterize temperature-dependent freezing by including randomness of ice-nucleating particle concentrations, *Atmos. Chem. Phys.*, 23, 10883-10900, 10.5194/acp-23-10883-2023, 2023.
- Funatsu, B. M., Claud, C., and Chaboureau, J.-P.: Potential of Advanced Microwave Sounding Unit to identify precipitating systems and associated upper-level features in the Mediterranean region: Case studies, 112, <https://doi.org/10.1029/2006JD008297>, 2007.
- 935 Gagin, A.: The Ice Phase in Winter Continental Cumulus Clouds, *J. Atmos. Sci.*, 32, 1604-1614, 10.1175/1520-0469(1975)032<1604:tipiwc>2.0.co;2, 1975.
- Ganor, E.: Analysis of atmospheric dust in Israel, *PhD Thesis*, Hebrew University, Jerusalem, 1975.
- Ganor, E., and Mamane, Y.: Transport of Saharan dust across the eastern Mediterranean, *Atmos. Environ.*, 16, 581-587, [https://doi.org/10.1016/0004-6981\(82\)90167-6](https://doi.org/10.1016/0004-6981(82)90167-6), 1982.
- 940 Ganor, E.: The composition of clay minerals transported to Israel as indicators of Saharan dust emission, *Atmospheric Environment. Part A. General Topics*, 25, 2657-2664, [https://doi.org/10.1016/0960-1686\(91\)90195-D](https://doi.org/10.1016/0960-1686(91)90195-D), 1991.
- Ganor, E.: The frequency of Saharan dust episodes over Tel Aviv, Israel, *Atmos. Environ.*, 28, 2867-2871, [https://doi.org/10.1016/1352-2310\(94\)90087-6](https://doi.org/10.1016/1352-2310(94)90087-6), 1994.

- 945 Ganor, E., Foner, H. A., Bingemer, H. G., Udisti, R., and Setter, I.: Biogenic sulphate generation in the Mediterranean Sea and its contribution to the sulphate anomaly in the aerosol over Israel and the Eastern Mediterranean, *Atmos. Environ.*, 34, 3453-3462, [https://doi.org/10.1016/S1352-2310\(00\)00077-7](https://doi.org/10.1016/S1352-2310(00)00077-7), 2000.
- Ganor, E., Osetinsky, I., Stupp, A., and Alpert, P.: Increasing trend of African dust, over 49 years, in the eastern Mediterranean, *J. Geophys. Res. Atmos.*, 115, D07201, 10.1029/2009JD012500, 2010.
- 950 Garcia, E., Hill, T. C. J., Prenni, A. J., DeMott, P. J., Franc, G. D., and Kreidenweis, S. M.: Biogenic ice nuclei in boundary layer air over two U.S. High Plains agricultural regions, *J. Geophys. Res. Atmos.*, 117, D18209, 10.1029/2012jd018343, 2012.
- Gat, D., Mazar, Y., Cytryn, E., and Rudich, Y.: Origin-dependent variations of atmospheric microbiome community in Eastern Mediterranean dust storms, *Environmental Science & Technology*, submitted, 2017.
- 955 Gat, D., Reicher, N., Schechter, S., Alayof, M., Tarn, M. D., Wyld, B. V., Zimmermann, R., and Rudich, Y.: Size-Resolved Community Structure of Bacteria and Fungi Transported by Dust in the Middle East, 12, 10.3389/fmicb.2021.744117, 2021.
- Ghasem, A., Shamsipour, A., Miri, M., and Safarrad, T.: Synoptic and remote sensing analysis of dust events in southwestern Iran, *Natural Hazards*, 64, 1625-1638, 10.1007/s11069-012-0328-9, 2012.
- 960 Gobbi, G. P., Barnaba, F., and Ammannato, L.: The vertical distribution of aerosols, Saharan dust and cirrus clouds in Rome (Italy) in the year 2001, *Atmos. Chem. Phys.*, 4, 351-359, 10.5194/acp-4-351-2004, 2004.
- Gong, X., Wex, H., Müller, T., Wiedensohler, A., Höhler, K., Kandler, K., Ma, N., Dietel, B., Schiebel, T., Möhler, O., and Stratmann, F.: Characterization of aerosol properties at Cyprus, focusing on cloud condensation nuclei and ice-nucleating particles, *Atmos. Chem. Phys.*, 19, 10883-10900, 10.5194/acp-19-10883-2019, 2019.
- 965 Hamzeh, N. H., Kaskaoutis, D. G., Rashki, A., and Mohammadpour, K.: Long-Term Variability of Dust Events in Southwestern Iran and Its Relationship with the Drought, 12, 1350, 2021.
- Harrison, A. D., Whale, T. F., Carpenter, M. A., Holden, M. A., Neve, L., O'Sullivan, D., Vergara Temprado, J., and Murray, B. J.: Not all feldspars are equal: a survey of ice nucleating properties across the feldspar group of minerals, *Atmos. Chem. Phys.*, 16, 10927-10940, 10.5194/acp-16-10927-2016, 2016.
- 970 Harrison, A. D., Lever, K., Sanchez-Marroquin, A., Holden, M. A., Whale, T. F., Tarn, M. D., McQuaid, J. B., and Murray, B. J.: The ice-nucleating ability of quartz immersed in water and its atmospheric importance compared to K-feldspar, *Atmos. Chem. Phys.*, 19, 11343-11361, 10.5194/acp-19-11343-2019, 2019.
- Harrison, A. D., O'Sullivan, D., Adams, M. P., Porter, G. C. E., Blades, E., Brathwaite, C., Chewitt-Lucas, R., Gaston, C., Hawker, R., Krüger, O. O., Neve, L., Pöhlker, M. L., Pöhlker, C., Pöschl, U., Sanchez-Marroquin, A., Sealy, A., Sealy, P., 975 Tarn, M. D., Whitehall, S., McQuaid, J. B., Carslaw, K. S., Prospero, J. M., and Murray, B. J.: The ice-nucleating activity of African mineral dust in the Caribbean boundary layer, *Atmos. Chem. Phys.*, 22, 9663-9680, 10.5194/acp-22-9663-2022, 2022.
- Hawker, R. E., Miltenberger, A. K., Wilkinson, J. M., Hill, A. A., Shipway, B. J., Cui, Z., Cotton, R. J., Carslaw, K. S., Field, P. R., and Murray, B. J.: The temperature dependence of ice-nucleating particle concentrations affects the radiative 980 properties of tropical convective cloud systems, *Atmos. Chem. Phys.*, 21, 5439-5461, 10.5194/acp-21-5439-2021, 2021.
- Herbert, R. J., Murray, B. J., Dobbie, S. J., and Koop, T.: Sensitivity of liquid clouds to homogenous freezing parameterizations, *Geophys. Res. Lett.*, 42, 1599-1605, 10.1002/2014gl062729, 2015.
- History article: "Fertile Crescent": <https://www.history.com/topics/pre-history/fertile-crescent>, access: September 2023, 2017.
- Hoffmann, F., Raasch, S., and Noh, Y.: Entrainment of aerosols and their activation in a shallow cumulus cloud studied with 985 a coupled LCM-LES approach, *Atmospheric Research*, 156, 43-57, <https://doi.org/10.1016/j.atmosres.2014.12.008>, 2015.

- Holden, M. A., Whale, T. F., Tarn, M. D., O’Sullivan, D., Walshaw, R. D., Murray, B. J., Meldrum, F. C., and Christenson, H. K.: High-speed imaging of ice nucleation in water proves the existence of active sites, *Sci. Adv.*, 5, eaav4316, 10.1126/sciadv.aav4316, 2019.
- 990 Holden, M. A., Campbell, J. M., Meldrum, F. C., Murray, B. J., and Christenson, H. K.: Active sites for ice nucleation differ depending on nucleation mode, *Proceedings of the National Academy of Sciences*, 118, e2022859118, 10.1073/pnas.2022859118, 2021.
- Hoose, C., and Möhler, O.: Heterogeneous ice nucleation on atmospheric aerosols: a review of results from laboratory experiments, *Atmos. Chem. Phys.*, 12, 9817-9854, 10.5194/acp-12-9817-2012, 2012.
- 995 Huffman, J. A., Prenni, A. J., DeMott, P. J., Pöhlker, C., Mason, R. H., Robinson, N. H., Fröhlich-Nowoisky, J., Tobo, Y., Després, V. R., Garcia, E., Gochis, D. J., Harris, E., Müller-Germann, I., Ruzene, C., Schmer, B., Sinha, B., Day, D. A., Andreae, M. O., Jimenez, J. L., Gallagher, M., Kreidenweis, S. M., Bertram, A. K., and Pöschl, U.: High concentrations of biological aerosol particles and ice nuclei during and after rain, *Atmos. Chem. Phys.*, 13, 6151-6164, 10.5194/acp-13-6151-2013, 2013.
- 1000 *Coriolis μ Air Sampler product information*, <https://www.bertin-technologies.com/product/air-samplers/coriolis-micro-air-sampler>, Bertin Technologies, access: 23rd January 2024.
- Israelevich, P. L., Levin, Z., Joseph, J. H., and Ganor, E.: Desert aerosol transport in the Mediterranean region as inferred from the TOMS aerosol index, 107, AAC 13-11-AAC 13-13, <https://doi.org/10.1029/2001JD002011>, 2002.
- Kahl, J. D., and Samson, P. J.: Uncertainty in Trajectory Calculations Due to Low Resolution Meteorological Data %J *Journal of Applied Meteorology and Climatology*, 25, 1816-1831, [https://doi.org/10.1175/1520-0450\(1986\)025<1816:UITCDT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1986)025<1816:UITCDT>2.0.CO;2), 1986.
- 1005 Kanji, Z. A., Ladino, L. A., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D. J., and Krämer, M.: Overview of Ice Nucleating Particles, *Meteorological Monographs*, 58, 1.1-1.33, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, 2017.
- Karaca, F., Anil, I., and Alagha, O.: Long-range potential source contributions of episodic aerosol events to PM10 profile of a megacity, *Atmos. Environ.*, 43, 5713-5722, <https://doi.org/10.1016/j.atmosenv.2009.08.005>, 2009.
- 1010 Kelley, C. P., Mohtadi, S., Cane, M. A., Seager, R., and Kushnir, Y.: Climate change in the Fertile Crescent and implications of the recent Syrian drought, 112, 3241-3246, doi:10.1073/pnas.1421533112, 2015.
- Kiselev, A., Bachmann, F., Pedevilla, P., Cox, S. J., Michaelides, A., Gerthsen, D., and Leisner, T.: Active sites in heterogeneous ice nucleation—the example of K-rich feldspars, *Science*, 355, 367-371, 10.1126/science.aai8034, 2017.
- 1015 Knopf, D. A., Alpert, P. A., Zipori, A., Reicher, N., and Rudich, Y.: Stochastic nucleation processes and substrate abundance explain time-dependent freezing in supercooled droplets, *NPJ Clim. Atmos. Sci.*, 3, 2, 10.1038/s41612-020-0106-4, 2020.
- Krasnov, H., Katra, I., and Friger, M.: Increase in dust storm related PM10 concentrations: A time series analysis of 2001–2015, *Environmental Pollution*, 213, 36-42, <https://doi.org/10.1016/j.envpol.2015.10.021>, 2016.
- Kubilay, N., Nickovic, S., Moulin, C., and Dulac, F.: An illustration of the transport and deposition of mineral dust onto the eastern Mediterranean, *Atmos. Environ.*, 34, 1293-1303, [https://doi.org/10.1016/S1352-2310\(99\)00179-X](https://doi.org/10.1016/S1352-2310(99)00179-X), 2000.
- 1020 Levi, Y., and Rosenfeld, D.: Ice Nuclei, Rainwater Chemical Composition, and Static Cloud Seeding Effects in Israel, *J. Appl. Meteorol.*, 35, 1494-1501, 10.1175/1520-0450(1996)035<1494:inrcca>2.0.co;2, 1996.
- Levin, Z., and Lindberg, J. D.: Size distribution, chemical composition, and optical properties of urban and desert aerosols in Israel, *J. Geophys. Res. Oceans*, 84, 6941-6950, 10.1029/JC084iC11p06941, 1979.
- 1025 Levin, Z., Teller, A., Ganor, E., and Yin, Y.: On the interactions of mineral dust, sea-salt particles, and clouds: A measurement and modeling study from the Mediterranean Israeli Dust Experiment campaign, *J. Geophys. Res. Atmos.*, 110, D20202, 10.1029/2005jd005810, 2005.

- Lindsley, W. G.: Filter Pore Size and Aerosol Sample Collection, in: NIOSH Manual of Analytical Methods (NMAM), 5th ed., Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, 2016.
- 1030 Lohmann, U., and Feichter, J.: Global indirect aerosol effects: a review, *Atmos. Chem. Phys.*, 5, 715-737, 10.5194/acp-5-715-2005, 2005.
- Mamane, Y., Ganor, E., and Donagi, A. E.: Aerosol composition of urban and desert origin in the Eastern Mediterranean. I, *Water Air Soil Pollut.*, 14, 29-43, 10.1007/BF00291824, 1980.
- 1035 Marinou, E., Tesche, M., Nenes, A., Ansmann, A., Schrod, J., Mamali, D., Tsekeri, A., Pikridas, M., Baars, H., Engelmann, R., Voudouri, K. A., Solomos, S., Sciare, J., Groß, S., Ewald, F., and Amiridis, V.: Retrieval of ice-nucleating particle concentrations from lidar observations and comparison with UAV in situ measurements, *Atmos. Chem. Phys.*, 19, 11315-11342, 10.5194/acp-19-11315-2019, 2019.
- Marple, V. A., Rubow, K. L., and Behm, S. M.: A Microorifice Uniform Deposit Impactor (MOUDI): Description, Calibration, and Use, *Aerosol Science and Technology*, 14, 434-446, 10.1080/02786829108959504, 1991.
- 1040 Mason, R. H., Chou, C., McCluskey, C. S., Levin, E. J. T., Schiller, C. L., Hill, T. C. J., Huffman, J. A., DeMott, P. J., and Bertram, A. K.: The micro-orifice uniform deposit impactor–droplet freezing technique (MOUDI-DFT) for measuring concentrations of ice nucleating particles as a function of size: improvements and initial validation, *Atmos. Meas. Tech.*, 8, 2449-2462, 10.5194/amt-8-2449-2015, 2015.
- 1045 Möhler, O., Benz, S., Saathoff, H., Schnaiter, M., Wagner, R., Schneider, J., Walter, S., Ebert, V., and Wagner, S.: The effect of organic coating on the heterogeneous ice nucleation efficiency of mineral dust aerosols, *Environ. Res. Lett.*, 3, 025007, 10.1088/1748-9326/3/2/025007, 2008.
- Murray, B. J., O'Sullivan, D., Atkinson, J. D., and Webb, M. E.: Ice nucleation by particles immersed in supercooled cloud droplets, *Chem. Soc. Rev.*, 41, 6519-6554, 10.1039/C2CS35200A, 2012.
- 1050 Murray, B. J., Carslaw, K. S., and Field, P. R.: Opinion: Cloud-phase climate feedback and the importance of ice-nucleating particles, *Atmos. Chem. Phys.*, 21, 665-679, 10.5194/acp-21-665-2021, 2021.
- National Geographic article: "Fertile Crescent": <https://education.nationalgeographic.org/resource/fertile-crescent/>, access: September 2023, 2022.
- 1055 Nissenbaum, D., Sarafian, R., Rudich, Y., and Raveh-Rubin, S.: Six types of dust events in Eastern Mediterranean identified using unsupervised machine-learning classification, *Atmos. Environ.*, 309, 119902, <https://doi.org/10.1016/j.atmosenv.2023.119902>, 2023.
- O'Sullivan, D., Murray, B. J., Malkin, T. L., Whale, T. F., Umo, N. S., Atkinson, J. D., Price, H. C., Baustian, K. J., Browse, J., and Webb, M. E.: Ice nucleation by fertile soil dusts: relative importance of mineral and biogenic components, *Atmos. Chem. Phys.*, 14, 1853-1867, 10.5194/acp-14-1853-2014, 2014.
- 1060 O'Sullivan, D., Adams, M. P., Tarn, M. D., Harrison, A. D., Vergara-Temprado, J., Porter, G. C. E., Holden, M. A., Sanchez-Marroquin, A., Carotenuto, F., Whale, T. F., McQuaid, J. B., Walshaw, R., Hedges, D. H. P., Burke, I. T., Cui, Z., and Murray, B. J.: Contributions of biogenic material to the atmospheric ice-nucleating particle population in North Western Europe, *Sci. Rep.*, 8, 13821, 10.1038/s41598-018-31981-7, 2018.
- Pardo, M., Kutra, I., Schaeur, J. J., and Rudich, Y.: Mitochondria-mediated oxidative stress induced by desert dust in rat alveolar macrophages, *GeoHealth*, 1, 4-16, <https://doi.org/10.1002/2016GH000017>, 2017.
- 1065 Peng, X., Gat, D., Paytan, A., and Rudich, Y.: The Response of Airborne Mycobiome to Dust Storms in the Eastern Mediterranean, 7, 802, 2021.
- Peters, T. M., Ott, D., and O'Shaughnessy, P. T.: Comparison of the Grimm 1.108 and 1.109 portable aerosol spectrometer to the TSI 3321 aerodynamic particle sizer for dry particles, *Ann. Occup. Hyg.*, 50, 843-850, 10.1093/annhyg/mel067, 2006.

- 1070 Petters, M. D., and Wright, T. P.: Revisiting ice nucleation from precipitation samples, *Geophys. Res. Lett.*, 42, 8758-8766, 10.1002/2015GL065733, 2015.
- Prodi, F., Santachiara, G., and Oliosi, F.: Characterization of aerosols in marine environments (Mediterranean, Red Sea, and Indian Ocean), *Journal of Geophysical Research: Oceans*, 88, 10957-10968, <https://doi.org/10.1029/JC088iC15p10957>, 1983.
- 1075 Reicher, N., Segev, L., and Rudich, Y.: The Weizmann Supercooled Droplets Observation on a Microarray (WISDOM) and application for ambient dust, *Atmos. Meas. Tech.*, 11, 233-248, 10.5194/amt-11-233-2018, 2018.
- Reicher, N., Budke, C., Eickhoff, L., Raveh-Rubin, S., Kaplan-Ashiri, I., Koop, T., and Rudich, Y.: Size-dependent ice nucleation by airborne particles during dust events in the eastern Mediterranean, *Atmos. Chem. Phys.*, 19, 11143-11158, 10.5194/acp-19-11143-2019, 2019.
- 1080 Reuters article: "Iraq's marshes, once drained by Saddam, named world heritage site": <https://www.reuters.com/article/us-un-heritage-iraq-idUSKCN0ZX0SN>, access: October 2023, 2016.
- Roesch, C., Roesch, M., Wolf, M. J., Zawadowicz, M. A., AlAloula, R., Awwad, Z., and Cziczko, D. J.: CCN and INP activity of middle eastern soil dust, *Aeolian Research*, 52, 100729, <https://doi.org/10.1016/j.aeolia.2021.100729>, 2021.
- Rosenfeld, D., and Woodley, W. L.: Deep convective clouds with sustained supercooled liquid water down to -37.5 degrees C, *Nature*, 405, 440-442, 10.1038/35013030, 2000.
- 1085 Salamini, F., Özkan, H., Brandolini, A., Schäfer-Pregl, R., and Martin, W.: Genetics and geography of wild cereal domestication in the near east, *Nature Reviews Genetics*, 3, 429-441, 10.1038/nrg817, 2002.
- Sanchez-Marroquin, A., West, J. S., Burke, I. T., McQuaid, J. B., and Murray, B. J.: Mineral and biological ice-nucleating particles above the South East of the British Isles, *Environmental Science: Atmospheres*, 1, 176-191, 10.1039/D1EA00003A, 2021.
- 1090 Šantl-Temkiv, T., Amato, P., Gosewinkel, U., Thyraug, R., Charton, A., Chicot, B., Finster, K., Bratbak, G., and Löndahl, J.: High-Flow-Rate Impinger for the Study of Concentration, Viability, Metabolic Activity, and Ice-Nucleation Activity of Airborne Bacteria, *Environmental Science & Technology*, 51, 11224-11234, 10.1021/acs.est.7b01480, 2017.
- 1095 Sari, D., Incecik, S., and Ozkurt, N.: Analysis of surface ozone episodes using WRF-HYSPLIT model at Biga Peninsula in the Marmara region of Turkey, *Atmospheric Pollution Research*, 11, 2361-2378, <https://doi.org/10.1016/j.apr.2020.09.018>, 2020.
- Schnell, R. C., and Vali, G.: Biogenic Ice Nuclei: Part I. Terrestrial and Marine Sources, *J. Atmos. Sci.*, 33, 1554-1564, 10.1175/1520-0469(1976)033<1554:BINPIT>2.0.CO;2, 1976.
- Schnell, R. C.: Kaolin and a biogenic ice nucleant: Some nucleation and identification studies, IXth Nucleation Conference, Galway, Ireland, 1977, 353-356,
- 1100 Schrod, J., Weber, D., Drücke, J., Keleshis, C., Pikridas, M., Ebert, M., Cvetković, B., Nickovic, S., Marinou, E., Baars, H., Ansmann, A., Vrekoussis, M., Mihalopoulos, N., Sciare, J., Curtius, J., and Bingemer, H. G.: Ice nucleating particles over the Eastern Mediterranean measured by unmanned aircraft systems, *Atmos. Chem. Phys.*, 17, 4817-4835, 10.5194/acp-17-4817-2017, 2017.
- 1105 Soo, J.-C., Monaghan, K., Lee, T., Kashon, M., and Harper, M.: Air sampling filtration media: Collection efficiency for respirable size-selective sampling, *Aerosol Sci. Tech.*, 50, 76-87, 10.1080/02786826.2015.1128525, 2016.
- Sprenger, M., and Wernli, H.: The LAGRANTO Lagrangian analysis tool – version 2.0, *Geosci. Model Dev.*, 8, 2569-2586, 10.5194/gmd-8-2569-2015, 2015.
- 1110 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *Bulletin of the American Meteorological Society*, 96, 2059-2077, 10.1175/bams-d-14-00110.1, 2016.

- Storelvmo, T.: Aerosol Effects on Climate via Mixed-Phase and Ice Clouds, *Annual Review of Earth and Planetary Sciences*, 45, 199-222, 10.1146/annurev-earth-060115-012240, 2017.
- 1115 Tarn, M. D., Sikora, S. N. F., Porter, G. C. E., Wyld, B. V., Alayof, M., Reicher, N., Harrison, A. D., Rudich, Y., Shim, J.-u., and Murray, B. J.: On-chip analysis of atmospheric ice-nucleating particles in continuous flow, *Lab Chip*, 20, 2889-2910, 10.1039/D0LC00251H, 2020.
- Tarn, M. D., Wyld, B. V., Reicher, N., Alayof, M., Gat, D., Sanchez-Marroquin, A., Sikora, S. N. F., Harrison, A. D., Rudich, Y., and Murray, B. J.: Dataset for “Atmospheric ice-nucleating particles in the Eastern Mediterranean”, University of Leeds. [Dataset]. <https://doi.org/10.5518/1487>, 2024.
- 1120 Trueblood, J. V., Nicosia, A., Engel, A., Zäncker, B., Rinaldi, M., Freney, E., Thyssen, M., Obernosterer, I., Dinasquet, J., Belosi, F., Tovar-Sánchez, A., Rodriguez-Romero, A., Santachiara, G., Guieu, C., and Sellegri, K.: A two-component parameterization of marine ice-nucleating particles based on seawater biology and sea spray aerosol measurements in the Mediterranean Sea, *Atmos. Chem. Phys.*, 21, 4659-4676, 10.5194/acp-21-4659-2021, 2021.
- Ullrich, R., Hoose, C., Möhler, O., Niemand, M., Wagner, R., Höhler, K., Hiranuma, N., Saathoff, H., and Leisner, T.: A New Ice Nucleation Active Site Parameterization for Desert Dust and Soot, *Journal of the Atmospheric Sciences*, 74, 699-717, 10.1175/JAS-D-16-0074.1, 2017.
- 1125 Vali, G.: Quantitative Evaluation of Experimental Results on the Heterogeneous Freezing Nucleation of Supercooled Liquids, *J. Atmos. Sci.*, 28, 402-409, 10.1175/1520-0469(1971)028<0402:QEOERA>2.0.CO;2, 1971.
- Vali, G.: Freezing Rate Due to Heterogeneous Nucleation, *J. Atmos. Sci.*, 51, 1843-1856, 10.1175/1520-0469(1994)051<1843:FRDTHN>2.0.CO;2, 1994.
- 1130 Vali, G.: Repeatability and randomness in heterogeneous freezing nucleation, *Atmos. Chem. Phys.*, 8, 5017-5031, 10.5194/acp-8-5017-2008, 2008.
- Vali, G.: Revisiting the differential freezing nucleus spectra derived from drop-freezing experiments: methods of calculation, applications, and confidence limits, *Atmos. Meas. Tech.*, 12, 1219-1231, 10.5194/amt-12-1219-2019, 2019.
- 1135 Vergara-Temprado, J., Murray, B. J., Wilson, T. W., O'Sullivan, D., Browse, J., Pringle, K. J., Ardon-Dryer, K., Bertram, A. K., Burrows, S. M., Ceburnis, D., DeMott, P. J., Mason, R. H., O'Dowd, C. D., Rinaldi, M., and Carslaw, K. S.: Contribution of feldspar and marine organic aerosols to global ice nucleating particle concentrations, *Atmos. Chem. Phys.*, 17, 3637-3658, 10.5194/acp-17-3637-2017, 2017.
- Welti, A., Müller, K., Fleming, Z. L., and Stratmann, F.: Concentration and variability of ice nuclei in the subtropical maritime boundary layer, *Atmos. Chem. Phys.*, 18, 5307-5320, 10.5194/acp-18-5307-2018, 2018.
- 1140 Whale, T. F., Murray, B. J., O'Sullivan, D., Wilson, T. W., Umo, N. S., Baustian, K. J., Atkinson, J. D., Workneh, D. A., and Morris, G. J.: A technique for quantifying heterogeneous ice nucleation in microlitre supercooled water droplets, *Atmos. Meas. Tech.*, 8, 2437-2447, 10.5194/amt-8-2437-2015, 2015.
- Yaalon, D. H., and Ganor, E.: East Mediterranean trajectories of dust-carrying storms from the Sahara and Sinai, in: *Saharan Dust: Mobilization, Transport, Deposition*, edited by: Morales, C., John Wiley and Sons New York, 187-193, 1979.
- 1145 Zaitchik, B. F., Evans, J. P., Geerken, R. A., and Smith, R. B.: Climate and vegetation in the Middle East: Interannual variability and drought feedbacks, *Journal of Climate*, 20, 3924-3941, 10.1175/JCLI4223.1, 2007.
- Zittis, G., Almazroui, M., Alpert, P., Ciais, P., Cramer, W., Dahdal, Y., Fnais, M., Francis, D., Hadjinicolaou, P., Howari, F., Jrrar, A., Kaskaoutis, D. G., Kulmala, M., Lazoglou, G., Mihalopoulos, N., Lin, X., Rudich, Y., Sciare, J., Stenchikov, G., Xoplaki, E., and Lelieveld, J.: Climate Change and Weather Extremes in the Eastern Mediterranean and Middle East, 60, e2021RG000762, <https://doi.org/10.1029/2021RG000762>, 2022.
- 1150