



Optimized procedure for the determination of alkylamines in airborne particulate matter of anthropized areas

5 Davide Spolaor¹, Lidia Soldà¹, Gianni Formenton², Marco Roverso¹, Denis Badocco¹,
Sara Bogialli¹, Fazel A. Monikh^{1,3}, Andrea Tapparo¹

¹Dipartimento di Scienze Chimiche, Università di Padova, via Marzolo 1, 35131 Padova, Italy

²ARPAV Environmental Regional Agency, Laboratory Department, via Lissa 6, 30171
Mestre, Venice, Italy

10 ³Institute for Nanomaterials, Advanced Technologies and Innovation, Technical University of
Liberec Bendlova 1409/7, 460 01, Liberec, Czech Republic

Correspondence: Andrea Tapparo (andrea.tapparo@unipd.it)

15 **Abstract**

Due to their role in the formation of secondary aerosol, the concentrations of the most abundant
aliphatic amines (methylamine (MA), dimethylamine (DMA), ethylamine (EA), diethylamine
(DEA), propylamine (PA), and buthylamine) present in the aerosol of a very anthropized area
were measured by an optimized analytical procedure. PM₁₀ samples were collected in the
20 tannery district of Vicenza (in the Po Valley, northern Italy) in autumn 2020. Alkylamines were
extracted in water and converted to carbamates through derivatization with Fmoc-OSu for
subsequent determination by UHPLC with fluorescence detection. The procedure has been
optimized and validated obtaining very satisfactory analytical performances: LODs were in the
range of 0.09-0.26 ng/m³, average uncertainty of 3.4 % and recoveries of 95-101 %. The mean
25 total concentration of the 6 amines measured in this study was 37 ± 17 ng/m³, with DMA
making the largest contribution. The proposed procedure may contribute to a better
characterization of the local aerosol. In our preliminary investigation, the Pearson's correlation
test showed that amines correlate strongly with each other and with secondary inorganic ions
(NH₄⁺, NO₃⁻ and SO₄²⁻), confirming that they compete with ammonia in the acid-base
30 atmospheric processes that lead to the formation of nitrate and sulphate particles. The developed
method allows to gather critical information about the load of aliphatic amines in PM to gain
more insights into the sources and fate of these chemicals in the atmosphere.



1. Introduction

35

Atmospheric particulate matter (PM) is the subject of many discussions about the health of humans and ecosystems. A polluted atmosphere contains hundreds of organic and inorganic compounds involved in the formation of secondary aerosols and that contribute to their toxicity. Atmospheric amines could be especially important due to their acid neutralizing capacity. These amines are derivatives of ammonia with one or more hydrogen atoms replaced by an alkyl or aryl group. Like ammonia, they are hydrophilic basic compounds that exist in the atmosphere in both gaseous and particulate phases. Amines could undergo rapid acid-base reactions with acids such as H_2SO_4 and HNO_3 , similar to the transformation that occurs for ammonia itself (Ge et al., 2011b). They can also contribute to the production of secondary organic aerosol (SOA) by reacting with organic acids and oxidants (e.g. O_3 or $\cdot\text{NO}_3$) (Tang et al., 2013; Nielsen et al., 2012; Malloy et al., 2009). Recent studies have highlighted that atmospheric amines, in particular dimethylamine, could promote the formation of new particles (NPF) (Loukonen et al., 2010; Almeida et al., 2013; Yao et al., 2018; Xiao et al., 2021). Most amines are toxic and irritants to the eyes, skin, and mucous membrane and some of their oxidation products can also be carcinogenic (e.g. nitrosamines) (Lee and Wexler, 2013). The most common organic amines found in the atmosphere are C1-C6 alkylamines, which can be emitted from a variety of anthropogenic and natural sources (well recognized in the review of Ge et al. (2011a)). The main natural sources include biomass burning and biodegradation. Animal husbandry is one of the main anthropogenic sources of atmospheric amines, along with various industrial activities, in particular the food industry. Other anthropogenic sources include fuel combustion (Shen et al., 2017), vehicular traffic, composting, the treatment of industrial waste, and the consumption of tobacco (Ge et al., 2011a). Tanneries may also represent a significant source (JRC reference reports, 2013). In urban areas anthropogenic nonagricultural sources could represent the main contribution to amines emission (Shen et al., 2017; Cheng et al., 2020; Liu et al., 2022). Furthermore, emerging emission sources of amines have been associated with their use in the CO_2 capture processes and, possibly, in catalytic processes for the reduction of NO_x emissions from engines (Ge et al., 2011a; Muthiya et al., 2022; Stanciulescu et al., 2010)

Measurement of the ambient concentrations of atmospheric amines is challenging due to their high volatility and polarity. Common techniques for the determination of low molecular weight alkylamine are gas chromatography (GC), high-performance liquid chromatography (HPLC), and ion chromatography (IC), coupled with different detectors. In direct analysis by IC the

65



separation and quantification of all analytes is often difficult due to interference from other cations (e.g., sodium or potassium) and the difficulty of separate some of the amines (e.g., diethylamine and trimethylamine) (VandenBoer et al., 2012; Huang et al., 2014; Zhou et al., 2018; Feng et al., 2020). Similarly, direct analysis by HPLC-MS may fail to separate amines with the same molecular mass (i.e. DMA/EA and PA/TMA) (Kieloaho et al. 2013; Tzitzikalaki et al., 2021). Direct analyses of amines can also be performed using GC (Bai et al., 2019). However, the polar nature of small aliphatic amines makes these analyses unfavorable due to the adsorption and decomposition of the solute in the column, resulting in peak tailing and analyte losses.

Because of the poor results often obtained with direct analyses of amines, the derivatization of the target analytes is recommended. Organic amines can be derivatized with various methods such as acylation, carbamate formation, Schiff base formation and silylation (Płotka-Wasyłka et al., 2015). Among these methods, acylation and carbamate formation are the ones most commonly used because they allow a simple and fast derivatization step, where amines are easily derivatized in aqueous solutions and mild conditions. The derivatization is usually followed by a GC or HPLC separation. Derivatization can also be used to add a chromophore (or a fluorophore) to derivatized analytes, allowing the detection by UV-vis absorption or fluorescence (Płotka-Wasyłka et al., 2015; Szulejko and Kim, 2014). In this work a simple and sensitive analytical method for the determination of aliphatic amines contained in atmospheric PM has been developed and validated. It makes use of a water extraction of amines, their derivatization by Fmoc-OSu (9-fluorenylmethoxycarbonyl-N-hydroxysuccinimide), and their instrumental analysis by UHPLC with fluorescence detection (FLD). The procedure has been applied for the determination of six volatile aliphatic amines (methylamine (MA), dimethylamine (DMA), ethylamine (EA), diethylamine (DEA), propylamine (PA), and butylamine (BA)) contained (as ammonium salts) in PM₁₀ samples of a highly anthropized area. The results are also preliminarily discussed to assess the contribution of aliphatic amines in the formation of secondary aerosol.

95

2. Materials and Methods

2.1 Chemicals

Ultrapure water (18.2 MΩ cm, ELGA LabWater PURELAB Chorus 1 Complete) and 37% HCl solution (ACS reagent, Riedel-de Haën) were used for the extraction of the analytes; Fmoc-

100



OSu (9-fluorenylmethoxycarbonyl-N-hydroxysuccinimide, Sigma Aldrich), acetonitrile (>99,95%, Carlo Erba), sodium tetraborate (Sodium tetraborate decahydrate, Normapur) and NaOH (>97%, ACS reagent, Sigma Aldrich) were used for the derivatization step. Methylamine (MA), dimethylamine (DMA), ethylamine (EA), diethylamine (DEA), propylamine (PA) were purchased as hydrochloride salt (>98%, Sigma Aldrich) while buthylamine was purchased as pure liquid (99.5%, Sigma Aldrich).

2.2 Aerosol Sampling

The sampling campaign was carried out in the Chiampo Valley (Po Valley, northern Italy), an area located near Vicenza and characterized by the presence of numerous leather industries (tanning district). PM₁₀ samples were collected for 24 h and measured (beta attenuation monitoring) using a SWAM 5a Monitor (FAI Instruments, Italy) working at a constant flow rate of 2.3 m³/h and equipped with 47 mm quartz filters (Whatman, QM-A quartz filters, Particle Retention Rating 98% at 2.2 μm). This sampling and measurement procedure (implemented in the air quality monitoring network of the Environmental Regional Agency, ARPAV) is the standardized “Ambient Air - Automated measuring system for the measurements of the concentration of particulate matter (PM₁₀; PM_{2.5})” (EN 16450:2017). It is worth noticing that artifacts can occur in the sampling of aliphatic amines, well described by Shen et al. (2017) although different hypotheses regarding DEA are proposed. We did not estimate these artifacts, but regarding sampling campaigns carried out in the cold season, the sampling losses of volatile and semivolatile compounds were probably limited. The sampling campaign was conducted in three different areas of the Chiampo Valley over a two-month period, from the 18th of September to the 7th of October 2020 (site of Montebello), from the 13th of October to the 28th of October 2020 (site of Montorso) and from the 30th of October to the 18th of November 2020 (site of Trissino). A total of 53 PM₁₀ samples were collected. Filter samples were kept at -20 °C until further analysis. Analysis of water-soluble ions (by ion chromatography), organic carbon, and elemental carbon (by Sunset OC/EC analyzer) was also performed using well-established procedures (Giorio et al., 2022; Khan et al., 2016).

2.3 Sample Preparation

Analytes were extracted from a portion of the filter (2/3) with 3 mL of a 0.1 M HCl solution in ultrapure water followed by sonication for 10 minutes at a temperature below 25 °C. The extract was then filtered (PTFE membrane, 0.2 μm pore size) inside a 1.5 mL test tube and stored for further analysis in a freezer at -18 °C. The derivatization reagent was prepared by dissolving a



135 few mg of Fmoc-OSu in acetonitrile to obtain a concentration of 3 g/L. Borate buffer was
prepared by dissolving 5 g of sodium tetraborate in 100 mL of milliQ water. An aliquot of 300
μL of the sample solution was added in a 1.5 mL test tube and neutralized by adding 5 μL of a
6 M NaOH solution. Then 150 μL of borate buffer and 250 μL of derivatization reagent were
added to the solution. The mixture was stirred for about 30 seconds and allowed to react for 10
140 minutes at 50 °C.

2.4 Instrumental analysis

The solution containing the derivatized analytes was injected (injection volume 5 μL) into an
HPLC (LC-20AD XR Shimadzu; Canby, OR, USA) equipped with a fluorescence detector (RF-
145 20° XS Shimadzu) and an autosampler (SIL-20AC XR Shimadzu). Excitation and emission
wavelengths were 264 and 313 nm, respectively. The separation was performed using a Luna
Omega 1.6 μm Polar C18 column (100 x 2.1 mm), with a constant flow rate of 0.2 mL/min and
a column temperature of 30°C. A gradient elution program using acetonitrile (eluent B) and
water (eluent A), to which formic acid was added (0.05%), was optimized and used to separate
150 the derivatized amines. The elution program started with a mixture containing eluent B at 40%,
after a minute the concentration of B was increased to 60% in the span of 15 minutes, the elution
mixture was kept constant for 5 minutes then the concentration of B was raised to 100%.
Concentrated standard solutions (1 g/L) were prepared by dissolving the hydrochloride salt of
the amines in ultrapure water or, in the case of butylamine, by diluting the pure amine. From
155 these solutions, a mixed working standard solution containing all amines was then prepared,
each at 10 mg/L. This solution was diluted to obtain calibration solutions at suitable
concentrations (10, 50, 100, 250 μg/L). These standards were analyzed in duplicate to obtain
the calibration functions. Laboratory blanks consisting of ultrapure water and matrix blanks
consisting of aqueous extract of blank quartz filters were treated and analyzed each day as
160 regular samples to control the contribution of both chemicals and filters to the analytical signals.
The real sample results were corrected for the matrix blank concentrations.

2.5 Analysis via ESI-QTOF-MS

The possible presence of chromatographic interferences was investigated by analyzing PM10
165 samples using independent detection techniques. Specifically, three selected PM10 samples
were analyzed using the same sample preparation procedure and the same chromatographic
method described in 2.3 and 2.4, but with two different detection systems: fluorescence and
ESI-QTOF-MS. The UHPLC-MS analysis was performed with an ESI source, operating in



positive acquisition mode. Source parameters were: 300 °C capillary temperature, 40 a.u. sheath
170 gas flow, 300 °C sheath gas temperature, 20 a.u. auxiliary gas flow, 350 °C auxiliary gas
temperature, 4 kV spray voltage. Data were acquired in the range of 100-1000 m/z (scan rate).

2.6 Statistical analysis

The correlations among the data acquired during the campaign (temperature, relative humidity,
175 atmospheric precipitation, concentration of atmospheric pollutants - NO_x, H₂S and NH₃ - Table
S1) and the measured concentrations of PM10 components (elemental and organic carbon,
inorganic ions and amines; Table S2, S3 and S4) were evaluated by the Pearson's correlation
coefficients. Principal component analysis (PCA) was also performed considering the full set
of PM10 samples and species analyzed. Atmospheric parameters such as temperature, relative
180 humidity, and precipitation quantities were used as supplementary variables.

3. Results and Discussion

185 3.1 Optimization of the analytical procedure

The aliphatic amines in the atmospheric particulate matter were determined here by extracting
the analytes from the sampling filters and subsequently analyzing by UHPLC-FLD, after a
derivatization procedure using Fmoc-OSu. The advantages of this method consist primarily of
its rapidity and simplicity; the amines are readily derivatized in an aqueous alkaline solution
190 with the addition of Fmoc-OSu and are directly injected into the HPLC column without the
need for further extraction steps.

It is worth noting that the derivatization step is generally mandatory for the separation of amines
via HPLC or GC (Huang et al., 2014) and most derivatizing agents are suitable only for primary
and secondary amines, thus precluding the analysis of tertiary amines. Nevertheless, the
195 derivatization of TMA is possible using Fmoc but the reaction is very slow (Szulejko et al.,
2014), especially in aqueous solutions, which is why no peak attributable to TMA were
identified in our chromatograms and no difference in the peak intensity of DMA were observed
after adding TMA to a standard solution (tertiary amines derivatized with Fmoc may dealkylate
yielding a product identical to the derivatized secondary amine).

200

3.1.1 Optimization of chromatographic conditions. Three RP-C18 columns were
preliminary tested for the UHPLC separation of the six derivatized aliphatic amines: Shimadzu



XR-ODS III C18, Phenomenex Luna Omega polar C18, Phenomenex Kinetex C18. The performances of the three columns are very similar in terms of selectivity, being able to separate the analytes in about twenty minutes using an acidic mixture of water/acetonitrile as the eluent. The Luna Omega column was finally chosen because of its slightly higher efficiency. The elution program (gradient elution using acidic water/acetonitrile) has been optimized to allow the proper separation of each amine, in particular the ethylamine/dimethylamine and butylamine/diethylamine pairs. The correct separation of methylamine was also problematic due to the presence of a large peak attributable to the by-products of Fmoc-OSu and to the derivatized NH₃, which are both present in the solution in large excess. The addition of formic acid in the two solvents (0.05%) did not have an effect on the retention time of the amines, but slightly reduced the retention time of the Fmoc by-products. Changes in the chromatographic method were aimed at reducing the elution time without reducing the separation efficiency. Figure 1 shows representative chromatograms obtained for a standard solution (100 µg/L) and a PM10 sample extract. The order of retention times of Fmoc derivatives is as follows: methylamine (MA), ethylamine (EA), dimethylamine (DMA), propylamine (PA), butylamine (BA), and diethylamine (DEA); the order is the same as that reported in other articles (Shen et al., 2017; Cheng et al., 2020).

220

3.1.2 Optimization of extraction and derivatization steps. Compared to the procedures of Shen et al. (2017) and Cheng et al. (2020) we added a larger amount of derivatizing agent (250 µL in 700 µL instead of just 400 µL in 10 mL of Fmoc-OSu in acetonitrile) obtaining stable peaks for each of the six target amines. Therefore, the extraction and reaction steps (time, temperature, and reaction mixture composition) were optimized to gain the maximum recovery for each of the analytes. The extraction was initially tested with ultrapure water, but the addition of hydrochloric acid (0.1 M) showed better results in terms of recoveries (from 80-95% to >95%).

The extraction time was initially set at 20 min, but no significant differences in instrumental signals were observed after reducing it to 10 min. The derivatization step was tested at different temperatures (25, 40, 50, 60 °C) obtaining equal reaction times between the various tests. The reaction was indeed very fast even at ambient temperature, as it takes less than 5 minutes to complete.

235




Figure 1. UHPLC-FLD chromatograms of (a) standard solution of the six amines at 100 µg/L each and (b) a PM10 sample extract (sample of 02.10.2021, PM10=24.5 µg/m³)

However, because the reaction time and temperature affected the amount of unreacted Fmoc in the solution (that may represent a serious chromatographic interference), the reaction was
240 carried out at 50 °C, for 10 min, to eliminate the chromatographic peak relative to the unreacted Fmoc that coelutes with the derivatized ethylamine. The same results were also obtained using Fmoc-Cl as the derivatizing agent. In conclusion, if compared with most previous published procedures that use a derivatization reagent (Shen et al., 2017; Cheng et al., 2020; Choi et al., 2020; Akyüz, 2008; Majedi and Lee, 2017; Parshintsev et al., 2015), our sample pre-treatment
245 appears faster as it does not need successive liquid-liquid extraction or preconcentration steps.



3.2 Method validation

The linearity of the calibration functions, limits of detection (LOD), precision (repeatability), and recoveries of the method were estimated. The calibration functions of the six aliphatic amines (ranged from 0 $\mu\text{g/L}$, the solvent blank, to 250 $\mu\text{g/L}$) were all assessed as linear after an F test between the linear and quadratic regression variance ($\alpha = 0.05$; $F_{(0.05,8,7)} = 3.73$; $F_{sp} < 1.85$; $p > 0.2$). Repeatability was evaluated as the relative standard deviation (RSD,%) over four equal parts of the same sample (a PM10 filter), each spiked with 15 μL of a mixture containing the six amines at 10 mg/L . The measured RSD were less than 5% for all the amines. The recoveries were evaluated by adding different quantities of a mixture containing each of six amines at 10 mg/L (0, 20, 40, 60 μL) to four equal parts of the same PM10 sample and determining the slope of each recovery function. It was found that the recoveries ranged from 95% to 101%. The limits of detection (LOD) were estimated from the calibration function using the Hubaux-Vos method (Hubaux and Vos, 1970), obtaining values ranging from 1.1 to 3.2 $\mu\text{g/L}$, which correspond to an atmospheric concentration of 0.088-0.26 ng/m^3 . These validation parameters are listed in Table 1.

These performances, compared to the ones obtained by previously developed methods (Huang et al., 2014; Choi et al., 2020; Parshintsev et al., 2015), are characterized by similar or higher LODs, mainly due to lower sampling volume and the absence of a preconcentration step, but also by better accuracy, with quantitative recovery functions, and good repeatability. Furthermore, our optimized sample preparation procedure allows for the correct quantification of DEA that was not carried out using analogous methods (Shen et al., 2017; Cheng et al., 2020).

Table 1 Detection limit, recovery, and relative standard deviation of the analytical method developed in this study

Analyte	Detection limit ($\mu\text{g/L}$)	Detection limit (ng/m^3)	RSD (%)*	Recovery (%)
Methylamine	2.79	0.23	3.89	100.7 \pm 6.0
Ethylamine	1.72	0.14	3.54	99.1 \pm 3.9
Dimethylamine	2.45	0.20	4.72	98 \pm 16
Propylamine	1.24	0.10	3.58	100.4 \pm 2.3
Buthylamine	3.16	0.26	3.19	95.2 \pm 2.0
Diethylamine	1.07	0.09	1.56	97.0 \pm 6.5

*estimated by the analysis of a real PM10 sample (spiked, added concentration 4.1 ng/m^3 of each amine, $n=4$)



3.2.1 Comparison of the results obtained by independent methods. Three samples were analyzed with two different methods as described in section 2.5. In the MS analysis, the signal of the parent ion was used to identify each of the derivatized amines, but it was too low to be used for quantification. As a consequence, the MS signal at m/z 179.0861 (common to each
275 amine, due to the in-source fragmentation of the derivatized analytes and attributable to a residual moiety of the derivatizing agent) was selected for the quantitative MS analysis. Butylamine was used as the internal standard, as it was not present in the analyzed samples. To ensure that each amine was present at a concentration high enough to be quantified, 10 μL of a mixture containing the amines at 10 mg/L was added to the three samples. From the comparison
280 of the results obtained by applying the two different methods (Table S5), it was found that the measured concentrations of the amine derivatives were statistically equivalent and that there was no significant systematic difference between the two detection methods. This result suggests the absence of significant chromatographic interferences in the optimized UHPLC-FLD analytical method.

285

3.3 Environmental concentrations of aliphatic amines

Table 2 shows the mean concentrations of the amines contained in PM10 samples collected in the three sampling sites considered in this study (Montebello, Montorso, Trissino, Figure S1). Taking into account the average concentration (\pm standard deviation) throughout the sampling
290 campaign (53 total samples), the amines present at higher concentrations were dimethylamine ($19.0 \pm 8.7 \text{ ng/m}^3$), methylamine ($9.7 \pm 5.5 \text{ ng/m}^3$) and diethylamine ($4.5 \pm 2.0 \text{ ng/m}^3$) while butylamine was measured only at concentrations below the detection limit and therefore was not considered in the present discussion. Absolute concentrations in the three sites of EA, PA, and DEA varied very little, while concentrations of MA and DMA were characterized by much
295 wider distributions. The concentration of most of the amines did not show great variations between the three locations considered, except for dimethylamine, where the average concentration ranged from 13.7 ng/m^3 in Montebello to 24.9 ng/m^3 in Trissino (details in the box plots, Fig. S2). The concentrations of aliphatic amines measured in this work are consistent with those described in other studies (Choi et al., 2020; Akyüz, 2008; Feng et al., 2020; Cheng
300 et al., 2020), being in the range between a few tenths of $\mu\text{g/m}^3$ and 1 ng/m^3 . In the work of Choi et al. (2020), the concentration of amines (EA, DMA, DEA, PA, BA) was measured in PM2.5 in the metropolitan area of Seoul (South Korea). The sum of the annual mean concentrations of the target amines was $5.56 \pm 2.76 \text{ ng/m}^3$ with a low dependence on the sampling season. Consistent with our study, the main contribution was attributed to DMA, while



305 the lowest was attributed to EA. However, the concentrations measured in Seoul were much
lower than those measured in the Chiampo Valley (Vicenza-IT, present study). Cheng et al.
(2020) measured EA, DMA, and DEA in Yangzhou, China. They obtained much higher values
for EA, $12.58 \pm 10.45 \text{ ng/m}^3$, but lower for the other two amines. Akyüz (2008) measured
amines in PM₁₀ sampled in the Zonguldak industrial area in Turkey, where pollution is very
310 high due to the production and consumption of coal. Concentrations in the Zonguldak industrial
area were rather high for all amines considered ($7\text{-}9 \text{ ng/m}^3$) and higher during winter. In the
work of Feng et al. (2020), which was carried out in Shanghai, the amines present at the highest
concentrations were found to be DMA and MA, which reached average concentrations around
 20 ng/m^3 . The differences between the measured concentrations in this study and those reported
315 in literature are probably due to the different emission sources present in the Chiampo Valley
compared to the other locations. In particular, DMA appears to be present in large quantities in
the Chiampo Valley compared to other amines, while BA was almost not detected ($<1 \text{ ng/m}^3$).
The high total concentration of amines can be attributed to the presence of significant emission
sources, such as vehicular exhaust or intense industrial activity (tanning industry, in particular)
320 and to the accumulation of pollutants that affects the entire Po Valley during winter and autumn
(Masiol et al., 2015).

Table 2 Amines average concentrations and standard deviations in the three sites and during the entire campaign (53 total samples)

	Montebello (n=19)		Montorso (n=15)		Trissino (n=19)		Total (N=53)	
	mean (ng/m ³)	SD (ng/m ³)	mean (ng/m ³)	SD (ng/m ³)	mean (ng/m ³)	SD (ng/m ³)	mean (ng/m ³)	SD (ng/m ³)
MA	18.5	5.2	10.9	5.9	11.0	5.0	9.7	5.5
EA	2.6	1.2	2.5	1.4	1.95	0.76	2.4	1.2
DMA	14.2	5.6	19.4	9.3	24.9	7.2	19.0	8.7
PA	1.5	1.1	1.8	1.5	2.1	1.2	1.8	1.2
DEA	4.3	2.1	5.5	2.2	4.0	1.5	4.5	2.0

325

3.4 Correlation among amine concentrations and other atmospheric variables

Regarding the contribution that this method can offer in the study of atmospheric phenomena,
in our preliminary study we observed that the amines contained in PM₁₀ samples correlate
330 rather strongly between themselves (Pearson's correlation coefficients are reported Table S6),
indicating that their presence in PM is due to similar processes, i.e. the rapid acid-base reactions
with atmospheric acids as described in previous works (Ge et al., 2011a). The concentration of



all amines correlates strongly or moderately with PM₁₀ ($r > 0.55$, except ethylamine $r = 0.38$), as expected for the aerosol components associated with the secondary fraction (that is significant
335 in Po Valley). In effect, amines are also strongly correlated with nitrate ($r > 0.52$) and ammonium ($r > 0.51$) supporting the hypothesis that amines are present in the particulate mainly as a result of acid-base processes that lead to the formation of nitrates and, in this case, to a lesser extent, of sulphates.

Regarding the sources of emission, it is worth noting that DMA (the amine present at higher
340 concentrations) shows a strong correlation with potassium ($r = 0.73$) supporting the hypothesis that biomass combustion processes are significant sources of amines. This hypothesis is further confirmed by the strong correlation ($r = 0.80$) between DMA and OC concentration in PM₁₀, taking into account that both OC and potassium in PM are closely associated with biomass burning emissions. However, OC could also be related to industrial emission sources, which
345 then would also contribute to the emission of DMA. On the other hand, the moderate correlation ($r = 0.51$) between DMA and calcium is not immediately justifiable. It is worth noticing that these correlations are not exactly reproduced for the other amines. For instance, the lower correlation of EA with both DMA and PM₁₀, also confirmed by PCA results (see below), is not easily explainable. A hypothesis regarding a specific source of EA (possibly associated to the
350 traffic combustion), different from those that characterize other amines (biomass combustion, tanneries) is of great interest and needs more investigations. To provide more insights into the sources of amines more studies on the ambient concentrations of amines, both in PM and in the air, with source apportionment techniques, are still needed.

The PCA loading plot depicted in Figure 2 shows that PC1, which describes 53.22% of the total
355 variance, is linked to the fine component of the atmospheric particulate, especially to the component of secondary origin, which includes all the amines. PC2, which expresses 14.98% of the variance, appears to be more related to the coarse component of the particulate matter. The loading plot confirms the strong correlation between organic carbon, PM₁₀, nitrate, and sulfate, as well as with different amines, in particular with DMA and MA. The graph also shows
360 that the gaseous substances (NO_x, NH₃, i.e. the precursors of the secondary compounds) do not correlate at all with the secondary components of the particulate. This result indicates that the concentration of these pollutants does not significantly affect the local concentration or the composition of PM₁₀ on a daily basis, which is influenced mainly by meteorological parameters and primary emission sources (Masiol et al., 2017).

365




Figure 2. PCA loading plot referred to the first two principal components. Blu vectors indicate active variables and red vectors refer to supplementary variables.

370

4. Conclusions

The optimization of a fast sample preparation procedure followed by a UHPLC-FLD analysis allowed us to accurately determine the main alkylamines in the aerosol samples. The proposed analytical method performs well in terms of precision, detection limit, recoveries, and linear response range. The concentrations of amines measured in PM10 sampled in the tannery district (Chiampo Valley, Italy) are quite high compared to the ones quantified in other works, possibly due to the presence of important local emission sources, such as vehicular traffic, intense industrial activity, and biomass burning. These sources may have contributed, in particular, to the production of dimethylamine. Amine concentrations are strongly correlated with PM10 and with secondary inorganic ions, supporting the hypothesis that amines are present in the particulate mainly as a result of acid-base processes that lead to the formation of nitrates and sulphates. Dimethylamine, which is the amine present at higher concentrations, shows strong correlations with both the organic component of PM10 and potassium, indicating the relevance

380



385 of biomass combustion process emissions. We expect that the procedure developed here will
find application in extensive environmental studies. In this connection, a promising perspective
for the UHPLC-FLD method is the possibility to validate the analytical data obtained by the
online aerosol analyzer (ATOF-MS, AMS) (Malloy et al., 2009; Bottenus et al., 2018; Huang
et al., 2012; Giorio et al., 2015).

390

Author contributions

DS and AT are the principal investigators; they are responsible for all research activities. LS
supported the laboratory analysis of PM samples. GF coordinated the sampling campaigns and
395 performed the OC/EC analysis. MR and SB were responsible for LC-MS analysis. DB was
involved in the statistical analysis of the results. FAM contributed to the interpretation of the
data and to the text revision.

Acknowledgements

400 We acknowledge the support from ARPA Veneto, Dept. of Vicenza (IT), in the PM10 sampling
campaign.

Competing interests

The authors declare that they have no conflict of interest.

405



References

- Akyüz, M.: Simultaneous determination of aliphatic and aromatic amines in ambient air and airborne particulate matters by gas chromatography-mass spectrometry, *Atmos. Environ.*, 42, 3809–3819, <https://doi.org/10.1016/j.atmosenv.2007.12.057>, 2008.
- 410 Almeida, J., Schobesberger, S., Kürten, A., Ortega, I. K., Kupiainen-Määttä, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T., Kvashin, A.
- 415 N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppä, J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara, P.,
- 420 Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H., and Kirkby, J.: Molecular understanding of sulphuric acid-amine particle nucleation in the atmosphere, *Nature*, 502, 359–363, <https://doi.org/10.1038/nature12663>, 2013.
- 425 Bai, J., Baker, S. M., Goodrich-Schneider, R. M., Montazeri, N., and Sarnoski, P. J.: Aroma Profile Characterization of Mahi-Mahi and Tuna for Determining Spoilage Using Purge and Trap Gas Chromatography-Mass Spectrometry, *J. Food Sci.*, 84, 481–489, <https://doi.org/10.1111/1750-3841.14478>, 2019.
- Bottenus, C. L. H., Massoli, P., Sueper, D., Canagaratna, M. R., VanderSchelden, G., Jobson, 430 B. T., and VanReken, T. M.: Identification of amines in wintertime ambient particulate material using high resolution aerosol mass spectrometry, *Atmos. Environ.*, 180, 173–183, <https://doi.org/10.1016/j.atmosenv.2018.01.044>, 2018.
- Cheng, G., Hu, Y., Sun, M., Chen, Y., Chen, Y., Zong, C., Chen, J., and Ge, X.: Characteristics and potential source areas of aliphatic amines in PM_{2.5} in Yangzhou, 435 China, *Atmos. Pollut. Res.*, 11, 296–302, <https://doi.org/10.1016/j.apr.2019.11.002>, 2020.
- Choi, N. R., Lee, J. Y., Ahn, Y. G., and Kim, Y. P.: Determination of atmospheric amines at Seoul, South Korea via gas chromatography/tandem mass spectrometry, *Chemosphere*, 258, 127367, <https://doi.org/10.1016/j.chemosphere.2020.127367>, 2020.
- Feng, H., Ye, X., Liu, Y., Wang, Z., Gao, T., Cheng, A., Wang, X., and Chen, J.: 440 Simultaneous determination of nine atmospheric amines and six inorganic ions by non-suppressed ion chromatography using acetonitrile and 18-crown-6 as eluent additive, *J. Chromatogr. A*, 1624, 461234, <https://doi.org/10.1016/j.chroma.2020.461234>, 2020.
- Ge, X., Wexler, A. S., and Clegg, S. L.: Atmospheric amines - Part I. A review, *Atmos. Environ.*, 45, 524–546, <https://doi.org/10.1016/j.atmosenv.2010.10.012>, 2011a.
- 445 Ge, X., Wexler, A. S., and Clegg, S. L.: Atmospheric amines - Part II. Thermodynamic properties and gas/particle partitioning, *Atmos. Environ.*, 45, 561–577, <https://doi.org/10.1016/j.atmosenv.2010.10.013>, 2011b.
- Giorio, C., D’Aronco, S., Di Marco, V., Badocco, D., Battaglia, F., Soldà, L., Pastore, P., and Tapparo, A.: Emerging investigator series: aqueous-phase processing of atmospheric



- 450 aerosol influences dissolution kinetics of metal ions in an urban background site in the Po Valley, *Environ. Sci. Process. Impacts*, 24, 884–897, <https://doi.org/10.1039/d2em00023g>, 2022.
- Giorio, C., Tapparo, A., Dallosto, M., Beddows, D. C. S., Esser-Gietl, J. K., Healy, R. M., and Harrison, R. M.: Local and regional components of aerosol in a heavily trafficked street
455 canyon in central London derived from PMF and cluster analysis of single-particle ATOFMS spectra, *Environ. Sci. Technol.*, 49, 3330–3340, <https://doi.org/10.1021/es506249z>, 2015.
- Huang, R. J., Li, W. B., Wang, Y. R., Wang, Q. Y., Jia, W. T., Ho, K. F., Cao, J. J., Wang, G. H., Chen, X., Ei Haddad, I., Zhuang, Z. X., Wang, X. R., Prévôt, A. S. H., O’Dowd, C. D.,
460 and Hoffmann, T.: Determination of alkylamines in atmospheric aerosol particles: A comparison of gas chromatography-mass spectrometry and ion chromatography approaches, *Atmos. Meas. Tech.*, 7, 2027–2035, <https://doi.org/10.5194/amt-7-2027-2014>, 2014.
- Hubaux, A. and Vos, G.: Decision and Detection Limits for Linear Calibration Curves, *Anal. Chem.*, 42, 849–855, <https://doi.org/10.1021/ac60290a013>, 1970.
- Huang, Y., Chen, H., Wang, L., Yang, X., and Chen, J.: Single particle analysis of amines in ambient aerosol in Shanghai, *Environ. Chem.*, 9, 202–210, <https://doi.org/10.1071/EN11145>, 2012.
- JRC, Best Available Techniques (BAT) Reference Document for the Tanning of Hides and
470 Skins, Report EUR 26130 EN, 290 pp., <https://doi.org/10.2788/13548>, 2013.
- Khan, M. B., Masiol, M., Formenton, G., Di Gilio, A., de Gennaro, G., Agostinelli, C., and Pavoni, B.: Carbonaceous PM_{2.5} and secondary organic aerosol across the Veneto region (NE Italy), *Sci. Total Environ.*, 542, 172–181, <https://doi.org/10.1016/j.scitotenv.2015.10.103>, 2016.
- 475 Kieloaho, A. J., Hellén, H., Hakola, H., Manninen, H. E., Nieminen, T., Kulmala, M., and Pihlatie, M.: Gas-phase alkylamines in a boreal Scots pine forest air, *Atmos. Environ.*, 80, 369–377, <https://doi.org/10.1016/j.atmosenv.2013.08.019>, 2013.
- Lee, D. Y. and Wexler, A. S.: Atmospheric amines - Part III: Photochemistry and toxicity, *Atmos. Environ.*, 71, 95–103, <https://doi.org/10.1016/j.atmosenv.2013.01.058>, 2013.
- 480 Liu, Z., Li, M., Wang, X., Liang, Y., Jiang, Y., Chen, J., Mu, J., Zhu, Y., Meng, H., Yang, L., Hou, K., Wang, Y., and Xue, L.: Large contributions of anthropogenic sources to amines in fine particles at a coastal area in northern China in winter, *Sci. Total Environ.*, 839, 156281, <https://doi.org/10.1016/j.scitotenv.2022.156281>, 2022.
- Loukonen, V., Kurtén, T., Ortega, I. K., Vehkamäki, H., Pádua, A. A. H., Sellegri, K., and Kulmala, M.: Enhancing effect of dimethylamine in sulfuric acid nucleation in the presence of water—a computational study, *Atmos. Chem. Phys.*, 10, 4961–4974, <https://doi.org/10.5194/acp-10-4961-2010>, 2010.
- 485 Majedi, S. M. and Lee, H. K.: Combined dispersive solid-phase extraction-dispersive liquid–liquid microextraction-derivatization for gas chromatography–mass spectrometric determination of aliphatic amines on atmospheric fine particles, *J. Chromatogr. A*, 1486, 86–95, <https://doi.org/10.1016/j.chroma.2016.06.079>, 2017.
- 490



- Malloy, Q. G. J., Qi, L., Warren, B., Cocker, D. R., Erupe, M. E., and Silva, P. J.: Secondary organic aerosol formation from primary aliphatic amines with NO₃ radical, *Atmos. Chem. Phys.*, 9, 2051–2060, <https://doi.org/10.5194/acp-9-2051-2009>, 2009.
- 495 Masiol, M., Benetello, F., Harrison, R. M., Formenton, G., De Gaspari, F., and Pavoni, B.: Spatial, seasonal trends and transboundary transport of PM_{2.5} inorganic ions in the Veneto region (Northeastern Italy), *Atmos. Environ.*, 117, 19–31, <https://doi.org/10.1016/j.atmosenv.2015.06.044>, 2015.
- 500 Masiol, M., Squizzato, S., Formenton, G., Harrison, R. M., and Agostinelli, C.: Air quality across a European hotspot: Spatial gradients, seasonality, diurnal cycles and trends in the Veneto region, NE Italy, *Sci. Total Environ.*, 576, 210–224, <https://doi.org/10.1016/j.scitotenv.2016.10.042>, 2017.
- 505 Muthiya, S. J., Natrayan, L., Yuvaraj, L., Subramaniam, M., Dhanraj, J. A., and Mammo, W. D.: Development of Active CO₂ Emission Control for Diesel Engine Exhaust Using Amine-Based Adsorption and Absorption Technique, *Adsorpt. Sci. Technol.*, 2022, <https://doi.org/10.1155/2022/8803585>, 2022.
- 510 Nielsen, C.J., D'Anna, B., Bossi, R., Bunkan, A.J.C., Dithmer, L., Glasius, M., Hallquist, M., Hansen, A.M.K., Lutz, A., Salo, K., Maguta, M.M., Nguyen, Q., Mikoviny, T., Müller, M., Skov, H., Sarrasin, E., Stenström, Y., Tang, Y., Westerlund, J., Wisthaler, A.: Atmospheric Degradation of Amines (ADA), Summary report from atmospheric chemistry studies of amines, nitrosamines, nitramines and amides, CLIMIT project no. 201604, 2012.
- 515 Parshintsev, J., Rönkkö, T., Helin, A., Hartonen, K., and Riekkola, M. L.: Determination of atmospheric amines by on-fiber derivatization solid-phase microextraction with 2,3,4,5,6-pentafluorobenzyl chloroformate and 9-fluorenylmethoxycarbonyl chloride, *J. Chromatogr. A*, 1376, 46–52, <https://doi.org/10.1016/j.chroma.2014.12.040>, 2015.
- Plotka-Wasyłka, J. M., Morrison, C., Biziuk, M., and Namieśnik, J.: Chemical Derivatization Processes Applied to Amine Determination in Samples of Different Matrix Composition, *Chem. Rev.*, 115, 4693–4718, <https://doi.org/10.1021/cr4006999>, 2015.
- 520 Shen, W., Ren, L., Zhao, Y., Zhou, L., Dai, L., Ge, X., Kong, S., Yan, Q., Xu, H., Jiang, Y., He, J., Chen, M., and Yu, H.: C1-C2 alkyl aminiums in urban aerosols: Insights from ambient and fuel combustion emission measurements in the Yangtze River Delta region of China, *Environ. Pollut.*, 230, 12–21, <https://doi.org/10.1016/j.envpol.2017.06.034>, 2017.
- Stanciulescu, M., Charland, J. P., and Kelly, J. F.: Effect of primary amine hydrocarbon chain length for the selective catalytic reduction of NO_x from diesel engine exhaust, *Fuel*, 89, 2292–2298, <https://doi.org/10.1016/j.fuel.2010.02.015>, 2010.
- 525 Szulejko, J. E. and Kim, K. H.: A review of sampling and pretreatment techniques for the collection of airborne amines, *TrAC - Trends Anal. Chem.*, 57, 118–134, <https://doi.org/10.1016/j.trac.2014.02.010>, 2014.
- 530 Tang, X., Price, D., Praske, E., Lee, S. A., Shattuck, M. A., Purvis-Roberts, K., Silva, P. J., Asa-Awuku, A., and Cocker, D. R.: NO₃ radical, OH radical and O₃-initiated secondary aerosol formation from aliphatic amines, *Atmos. Environ.*, 72, 105–112, <https://doi.org/10.1016/j.atmosenv.2013.02.024>, 2013.
- 535 Tzitzikalaki, E., Kalivitis, N., and Kanakidou, M.: Observations of gas-phase alkylamines at a coastal site in the east mediterranean atmosphere, *Atmosphere (Basel)*, 12, <https://doi.org/10.3390/atmos12111454>, 2021.



- VandenBoer, T. C., Markovic, M. Z., Petroff, A., Czar, M. F., Borduas, N., and Murphy, J. G.: Ion chromatographic separation and quantitation of alkyl methylamines and ethylamines in atmospheric gas and particulate matter using preconcentration and suppressed conductivity detection, *J. Chromatogr. A*, 1252, 74–83, <https://doi.org/10.1016/j.chroma.2012.06.062>, 2012.
- 540 Xiao, M., Hoyle, C. R., Dada, L., Stolzenburg, D., Kürten, A., Wang, M., Lamkaddam, H., Garmash, O., Mentler, B., Molteni, U., Baccharini, A., Simon, M., He, X. C., Lehtipalo, K., Ahonen, L. R., Baalbaki, R., Bauer, P. S., Beck, L., Bell, D., Bianchi, F., Brilke, S., Chen, D., Chiu, R., Dias, A., Duplissy, J., Finkenzeller, H., Gordon, H., Hofbauer, V., Kim, C., Koenig, T. K., Lampilahti, J., Lee, C. P., Li, Z., Mai, H., Makhmutov, V., Manninen, H. E., 545 Marten, R., Mathot, S., Mauldin, R. L., Nie, W., Onnela, A., Partoll, E., Petäjä, T., Pfeifer, J., Pospisilova, V., Quéléver, L. L. J., Rissanen, M., Schobesberger, S., Schuchmann, S., Stozhkov, Y., Tauber, C., Tham, Y. J., Tomé, A., Vazquez-Pufleau, M., Wagner, A. C., Wagner, R., Wang, Y., Weitz, L., Wimmer, D., Wu, Y., Yan, C., Ye, P., Ye, Q., Zha, Q., Zhou, X., Amorim, A., Carslaw, K., Curtius, J., Hansel, A., Volkamer, R., Winkler, P. M., 550 Flagan, R. C., Kulmala, M., Worsnop, D. R., Kirkby, J., Donahue, N. M., Baltensperger, U., El Haddad, I., and Dommen, J.: The driving factors of new particle formation and growth in the polluted boundary layer, *Atmos. Chem. Phys.*, 21, 14275–14291, <https://doi.org/10.5194/acp-21-14275-2021>, 2021.
- 555 Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S. B., Ehn, M., Paasonen, P., Sipilä, M., Wang, M., Wang, X., Xiao, S., Chen, H., Lu, Y., Zhang, B., Wang, D., Fu, Q., Geng, F., Li, L., Wang, H., Qiao, L., Yang, X., Chen, J., Kerminen, V. M., Petäjä, T., Worsnop, D. R., Kulmala, M., and Wang, L.: Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278–281, <https://doi.org/10.1126/science.aao4839>, 2018.
- 560 Zhou, S., Lin, J., Qin, X., Chen, Y., and Deng, C.: Determination of atmospheric alkylamines by ion chromatography using 18-crown-6 as mobile phase additive, *J. Chromatogr. A*, 1563, 154–161, <https://doi.org/10.1016/j.chroma.2018.05.074>, 2018.