

Particle deliquescence in a turbulent humidity field

Dennis Niedermeier^{1,*}, Rasmus Hoffmann^{1,*}, Silvio Schmalfuss¹, Wiebke Frey^{1,2}, Fabian Senf¹, Olaf Hellmuth¹, Mira Pöhlker¹, and Frank Stratmann¹

¹Leibniz Institute for Tropospheric Research, Leipzig, Germany

²Wageningen University and Research, Wageningen, Netherlands

*These authors contributed equally to this work.

Correspondence: Dennis Niedermeier (dennis.niedermeier@tropos.de)

Abstract. The atmosphere contains aerosol particles, some of which are hygroscopic in nature. These particles have direct and indirect effects on weather and climate. Furthermore, turbulence causes fluctuations in temperature, water vapor content, and relative humidity (RH). Turbulent humidity fluctuations may influence, among others, the phase state of specific hygroscopic particles. One process of particular interest in that context is particle deliquescence which is the phase transition of solid particles to solution droplets. It occurs at a certain RH, the so-called deliquescence relative humidity (DRH), which in turn depends on e.g., the particle substance. This study investigates the deliquescence behavior of sodium chloride particles in a turbulent humidity field, in particular addressing the questions whether and how turbulent relative humidity fluctuations affect the number / number fraction of deliquesced particles. The turbulent moist air wind tunnel LACIS-T (Turbulent Leipzig Aerosol Cloud Interaction Simulator) is used for this study. The results show that the number of deliquesced particles is influenced by turbulent RH fluctuations. On the one hand, particle deliquescence can be observed although the mean RH is smaller than DRH. On the other hand, there are cases for which non-deliquesced particles are present even though the mean RH is larger than DRH. In general, the number fraction of deliquesced particles depends on a combination of mean relative humidity, strength of humidity fluctuations, and residence time of the particles in the turbulent humidity field. The study concludes that relying solely on the mean relative humidity is inadequate for determining the phase state of deliquescent particle species in the atmosphere. It is necessary to additionally consider both the humidity fluctuations and the particle history.

1 Introduction

A large portion of the atmospheric aerosol consists of hygroscopic particles, which feature different sizes and phase states, and may have immense impacts on Earth's weather and climate. The particle size affects, for example, particle radiative properties as well as the particles' potential to function as cloud condensation nuclei (CCN). The phase state also influences particle radiative properties, as hygroscopically grown particles have different angular scattering properties and refractive indices compared to their dry counterparts (Titos et al., 2016). The liquid particle fraction scatters more light than the respective solid fraction (e.g., Toon et al., 1976; Sloane, 1984). Furthermore, the phase state influences gas-particle partitioning, heterogeneous and multi-phase chemistry processes as, for example, the presence of water allows for reactions with atmospheric pollutants (e.g., Finlayson-Pitts and Hemminger, 2000; Bahadur and Russell, 2008; Liu et al., 2019).

25 The phase state depends on the particle properties, i.e., chemical composition and size, as well as the atmospheric conditions, such as temperature and relative humidity (RH). One process of particular interest in that context is particle deliquescence, for which a solid, water-soluble particle turns into an aqueous solution droplet, thereby increasing its size significantly (Khvorostyanov and Curry, 2014, Ch. 11, pp. 547-575). This phase transition occurs at a certain RH, called deliquescence RH (DRH), and is specific for each deliquescent particle substance. The DRH also depends on temperature (Seinfeld and Pandis, 30 2006, Ch. 10.2.2, pp. 453-454; Khvorostyanov and Curry, 2014, Ch. 11.4.3, pp. 562-563) and particle size (Bahadur and Russell, 2008). Beyond the DRH, a further increase of the RH leads to the growth of the formed solution droplet. However, if the RH decreases below DRH, the material dissolved in the solution droplet will not re-crystallize. The solution will become supersaturated and will remain in a metastable state until it reaches another specific RH, called the efflorescence RH (ERH, with $ERH < DRH$) at which re-crystallization, i.e., efflorescence occurs. This behavior leads to a hysteresis curve which implies that the phase state of a soluble, deliquescent particle, which is at a RH between ERH and DRH, depends on its history (Seinfeld and Pandis, 2006, Ch. 10.2, pp. 449-461; Khvorostyanov and Curry, 2014, Ch. 11, pp. 547-575; Titos et al., 2016).

A wide range of theoretical and experimental studies have been performed in the past and the obtained results significantly increased both the fundamental and quantitative understanding of aerosol particle deliquescence (e.g., Seinfeld and Pandis, 2006; Shchekin et al., 2008, 2013; Hellmuth et al., 2013; Khvorostyanov and Curry, 2014; Hellmuth and Shchekin, 2015; Tang et al., 2019; Peng et al., 2022). This holds even for particles with complex chemical composition. Most of the experimental investigations have focused on the process itself using various techniques (Tang et al., 2019). A majority of experiments were carried out under no-flow conditions. Continuous flow type experiments (such as in Wex et al. (2007)) were carried out under laminar flow conditions. However, the atmosphere is turbulent and turbulent mixing leads to strong fluctuations in temperature, water vapor concentrations, and consequently RH (Siebert et al., 2006; Bodenschatz et al., 2010), which could affect the phase 45 state of deliquescent particles. To our knowledge, the behavior of deliquescent particles in turbulent humidity fields has not yet been investigated in detail. The questions are whether and how turbulent RH fluctuations affect the number of deliquesced particles, and whether or not the number of deliquesced particles is time dependent due to the hysteresis effect.

With the turbulent moist-air wind tunnel LACIS-T (Turbulent Leipzig Aerosol Cloud Interaction Simulator, Niedermeier et al. (2020)), we wanted to address these fundamental questions. We performed various experiments for different mean RHs, 50 RH fluctuation intensities as well as residence times, and determined the number fraction of deliquesced particles. We used size-selected, monodisperse NaCl particles in this study. NaCl was chosen as it is an atmospherically relevant substance being, e.g., the main component of sea salt particles (Niedermeier et al., 2008). Furthermore, the deliquescence properties of NaCl are well characterized: The DRH at 15°C, the temperature used in our experiments, is about 75.5% (Seinfeld and Pandis, 2006, Ch. 10.2.2, pp. 453-454), the temperature dependence of the DRH is weak for NaCl (Seinfeld and Pandis, 2006, Ch. 10.2.2, pp. 453-454; Khvorostyanov and Curry, 2014, Ch. 11.4.3, pp. 562-563), and for a given temperature, DRH is constant for particle 55 diameters larger than 100 nm (Bahadur and Russell, 2008).

The experimental investigations at LACIS-T were accompanied by fluid dynamics simulations performed with OpenFOAM. These large-eddy simulations (LES) assist in setting up the experiments and aid in interpreting the observations. For example, the simulations provide the strength of RH fluctuations that cannot be obtained experimentally.

60 2 Experimental setup

This section describes the generation, size selection as well as pre-conditioning of the NaCl particles, and introduces the basic functionality of LACIS-T, the investigated parameter space, as well as the applied instrumentation.

2.1 Particle generation, size selection and pre-conditioning

The NaCl particles were generated by means of an atomizer (TSI 3075, TSI Inc., St. Paul, Minnesota, USA) atomizing an aqueous solution of 1 g NaCl per liter of double de-ionized water. The resulting aerosol was dried in a diffusion dryer to a RH lower than 20%, i.e., a RH well below the ERH of NaCl particles, which is between 43% and 45% for the investigated particle size range (Tang et al., 1977; Cziczo et al., 1997; Gao et al., 2007).

Downstream of the dryer, the particles are charged by means of a neutralizer. A Differential Mobility Analyzer (DMA, Knutson and Whitby (1975), type “Vienna medium”) is used to select a narrow particle size fraction. Inside the DMA, a further drying occurred as the RH of the DMA sheath air was always $\leq 5\%$. For the experiments, we selected a mobility diameter of $Dp_{\text{mob}} = 400$ nm in order to be able to detect dry solid particles optically by means of a Promo 2000 with welas 2300 aerosol spectrometer (Palas GmbH, Karlsruhe, Germany) inside LACIS-T as well as to minimize the amount of larger, doubly charged particles being present. As it turns out, their proportion to the total number of selected particles is less than 1%, i.e., doubly charged particles do not affect our results. Note that a particle shape factor has to be considered when converting the mobility diameter to a mass equivalent diameter, because of the non-spherical shape of solid NaCl particles. This factor is 1.08 for NaCl particles according to Kelly and McMurry (e.g., 1992). Earlier measurements with the laminar flow-tube LACIS (Wex et al., 2005; Niedermeier et al., 2008) confirm that this shape factor is valid for NaCl particles in the selected size range. In consequence, $Dp_{\text{mob}} = 400$ nm corresponds to a mass equivalent diameter of $Dp_{\text{me}} = 370$ nm.

The number concentration of the selected particles was determined utilizing a Condensational Particle Counter (CPC, TSI 3010, TSI Inc., St. Paul, Minnesota, USA), and was kept at about 1000 cm^{-3} by means of a dilution system upstream of the DMA. The dilution system consists of a by-pass, a filter and two valves for adjustment of the flows. All flows, i.e., aerosol flow, DMA sheath, and excess air-flow as well as CPC sample flow were controlled by mass flow controllers (Brooks Instrument GmbH, Dresden, Germany) and checked with a bubble flow meter (Gilian® GilibratorTM2, Sensidyne Inc., Clearwater, Florida, USA) on a daily basis.

For the experiments, three different types of pre-conditioning were applied concerning the NaCl particles which were fed into LACIS-T:

Case (i): The particles were left dry (i.e., $\text{RH} \leq 5\%$) so that solid NaCl particles entered the measurement section of LACIS-T. This setup was only used to determine the optical size distribution of solid NaCl particles (see Sect. 4.2), as dry particle injection into LACIS-T reduces the mean RH in the mixing zone of the measurement section significantly (see Appendix A).

Case (ii): The particles were pre-humidified to a dew-point temperature of $T_d = 9.6^\circ\text{C} (\pm 0.1^\circ\text{C})$ by mixing the dry aerosol with pre-humidified particle-free air by means of a saturator (Perma Pure MH-110-12S-4, Perma Pure LLC,




Figure 1. Schematic of LACIS-T including main components and possible conditions. Schematic copyrights: Ingenieurbüro Mathias Lippold, VDI; TROPOS. The figure is adapted from Niedermeier et al. (2020) and modified.

Toms River, New Jersey, USA). This led to a RH of 70.2% ($\pm 0.3\%$) at 15°C, i.e., the NaCl particles were still non-deliquesced before entering LACIS-T. With RH at the LACIS-T inlet being closer to DRH, the mean RH inside the measurement section of LACIS-T is not reduced significantly (see LES results in Sect. 4.1). This setup was used for the LACIS-T experiments for different sets of mean RH as well as strength of RH fluctuations.

Case (iii): The particles were pre-humidified at RH = 100% ($T_d = 12.5^\circ\text{C} (\pm 0.1^\circ\text{C})$) by means of a saturator (Perma Pure MH-110-12S-4, Perma Pure LLC, Toms River, New Jersey, USA) so that the NaCl particles deliquesce before entering LACIS-T (RH = 85.0% at 15°C). This setup was used to determine the optical size distribution of fully deliquesced NaCl particles inside LACIS-T for the different sets of conditions.

2.2 LACIS-T

LACIS-T is a unique turbulent moist-air wind tunnel which has been established to study cloud physical processes, in general, and cloud microphysics – turbulence interactions, in particular. LACIS-T can be operated under a wide range of well-characterized and reproducible initial and boundary conditions resembling atmospheric warm, mixed-phase and cold clouds. The design, functionality and capabilities of the setup are described in detail in Niedermeier et al. (2020) and only a brief description will be given here.

LACIS-T is a closed-loop wind tunnel of Göttingen type (Fig. 1). The unique characteristic of LACIS-T is that it features two parallel flow branches, in which two particle-free air-flows 'A' and 'B' can be separately conditioned and controlled with respect to volume flow rate (through the radial blowers and valves, up to 6000 l/min each), water vapor content in terms

110 of a dew-point temperature (through the humidification system, dew-point temperature range $-40^{\circ}\text{C} < T_d < 25^{\circ}\text{C}$, with an accuracy of 0.1 K), and temperature (through the heat exchangers, temperature range $-40^{\circ}\text{C} < T < 25^{\circ}\text{C}$, with an accuracy of 0.05 K). These two air-flows pass passive square-mesh grids through which defined turbulence is induced. They then enter the measurement section, at the entrance of which the particle-free air-flows are combined and turbulently mixed. The measurement section is of cuboid shape with the dimensions of 200 cm x 80 cm x 20 cm. The aerosol particles are introduced into the mixing
 115 zone of the two turbulent particle-free air-flows. This mixing zone provides an ideal environment for studying the influence of the turbulent fluctuations on aerosol and cloud microphysical processes. Downstream of the measurement section, the flow is directed towards an adsorption dehumidifying system where it is dried and heated. Afterwards, the flow is split up into the two branches, cleaned by particle filters, and the whole cycle starts again.

In this study, the flow rate was set to 4300 l/min for each particle-free air-flow leading to a mean flow velocity of 1.35 m/s
 120 in the measurement section. Turbulence conditions inside the measurement section are determined by means of Hot Wire anemometry (Dantec Dynamics Inc., Skovlunde, Denmark) and LES and are very similar to those described in Niedermeier et al. (2020). For example, the eddy turnover time τ_{mix} which is a measure for the turbulent mixing time scale, is between $\tau_{\text{mix}} = 0.1 - 0.7$ s. It increases due to the decaying turbulence inside the measurement section and is given as $\tau_{\text{mix}} = (l_T^2/\varepsilon)^{(1/3)}$ (e.g., Baker et al., 1984; Lehmann et al., 2009). The quantity ε represents the energy dissipation rate, while l_T denotes the Taylor
 125 microscale. In the context of atmospheric conditions, the Taylor microscale is frequently considered to be the appropriate mixing length scale (Lehmann et al., 2009).

We use isothermal conditions in this study, i.e., the temperature in both particle-free air-flows was identical, set to $T_A = T_B = 15^{\circ}\text{C}$ and monitored via PT100 temperature sensors (1/10 class B, DIN EN 60751, accuracy of $\pm(0.0300^{\circ}\text{C} + 0.0005 \times T)$). The dew-point temperature was set individually in each particle-free airflow and monitored by means of two dew-point mirrors
 130 (DPM, model 973 by MBW Calibration AG, accuracy of $\leq \pm 0.1$ K; reproducibility of $\leq \pm 0.05$ K), sampling air in each flow branch between heat exchanger and turbulence grid. Due to the individual settings of the dew-point temperatures, and thus the individual RH values in each flow branch, various RH mixing conditions could be established. That means different mean RH values, and different strengths of RH fluctuations, could be set in the mixing region (see Fig. 2a as an example for a specific set of RH conditions in branch A and B). In general, the larger the difference between RH_A and RH_B , the larger the strength of
 135 the RH fluctuations. However, no direct measurement of dew point temperature fluctuations and consequently RH fluctuations is available so far. The strengths of the fluctuations are determined by means of the LES which is described in Sect. 4.1.

Finally, particle size distributions are determined optically via a Promo 2000 with welas 2300 aerosol spectrometer (Palas GmbH, Karlsruhe, Germany). The welas 2300 sensor is placed inside the measurement section having a 15 cm long stainless-steel tube (5 mm inner diameter) on top of its own inlet tube. This additional long tube ensures particle extraction and detection
 140 from the flow field, which is only very weakly influenced by the body of welas 2300, which disturbs the flow field in its closer vicinity. Please note that the distance of the tip of this stainless-steel tube relative to the aerosol inlet of LACIS-T, called ‘ z ’, is used as a reference in the later data description.

In the first set of experiments, welas 2300 is placed at a fixed position inside the measurement section leading to a fixed residence time of the particles until the detection occurs. The inlet tube is placed at $z = 30$ cm below the aerosol inlet of

145 LACIS-T. In the second set of experiments, the position of the tube together with the welas 2300 is varied leading to different mean residence times of the particles.

3 Numerical setup

Large-eddy simulations directly compute the large-scale motions of the flow fields from the Navier-Stokes equations, and parameterize the unresolved small-scale/sub-grid scale (SGS) motions. We use the dynamic k-equation LES model in Open-
150 FOAM based on Chai and Mahesh (2012), who developed a new transport equation for the subgrid-scale kinetic energy, which has proven to be a good model for decaying turbulence and the transport of thermodynamic quantities. A hexaeder dominated mesh is applied, refined near turbulence grid and walls, with $7.6 \cdot 10^6$ cells in total. We use an Euler-Lagrange approach for the simulations, i.e., temperature and water vapor mixing ratio are calculated at the grid points while each individual particle is tracked along its trajectory through the thermodynamic field. Details about the general numerical setup, especially about how
155 to set the initial and boundary conditions, are given in Niedermeier et al. (2020).

Overall, the main goals of these simulations were a) the determination of the RHs the particles experience along their way through the measurement section for the different experimental conditions, and b) the determination of the deliquesced particle fraction for comparison with the gained experimental data. The RHs, and RH fluctuations the particles experience, which cannot be detected experimentally, were quantified in terms of mean standard deviation of simulation-derived RH probability
160 density functions (pdfs). Thereto, the RH pdfs are calculated for specific locations along the mixing region inside LACIS-T's measurement section. Furthermore, the numbers of deliquesced and non-deliquesced particles were derived from the simulations and compared with the experimentally obtained ones. Concerning the latter, a dynamical growth law is implemented for the Lagrangian particle simulations accounting for diffusive vapor transport and deposition to the particle surface, and latent heat release (Wilck, 1999) as well as deliquescence and efflorescence. However, the feedback effects on heat transport and
165 mass transfer in the gas phase are ignored as the droplet concentrations (mass and volume ratio) considered in this study are too low to influence the continuous phase. As we are modeling the particle hygroscopic growth dynamically, the model requires information concerning the deliquescence time scale for the considered NaCl particles. We assumed deliquescence to occur on the time scale τ_{del} of 10^{-4} s (the numerical time step is also 10^{-4} s) once $\text{DRH} = 75.5\%$ is reached (Note that this time scale is 3 orders of magnitude smaller than the mixing time scale τ_{mix} which is between 0.2 and 0.7 s). This assumption is based on
170 molecular dynamics simulations performed by Bahadur and Russell (2008) who give the time scale τ_{del} required for complete deliquescence of an infinitely extended "planar" NaCl slab to be $\tau_{\text{del}} = 9.96 \cdot 10^{-5}$ s. The extent to which this assumption is reasonable will be the subject of discussion in Appendix B.

For particle efflorescence, we also assume that the re-crystallization occurs on the time scale of 10^{-4} s once $\text{ERH} = 45.0\%$ is reached. Tang and Munkelwitz (1984) show that the re-crystallization of a single micrometer-sized NaCl particle can occur
175 very quickly, however, a clear time scale cannot be determined from Fig. 3 in their paper. Other studies like Ma et al. (2019) show that the time scale for particle efflorescence, which is a nucleation process, depends on the ambient RH and is inversely proportional to the nucleation rate. Various studies show that the ERH of NaCl particles is between 43% and 45% for the

investigated particle size range (Tang et al., 1977; Cziczo et al., 1997; Gao et al., 2007). As it turned out, the minimum RH reached in our experiments is well above 45%. From this point of view, our assumption about the time scale of particle
180 efflorescence will supposedly not affect the simulation results as efflorescence is unlikely to occur in our investigations.

4 Results

This section presents the results of the numerical and experimental investigations. First of all, the LES provide the RH experienced by the particles. This is then applied to determine the strength of the RH fluctuations which are later used for the interpretation of the experimental results. The experiments themselves are performed for different mean RH, strengths of RH
185 fluctuations, as well as for fixed and variable particle residence times. Number fractions of deliquesced particles are determined and discussed.

4.1 Turbulent RH field obtained via LES

At first, the simulations are employed to determine the RH experienced by the particles, as well as to ascertain the impact of the particles themselves on the RH field within the mixing region inside the measurement section. This is presented exemplarily for
190 the setting of $RH_A = 60\%$ and $RH_B = 85\%$ leading to a mean RH of 72.5% along the center line. About 78 000 monodisperse NaCl particles with a mass equivalent diameter of $Dp_{me} = 370$ nm are tracked inside the measurement section. The NaCl particles are pre-conditioned according to case (ii), i.e., the aerosol is injected with $RH = 70.2\%$ (as introduced in sect. 2.1). In Fig. 2a, a snapshot of the instantaneous RH field in the symmetry plane is shown including the respective particles deliquescing and growing along the inverse vertical axis. The dashed line represents the center line, where the strongest mixing occurs, as
195 well as where the particles are detected further downstream. As the particles shown move through the channel, statistics are done at distinct horizontal planes over time, i.e., data from particles that pass a given horizontal plane is stored and later on analyzed.

As mentioned, the RH of the introduced aerosol is slightly lower than the mean RH along the center line of the measurement section. This influences the RH field along the center line, which is illustrated in Fig. 2b in terms of median RH (RH_{med}),
200 mean RH (RH_{mean}) and standard deviation of the RH (σ_{RH}). First of all, it can be seen that generally RH_{med} and RH_{mean} fall together. Secondly, RH_{mean} is observed to be slightly lower close to the location of aerosol injection. However, it increases on the very first centimeters and then reaches a constant value of 72.5% at a height of approximately $z = 15$ cm. Additionally, σ_{RH} demonstrates an increase within the initial 15 cm, subsequently reaching a constant value of 4.9%. Consequently, the data obtained for $z \geq 15$ cm will be employed for the subsequent data interpretation. As an example, the RH experienced by
205 particles near the center line when passing the horizontal plane at $z = 30$ cm is shown as normalized frequencies in Fig. 2c and the obtained RH distribution has a Gaussian shape.

Note that there is an influence of the SGS motions onto the simulated fluctuations and consequently σ_{RH} . Based on currently ongoing work and in accordance to a study performed by Chandrakar et al. (2022), we estimate the contributions from these




Figure 2. a) A snapshot of the particle simulation with the fluctuating RH field in the background (flow direction from top to bottom). It depicts the scenario of $RH_A = 60\%$ (right side) and $RH_B = 85\%$ (left side) resulting in a mean RH of 72.5% along the center line (dashed white line). Note that only 10% of the particles are depicted. The particles are colored and sized (not to scale) according to their diameter. b) The obtained temporal median RH, mean RH, and RH fluctuations in terms of $\pm \sigma_{RH}$ plotted along the center line as a function of the distance z to the aerosol inlet of the measurement section. Note that the pre-humidified aerosol is injected (case(ii): $RH = 70.2\%$ at 15°C). c) The normalized frequency of the RH experienced by the particles. The distribution has a Gaussian shape as evidenced by the normal distribution with $RH_{\text{mean}} = 72.5\%$ and $\sigma_{RH} = 4.9\%$.

SGS fluctuations to the total fluctuations to be max. $\pm 0.04 \times \sigma_{RH}$, which represents the uncertainty of the determined σ_{RH} values.

4.2 Obtained particle size distributions and determined deliquesced particle fraction

In order to be able to determine the fraction of deliquesced particles, first the size distributions of both, dry and deliquesced particles, as well as those of a mixture of both need to be considered. Therefore, in a first set of experiments, different RH conditions are adjusted in order to obtain and interpret the corresponding size distributions. The corresponding RH_{mean} values at the center line are calculated from the RH values of the individual particle-free airflows. In general, it turns out from repeated measurements of temperature and dew-point temperature inside the tunnel that there is an absolute uncertainty in RH_{mean} of up to 0.6%. The width of the RH distribution which is given in terms of σ_{RH} is obtained from the LES. For this first set of experiments, the welas 2300 position was fixed with its inlet being placed at $z = 30$ cm leading to a mean residence time of

Table 1. NaCl particle and LACIS-T conditions for four different experiments. Column 1 gives the Experiment number. Column 2 shows the condition of the NaCl particles before being inserted into LACIS-T. Columns 3 and 4 show the corresponding RH conditions inside LACIS-T in terms of RH_{mean} and σ_{RH} .

Experiment #	NaCl particle conditions*	LACIS-T conditions	
		RH_{mean} [%]	σ_{RH} [%]
(1)	case (i) - solid, dry	20.0	0
(2)	case (ii) - solid, pre-humidified	72.5	0
(3)	case (iii) - fully deliquesced	72.5	4.9
(4)	case (ii) - solid, pre-humidified	72.5	4.9

* see Sect. 2.1 for details

the particles before detection of approx. 0.22 s as the mean velocity is 1.35 m/s. Figure 3 shows the results of four different experiments for the conditions given in Table 1.




Figure 3. Normalized particle number size distributions determined for the four different experiments being summarized in Table 1. Experiments (1), (2), and (3) are given by the solid red, dotted red, and solid blue lines, respectively. Note that experiment (4) leads to a bimodal distribution @ $RH_{\text{mean}} = 72.5\%$, $\sigma_{RH} = 4.9\%$ featuring both solid and deliquesced NaCl particles (brown solid curve). The dashed brown curve represents a bimodal fit to curve of experiment (4) in order to determine the solid and deliquesced particle fractions.

No distinct difference in the obtained particle size distributions is observed for experiment (1) and (2). During experiment (2), the particles most likely start to take up water molecules leading to a thin liquid shell surrounding the particles, however, particles are still non-deliquesced under these humid conditions. This is in agreement to observations obtained by Krueger et al. (2003) and Wise et al. (2008) for single NaCl particles.

225 Experiment (3) shows a clear increase in particle size compared to (1) and (2). In this case, the NaCl particles are fully deliquesced. The mean particle diameter determined optically agrees with the diameter calculated with the Köhler equation (Niedermeier et al., 2008) for this RH_{mean} .

Finally, experiment (4) results in two modes. The first mode fits with the one for solid NaCl particles, while the second mode fits with the one obtained for the deliquesced particles. That means, we observe both solid and deliquesced particles at
230 the same time. In other words, the humidity fluctuations lead to particle deliquescence although RH_{mean} is smaller than DRH which is about 75.5% at 15°C (Seinfeld and Pandis, 2006, Ch. 10.2.2, pp. 453–454). By fitting both modes with two log-normal distributions (see Fig. 3) the solid and deliquesced particles fractions can be determined which in the example are 0.57 and 0.43, respectively.

To systematically quantify this effect, such experiments have been performed for various thermodynamic and flow condi-
235 tions, as well as particle residence time (i.e., t_{res} 2300 sensor positions).

4.3 Dependence of deliquesced particle fraction on RH field and residence time

First of all, the influence of RH fluctuations on the deliquesced particle fraction was determined for a fixed t_{res} 2300 position, i.e. a fixed mean residence time. Thereto, the t_{res} 2300 inlet was placed again at $z = 30$ cm. RH_{mean} was varied between 71.6% and 78.1%, i.e., from below to above DRH of the investigated NaCl particles, and σ_{RH} was set between 0% and 9.5%.
240 A few studies exist (e.g., MacPherson et al., 1992; Lenchow et al., 1994; Kulmala et al., 1997; Siebert and Shaw, 2017), which show σ_{RH} values in the range of 1% to 4.6%, depending on the environmental conditions. It means that our measurements cover this range of observed strengths of RH fluctuations.

The obtained deliquesced particles fractions f_{del} are depicted in Fig. 4 as a function of σ_{RH} for four different RH_{mean} values.

Generally, it becomes obvious that the humidity fluctuations have a strong influence on the fraction of deliquesced particles.
245 Particle deliquescence can be observed although $RH_{\text{mean}} < \text{DRH}$. Furthermore, we detect an increase of the deliquesced particle fraction with increasing σ_{RH} as long as $RH_{\text{mean}} < \text{DRH}$. In this case, an increasing σ_{RH} increases the probability that solid NaCl particles are in a RH field with RH-values higher than DRH. The slope of f_{del} flattens for RH_{mean} being close to DRH and becomes negative for $RH_{\text{mean}} > \text{DRH}$ as now the increasing σ_{RH} increases the probability that solid NaCl particles experience a RH field with RH-values lower than DRH, and therefore do not deliquesce. In other words, in this case
250 not all particles deliquesce although $RH_{\text{mean}} > \text{DRH}$.

Secondly, the influence of RH fluctuations on the deliquesced particle fraction was determined as a function of mean particle residence time t_{res} in a fluctuating RH field. This was achieved by changing the position of the t_{res} 2300. The purpose of this type of experiment is as follows. Particles once deliquesced will not recrystallize / effloresce at DRH. They would do so at the ERH, which is between 43% and 45% for NaCl particles of the investigated size range (Tang et al., 1977; Cziczo et al.,
255 1997; Gao et al., 2007). However, in our experiments, even for the broadest RH distributions we do not reach this ERH. That means if solid NaCl particles are inserted in fluctuating RH fields, the residence time in this field might influence the number and fraction of deliquesced particles.




Figure 4. Deliquesced particle fraction f_{del} as a function of the standard deviation of the RH distribution σ_{RH} . The width of the symbols represents the range of uncertainty of σ_{RH} based on the LES results. The shown error bars originate from repeated measurements (three to five times) giving the minimal and maximal obtained f_{del} , respectively.

Figure 5 shows a series of normalized particle size distributions for $RH_{mean} = 75.9\%$ and $\sigma_{RH} = 4.9\%$, with different t_{res} inside the measurement section. It can be seen that the deliquesced particle mode of the distributions (right mode) increases with increasing t_{res} while the solid particle mode (left mode) decreases. In other words, more particles deliquesce with increasing t_{res} due to the hysteresis effect.

Measurements as presented in Fig. 5 have been carried out for other sets of RH_{mean} and σ_{RH} values. The respective results are summarized in Fig. 6. Here, the fraction of deliquesced particles f_{del} is depicted as a function of t_{res} (logarithmic scale) inside the fluctuating RH environment. In the left plot, RH_{mean} is fixed and σ_{RH} is varied. In the right plot, it is vice versa. The key findings are that the fraction of deliquesced particles increases with increasing t_{res} for all investigated cases. For fixed RH_{mean} , the slope of the three curves looks very similar pointing towards an approximately exponential relationship between f_{del} and t_{res} , with the curves shifting upwards to higher values of f_{del} as σ_{RH} increases. When looking at the curves for fixed σ_{RH} it can be observed that the slopes become flatter with increasing RH_{mean} , as a greater proportion of the particles is already deliquesced at the lowest measured residence times. Consequently, fewer solid particles are available for deliquescence.

In summary, it can be said that the combination of turbulent RH fluctuations and hysteresis in aerosol particle deliquescence and efflorescence, has a significant impact on the fraction of deliquesced particles over time. The time required for all NaCl




Figure 5. Normalized particle number size distributions determined experimentally for $\text{RH}_{\text{mean}} = 75.9\%$ and $\sigma_{\text{RH}} = 4.9\%$, but for different residence times t_{res} inside the measurement section. The colors of the different number distributions correspond to the colormap on the right-hand side of the plot denoting t_{res} .

particles to deliquesce depends on both the mean RH and the strength of the fluctuations, or in other words, the proportion of deliquesced particles is dependent on RH_{mean} , σ_{RH} , and residence time.

Finally, we simulated the deliquescence of the NaCl particles in the turbulent RH field by running LES in OpenFOAM. In general, the time dependence of f_{del} can also be identified in the model results. However, the simulations appear to overestimate the observed deliquesced particle fractions, which might be caused by our assumption of a constant time scale for the deliquescence process itself. A detailed discussion about this discrepancy and its possible reasoning is given in Appendix B. Nevertheless it is important to point out that this discrepancy between measurement and simulation does not invalidate the experimental observations and central statements. Furthermore, it motivates us to investigate the suitability of different theories in the future (more details given in Appendix B).

5 Summary and conclusion

We investigated the deliquescence behavior of size-selected, monodisperse NaCl particles in a turbulent humidity field with LACIS-T. The mean RH, the strength of RH fluctuations and the residence time of the particles in the turbulent humidity field were varied. In general, we found that turbulence affects the number of deliquesced particles in a particle population and this number depends on the combination of all three of the aforementioned variables. Fluctuations in RH can lead to particle deliquescence, despite the mean RH being below the deliquescence RH. Conversely, particle deliquescence can be hindered even though the mean RH is above the deliquescence RH. However, a population of solid, non-deliquesced NaCl particles




Figure 6. Deliquesced particle fraction f_{del} as a function of particle residence time t_{res} . Left figure: RH_{mean} was set to 72.5% and σ_{RH} was varied. Right figure: σ_{RH} was set to 4.9% and RH_{mean} was varied.

introduced into a fluctuating RH field, where the RH is always greater than the efflorescence RH, and the RH fluctuations exceed DRH will deliquesce completely due the combination of the turbulent RH fluctuations and the hysteresis effect. The time required to reach this fully deliquesced state is contingent upon the mean RH and the strength of the fluctuations.

We are able to simulate the general behavior of the NaCl particle deliquescence in the turbulent RH field by running LES in OpenFOAM. However, the simulations tend to overestimate the observed deliquesced particle fractions, which might be caused by our assumption of a constant time scale for the deliquescence process itself. In order to achieve a more accurate representation of the experimental observations, different theories for the derivation of the deliquescence time scale τ_{del} will be tested in the future. Following, for example, the argumentation of Khvorostyanov and Curry (2014, Ch. 11.3, pp. 553-558), nucleation rates, and with that characteristic deliquescence time scales - which depend on RH among other things - will be determined and implemented into the LES model.

Ultimately, the observation of the onset of particle deliquescence below, and the presence of non-deliquesced particles above the DRH in a turbulent humid field is inline with the argumentation of Prigogine (1979), according to which a consistent macroscopic description is no longer given in the vicinity of non-equilibrium phase transitions (such as the deliquescence transition). Near the phase transition (here the DRH), the turbulent fluctuations become as important as the mean values. Macroscopic values represent the most likely ones, which are identical with the mean values only if fluctuations can be neglected. In the real atmosphere, however, turbulent fluctuations are always present. A few studies exist (e.g., MacPherson et al., 1992; Lenchow

et al., 1994; Kulmala et al., 1997; Siebert and Shaw, 2017), which show σ_{RH} values in the range of 1% to 4.6%, depending
305 on the environmental conditions. Our investigations cover this range of observed RH fluctuations and we observe a distinct
influence of the turbulent RH fluctuation on the fraction of deliquesced particles at these σ_{RH} values. This indicates that our
results are relevant for the atmosphere.

Altogether, this implies that the description of hygroscopic growth and shrinking during humidification and dehumidification
of the ambient atmosphere, respectively, requires the consideration of both (i) the hysteresis effects during particle evolution
310 and in addition (ii) the turbulent character of the thermodynamic conditions of the ambient atmosphere, which affect the
macroscopic boundary conditions of phase transition. This is of particular importance, among others, for the purposes of
atmospheric modeling as the optical properties of solid and deliquesced particles differ, which needs to be carefully considered,
for example, in radiative transfer schemes in global atmospheric models (Haarig et al., 2017). In that sense, we also suggest to
collect additional data on atmospheric RH fluctuations, to be carried out in terrestrial and marine environments, at ground level
315 and above.

Data availability. Data sets will be made available via ACTRIS data centre in case of manuscript acceptance.

Appendix A: Comparison of two different NaCl pre-conditioning settings and its effect on the RH field inside LACIS-T

Here, we compare the influence of two different types of NaCl particle pre-conditioning, i.e., case (i) and case (ii) as introduced
320 in sect. 2.1, onto the RH field along the center line within the measurement section where the particles are injected, transported
and later detected. The comparison is based on the LES performed in OpenFOAM. In both cases, the settings are $RH_A = 60\%$
and $RH_B = 85\%$ leading to a mean RH of 72.5% along the center line and about 78 000 monodisperse NaCl particles with a
mass equivalent diameter of $Dp_{me} = 370$ nm are tracked inside the measurement section. The RH that the particles experience
is shown in Fig. A1 in terms of median RH (RH_{med}), mean RH (RH_{mean}) and standard deviation of the RH (σ_{RH}). For both
325 cases, RH_{med} and RH_{mean} fall together. The RH fluctuation distributions (not shown here) have a Gaussian shape. For case
(i), the aerosol is injected with $RH \leq 5\%$ and it can be observed that RH_{mean} in the mixing zone is lowered significantly close
to the aerosol inlet due to the low aerosol RH. The obtained RH_{mean} is about 46% which is 26.5% lower as it would occur in
case of no particle injection. With increasing distance to the aerosol inlet, RH_{mean} rises until about $z = 30$ cm distance where
it starts to become constant. Due to this increase of RH_{mean} , the whole RH distribution gets shifted towards larger RH values
330 which additionally influences σ_{RH} . This steep increase of RH_{mean} combined with the increase of σ_{RH} would complicate the
data interpretation. Therefore, this setup was not used for our investigations.

For case (ii), the aerosol is injected with $RH = 70.2\%$ which results in an only slightly lowered RH_{mean} in the mixing
zone. σ_{RH} increases within the first 15 cm, becoming constant to a value of 4.9%. Therefore, this setup was applied for our
experiments and data obtained for $z \geq 15$ cm was used for the later data interpretation.




Figure A1. LES results for the scenario of $RH_A = 60\%$ and $RH_B = 85\%$ which results in a mean RH of 72.5% along the center region (i.e., the main mixing zone). The obtained median RH, mean RH and RH fluctuations in terms of $\pm\sigma_{RH}$ are plotted for the center region as a function of the distance z to the aerosol inlet of the measurement section for two different types of particle injection. Left figure: Injection of dry aerosol (case (i): $RH \leq 5\%$ at 15°C). Right figure: Injection of pre-humidified aerosol (case(ii): $RH = 70.2\%$ at 15°C).

335 Appendix B: Comparison between measurements and simulations of f_{del} as a function of t_{res}

The model study was performed to test whether or not we are sensitive enough to observe the effects of turbulent RH fluctuations on the fraction of deliquesced particles. In Figure B1, the measured and simulated deliquesced particle fractions f_{del} are plotted as a function of particle residence time t_{res} exemplarily for $RH_{\text{mean}} = 72.5\%$ and three different σ_{RH} values. In general, the time dependence of f_{del} can also be identified in the model results. However, the simulations tend to overestimate f_{del} for the cases of $\sigma_{RH} = 4.9\%$ and 9.5% , and for residence times below 0.3 s. For the $\sigma_{RH} = 2.8\%$ case, there is a closer agreement between the measurement and the simulation. However, the simulated slope differs from the experimentally determined one. This latter observation also holds for the other two cases.

The question arises about the reason for this discrepancy. In general, the LES model has proven to be an effective tool for simulating the thermodynamic and fluidic conditions inside the measurement section as well as the hygroscopic and dynamic growth of droplets which formed on NaCl particles (Niedermeier et al., 2020). In view of the lack of full information of the microscopic state of the investigated system, the modeling setup required some closure assumptions, which inheres uncertainties. Perhaps our assumption about the time scale of deliquescence (τ_{del}) is too simplified. We assumed - as mentioned before - that NaCl deliquescence occurs on the time scale of 10^{-4} s once $DRH = 75.5\%$ is reached based on molecular dynamics simulations performed by Bahadur and Russell (2008). An increase of the deliquescence time scale τ_{del} to 10^{-3} s does not lead to a significant change of the simulated deliquesced particle fractions (not shown) because the mixing time scale τ_{mix}




Figure B1. Measured (symbols) and simulated (lines) deliquesced particle fraction f_{del} as a function of particle residence time t_{res} for $\text{RH}_{\text{mean}} = 72.5\%$.

(in the order of 10^{-1} s) is still two orders of magnitude larger than τ_{del} so that the microphysical system is able to react on thermodynamic changes.

Several other theories of deliquescence have been developed during the last decades (McGraw and Lewis, 2009; Lamb and Verlinde, 2011, Ch. 7.1, pp. 290-295; Hellmuth et al., 2013, Ch. 12.2, pp. 319-334; Shchekin et al., 2013;0 Khvorostyanov and Curry, 2014, Ch. 11.3, pp. 553-558) which are based on different approaches. For example, Khvorostyanov and Curry (2014, Ch. 11.3, pp. 553-558) describe deliquescence as a nucleation process (being in line with experiments performed by Cantrell et al. (2002)) beginning with the formation of a liquid solution germ on a crystal surface. Consequently, they derive a nucleation rate in analogy to surface melting. From this nucleation rate, τ_{del} for the deliquescence process of NaCl particles could be determined which depends, among others, on the actual RH. However, it should be noted that we were not able to observe / resolve the nucleation process itself in our laboratory experiments.

One could also think about an experimental approach in order to determine τ_{del} . However, for these measurements energy dissipation rates ranging from 10^{-2} up to 10^6 m^2/s^3 would be needed so that the mixing time scale τ_{mix} could be varied, ranging from values larger than τ_{del} to values smaller than it. In the two extremes, the microphysical system will either be able to react on thermodynamic changes (fast microphysics, $\tau_{\text{mix}} \gg \tau_{\text{del}}$) or not (slow microphysics, $\tau_{\text{mix}} \ll \tau_{\text{del}}$). The transition region ($\tau_{\text{mix}} \approx \tau_{\text{del}}$) could give an estimate for the deliquescence time scale.

However, testing different theories and performing additional experiments for the derivation of τ_{del} is beyond the scope of this study. It has to be the objective of future studies to investigate the suitability of different theories and experimental approaches in this context.

Author contributions. DeNi and RaHo wrote the manuscript with contributions from all co-authors. LACIS-T measurements and data evaluation were performed by DeNi and RaHo with contributions from SiSc, WiFr, MiPö and FrSt. Numerical simulations were performed by SiSc with contributions from DeNi, RaHo, FaSe, OIHe, and FrSt. All authors discussed the experimental and numerical results.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We received funding for LACIS-T from ACTRIS-D which the German contribution to the Aerosol, Cloud, Trace Gases Research Infrastructure ACTRIS. ACTRIS-D is funded by the German Federal Ministry for Education and Research (BMBF) under grant agreements 01LK2001A-K & 01LK2002A-G. Wiebke Frey has received funding from the European Union's Horizon 2020 RI programme as an Individual Fellowship under the Marie Skłodowska-Curie grant agreement number 835305.

References

- Bahadur, R. and Russell, L. M.: Water uptake coefficients and deliquescence of NaCl nanoparticles at atmospheric relative humidities from molecular dynamics simulations, *The Journal of Chemical Physics*, 129, 094 508, <https://doi.org/10.1063/1.2971040>, 2008.
- 380 Baker, M. B., Breidenthal, R. E., Choulaton, T. W., and Latham, J.: The effects of turbulent mixing in clouds, *Journal of Atmospheric Sciences*, 41, 299–304, [https://doi.org/10.1175/1520-0469\(1984\)041%3C0299:TEOTMI%3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1984)041%3C0299:TEOTMI%3E2.0.CO;2), 1984.
- Bodenschatz, E., Malinowski, S. P., Shaw, R. A., and Stratmann, F.: Can we understand clouds without turbulence?, *Science*, 327, 970–971, doi:10.1126/science.1185138, 2010.
- Cantrell, W., McCrory, C. and Ewing, G. E., Nucleated deliquescence of salt, *The Journal of Chemical Physics*, 116(5), 2116–2120, <https://doi.org/10.1063/1.1429924>, 2002.
- 385 Chai, X. and Mahesh, K.: Dynamic-equation model for large-eddy simulation of compressible flows, *Journal of Fluid Mechanics*, 699, 385–413, <https://doi.org/10.1017/jfm.2012.115>, 2012.
- Chandrakar, K. K., Morrison, H., Grabowski, W. W., Bryan, G. H., and Shaw, R. A., Supersaturation variability from scalar mixing: Evaluation of a new subgrid-scale model using direct numerical simulations of turbulent Rayleigh–Bénard convection, *Journal of the Atmospheric Sciences*, 79(4), 1191–1210, <https://doi.org/10.1175/JAS-D-21-0250.1>, 2022.
- 390 Cziczo, D. J., Nowak, J. B., Hu, J. H., and Abbatt, J. P. D.: Infrared spectroscopy of model tropospheric aerosols as a function of relative humidity: Observation of deliquescence and crystallization, *Journal of Geophysical Research: Atmospheres*, 102, 18 843–18 850, <https://doi.org/10.1029/97JD01361>, 1997.
- Finlayson-Pitts, B. J. and Hemminger, J. C.: Physical Chemistry of airborne sea salt particles and their components, *The Journal of Physical Chemistry A*, 104, 11 463–11 477, <https://doi.org/10.1021/jp002968n>, 2000.
- 395 Gao, Y., Chen, S. B., and Liya, E. Y.: Efflorescence relative humidity of airborne sodium chloride particles: A theoretical investigation, *Atmospheric Environment*, 41, 2019–2023, <https://doi.org/10.1016/j.atmosenv.2006.12.014>, 2007.
- Haarig, M., Ansmann, A., Gasteiger, J., Kandler, K., Althausen, D., Baars, H., Radenz, M., and Farrell, D. A.: Dry versus wet marine particle optical properties: RH dependence of depolarization ratio, backscatter, and extinction from multiwavelength lidar measurements during SALTRACE, *Atmospheric Chemistry and Physics*, 17, 14 199–14 217, <https://doi.org/10.5194/acp-17-14199-2017>, 2017.
- 400 Hellmuth, O., Khvorostyanov, V. I., Curry, J. A., Shchekin, A. K., Schmelzer, J. W. P., Feistel, R., Djikaev, Y. S., and Baidakov, V. G.: Nucleation Theory and Applications: Special issues. Review series on selected topics of atmospheric sol formation Vol. 1, Joint Institute for Nuclear Research (JINR), Dubna, ISBN: 978-5-9530-0349-0, 2013.
- Hellmuth, O., and Shchekin, A. K.: Determination of interfacial parameters of a soluble particle in a nonideal solution from measured deliquescence and efflorescence humidities, *Atmospheric Chemistry and Physics*, 15, 3851–3871, <https://doi.org/10.5194/acp-15-3851-2015>, 2015.
- 405 Kelly, W. P. and McMurry, P. H.: Measurement of particle density by inertial classification of differential mobility analyzer-generated monodisperse aerosols, *Aerosol Science and Technology*, 17, 199–212, <https://doi.org/10.1080/02786829208959571>, 1992.
- Khvorostyanov, V. I. and Curry, J. A.: Thermodynamics, kinetics, and microphysics of clouds, Cambridge University Press, 2014.
- 410 Knutson, E. O. and Whitby, K. T.: Aerosol classification by electric mobility: apparatus, theory, and applications, *Journal of Aerosol Science*, 6, 443–451, [https://doi.org/10.1016/0021-8502\(75\)90060-9](https://doi.org/10.1016/0021-8502(75)90060-9), 1975.

- Krueger, B. J., Grassian, V. H., Iedema, M. J., Cowin, J. P., and Laskin, A.: Probing heterogeneous chemistry of individual atmospheric particles using scanning electron microscopy and energy-dispersive X-ray analysis, *Analytical Chemistry*, 75, 5170–5179, <https://doi.org/10.1021/ac034455t>, 2003.
- 415 Kulmala, M., Rannik, Ü., Zapadinsky, E. L., and Clement, C. F., The effect of saturation fluctuations on droplet growth, *Journal of Aerosol Science*, 28(8), 1395–1409, [https://doi.org/10.1016/S0021-8502\(97\)00015-3](https://doi.org/10.1016/S0021-8502(97)00015-3), 1997.
- Lamb, D. and Verlinde, J.: *Physics and chemistry of clouds*, Cambridge University Press, 2011.
- Lehmann, K., Siebert, H., and Shaw, R. A.: Homogeneous and inhomogeneous mixing in cumulus clouds: Dependence on local turbulence structure, *Journal of the Atmospheric Sciences*, 66, 3641–3659, <https://doi.org/10.1175/2009JAS3012.1>, 2009.
- 420 Lenchow, D. H., Mann, J. and Kristensen, L., How long is long enough when measuring fluxes and other turbulence statistics?. *Journal of Atmospheric and Oceanic Technology*, 11, 661–673, [https://doi.org/10.1175/1520-0426\(1994\)011%3C0661:HLILEW%3E2.0.CO;2](https://doi.org/10.1175/1520-0426(1994)011%3C0661:HLILEW%3E2.0.CO;2), 1994.
- Liu, Y., Wu, Z., Huang, X., Shen, H., Bai, Y., Qiao, K., Meng, X., Hu, W., Tang, M., and He, L.: Aerosol phase state and its link to chemical composition and liquid water content in a subtropical coastal megacity, *Environmental Science & Technology*, 53, 5027–5033, <https://doi.org/10.1021/acs.est.9b01196>, 2019.
- 425 Ma, S.-S., Yang, W., Zheng, C.-M., Pang, S.-F., and Zhang, Y.-H.: Subsecond measurements on aerosols: From hygroscopic growth factors to efflorescence kinetics, *Atmospheric Environment*, 210, 177–185, <https://doi.org/10.1016/j.atmosenv.2019.04.049>, 2019.
- McGraw, R. and Lewis, E. R.: Deliquescence and efflorescence of small particles, *The Journal of Chemical Physics*, 131, 2009.
- MacPherson, J. I., Grossman, R. L., and Kelly, R. D.: Intercomparison results for FIFE flux aircraft, *Journal of Geophysical Research: Atmospheres*, 97 (D17), 18 499–18 514, <https://doi.org/10.1029/92JD00272>, 1992.
- 430 Niedermeier, D., Wex, H., Voigtländer, J., Stratmann, F., Brüggemann, E., Kiselev, A., Henk, H., and Heintzenberg, J.: LACIS-measurements and parameterization of sea-salt particle hygroscopic growth and activation, *Atmospheric Chemistry and Physics*, 8, 579–590, <https://doi.org/10.5194/acp-8-579-2008>, 2008.
- Niedermeier, D., Voigtländer, J., Schmalfuß, S., Busch, D., Schumacher, J., Shaw, R. A., and Stratmann, F.: Characterization and first results from LACIS-T: a moist-air wind tunnel to study aerosol–cloud–turbulence interactions, *Atmospheric Measurement Techniques*, 13, 2015–2033, <https://doi.org/10.5194/amt-13-2015-2020>, 2020.
- 435 Peng, C., Chen, L., and Tang, M.: A database for deliquescence and efflorescence relative humidities of compounds with atmospheric relevance, *Fundamental Research*, 2, 578–587, <https://doi.org/10.1016/j.fmre.2021.11.021>, 2022.
- Prigogine, I.: *Vom Sein zum Werden: Zeit und Komplexität in den Naturwissenschaften*. Nobelpreis 1977, R. Piper & Co. Verlag.
- 440 München/Zürich, 261pp, 1979.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2006.
- Shchekin, A. K., Shabaev, I. V., and Rusanov, A. I.: Thermodynamics of droplet formation around a soluble condensation nucleus in the atmosphere of a solvent vapor, *The Journal of Chemical Physics*, 129, 214111, <https://doi.org/10.1063/1.3021078>, 2008.
- Shchekin, A. K., Shabaev, I. V., and Hellmuth, O.: Thermodynamic and kinetic theory of nucleation, deliquescence and efflorescence transitions in the ensemble of droplets on soluble particles, *The Journal of Chemical Physics*, 138, 054704, <https://doi.org/10.1063/1.4789309>, 2013.
- 445 Siebert, H., Franke, H., Lehmann, K., Maser, R., Saw, E. W., Schell, D., Shaw, R. A., and Wendisch, M.: Probing finescale dynamics and microphysics of clouds with helicopter-borne measurements, *Bulletin of the American Meteorological Society*, 87, 1727–1738, <https://doi.org/10.1175/BAMS-87-12-1727>, 2006.

- 450 Siebert, H. and Shaw, R. A.: Supersaturation fluctuations during the early stage of cumulus formation, *Journal of the Atmospheric Sciences*, 74, 975–988, <https://doi.org/10.1175/JAS-D-16-0115.1>, 2017.
- Sloane, C.: Optical properties of aerosols of mixed composition, *Atmospheric Environment*, 18, 871–878, [https://doi.org/10.1016/0004-6981\(84\)90273-7](https://doi.org/10.1016/0004-6981(84)90273-7), 1984.
- Tang, I. N. and Munkelwitz, H. R.: An investigation of solute nucleation in levitated solution droplets, *Journal of Colloid and Interface*
455 *Science*, 98, 430–438, [https://doi.org/10.1016/0021-9797\(84\)90167-X](https://doi.org/10.1016/0021-9797(84)90167-X), 1984.
- Tang, I. N., Munkelwitz, H. R., and Davis, J. G.: Aerosol growth studies - II. Preparation and growth measurements of monodisperse salt aerosols, *Journal of Aerosol Science*, 8, 149–159, [https://doi.org/10.1016/0021-8502\(77\)90002-7](https://doi.org/10.1016/0021-8502(77)90002-7), 1977.
- Tang, M., Chan, C. K., Li, Y. J., Su, H., Ma, Q., Wu, Z., Zhang, G., Wang, Z., Ge, M., Hu, M., He, H., and Wang, X.: A review of experimental techniques for aerosol hygroscopicity studies, *Atmospheric Chemistry and Physics*, 19, 12 631–12 686, [https://doi.org/10.5194/acp-19-](https://doi.org/10.5194/acp-19-12631-2019)
460 12 631–2019, 2019.
- Titos, G., Cazorla, A., Zieger, P., Andrews, E., Lyamani, H., Granados-Muñoz, M. J., Olmo, F. J., and Alados-Arboledas, L.: Effect of hygroscopic growth on the aerosol light-scattering coefficient: A review of measurements, techniques and error sources, *Atmospheric Environment*, 141, 494–507, <https://doi.org/10.1016/j.atmosenv.2016.07.021>, 2016.
- Toon, O. B., Pollack, J. B., and Khare, B. N.: The optical constants of several atmospheric aerosol species: Ammonium sulfate, aluminum oxide, and sodium chloride, *Journal of Geophysical Research: Oceans and Atmospheres*, 81, 5733–5748, <https://doi.org/10.1029/JC081i033p05733>, 1976.
465
- Wex, H., Kiselev, A., Stratmann, F., Zoboki, J., and Brechtel, F.: Measured and modeled equilibrium sizes of NaCl and (NH₄)₂SO₄ particles at relative humidities up to 99.1%, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2004JD005507>, 2005.
- Wex, H., Ziese, M., Kiselev, A., Henning, S., and Stratmann, F.: Deliquescence and hygroscopic growth of succinic acid particles measured with LACIS, *Geophysical Research Letters*, 34, L17810, <https://doi.org/10.1029/2007GL030185>, 2007.
470
- Wilck, M.: Modal Modelling of Multicomponent Aerosols: Modale Modellierung Von Mehrkomponentenaerosolen, Verlag für Wissenschaft und Forschung (VWF), 1999.
- Wise, M. E., Martin, S. T., Russell, L. M., and Buseck, P. R.: Water uptake by NaCl particles prior to deliquescence and the phase rule, *Aerosol Science and Technology*, 42, 281–294, <https://doi.org/10.1080/02786820802047115>, 2008.