


# Supporting Information: A Cluster-of-Functional-Groups Approach for Studying Organic Enhanced Atmospheric Cluster Formation

Astrid Nørskov Pedersen, Yosef Knattrup, and Jonas Elm


Department of Chemistry, Aarhus University, Langelandsgade 140, 8000 Aarhus C, Denmark

**Correspondence:** Jonas Elm (jelm@chem.au.dk)

## S1 Cluster Structures



**Figure S1.** The  $(SA)_2(TMA)_2(\text{Carboxlic Acid})_3$  cluster geometry lowest in free energy. Calculated at the DLPNO-CCSD( $T_0$ )/aug-cc-pVTZ// $\omega$ B97X-D/6-31++G(d,p) level of theory with quasi-harmonic cutoff of  $100\text{ cm}^{-1}$ , at  $298.15\text{ K}$  and  $1\text{ atm}$ . White = hydrogen, brown = carbon, red = oxygen, yellow = sulfur, blue = nitrogen.



**Figure S2.** The  $(SA)_1(DMA)_1(Aldehyde)_1(Carboxlic\ Acid)_1$  cluster geometry second lowest in free energy (+0.62 kcal/mol). Calculated at the DLPNO-CCSD( $T_0$ )/aug-cc-pVTZ// $\omega$ B97X-D/6-31++G(d,p) level of theory with quasi-harmonic cutoff of  $100\text{ cm}^{-1}$ , at  $298.15\text{ K}$  and 1 atm. White = hydrogen, brown = carbon, red = oxygen, yellow = sulfur, blue = nitrogen.

## S2 Fluxes

**Table S1.** Sum of outgrowing channels that contribute over 2 % to the total flux and contain OOM. The SA concentration was fixed at  $10^6 \text{ cm}^{-3}$ .

System	$C_{\text{OOM}}(\text{ppt})$	$C_{\text{base}}(\text{ppt})$	Outgrowing (%)
SA-AM-OOM	1	10	97.96
	1	10000	97.95
	10	10	99.80
	10	10000	99.81
SA-MA-OOM	1	1	98.86
	1	100	99.05
	10	1	99.72
	10	100	99.74
SA-DMA-OOM	1	1	72.44
	1	10	83.77
	10	1	99.03
	10	10	99.67
SA-TMA-OOM	1	1	7.52
	1	10	12.39
	10	1	78.97
	10	10	87.22