

1 Drivers governing the seasonality of new particle formation in 2 the Arctic

3 Dominic Heslin-Rees^{1,2}, Peter Tunved^{1,2}, Diego Aliaga¹, Janne Lampilahti³, Ilona Riipinen^{1,2},
4 Annica M. L. Ekman^{4,2}, Ki-Tae Park⁵, Martina Mazzini⁶, Stefania Gilardoni⁷, Roseline
5 Thakur³, Kihong Park⁸, Young Jun Yoon⁹, Kitack Lee¹⁰, Mikko Sipilä³, Mauro Mazzola¹¹,
6 Radovan Krejci¹²

7
8
9 ¹Department of Environmental Science (ACES), Stockholm, Sweden.

10 ²Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

11 ³Institute for Atmospheric and Earth System Research (INAR), Helsinki, Finland.

12 ⁴Department of Meteorology, Stockholm, Sweden.

13 ⁵Hallym University, Chuncheon, Gangwon, Republic of Korea

14 ⁶Institute of Atmospheric Science and Climate (CNR-ISAC), Bologna, Italy.

15 ⁷Institute of Polar Sciences (CNR-ISP), Milan, Italy.

16 ⁸School of Earth Sciences and Environmental Engineering, Gwangju Institute of Science and
17 Technology, Buk-gu, Gwangju, Republic of Korea

18 ⁹Korea Polar Research Institute, Yeosu-gu, Incheon, Republic of Korea

19 ¹⁰Division of Environmental Science and Engineering, Pohang University of Science and
20 Technology, Pohang, Republic of Korea

21 ¹¹Institute of Polar Sciences, National Research Council (CNR-ISP), Bologna, Italy

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

Correspondence to: Dominic Heslin-Rees (dominic.heslin-rees@aces.su.se) and Radovan Krejci (radovan.krejci@aces.su.se)

45 Abstract

46 New particle formation (NPF) is the phenomenon wherein gaseous precursors form critical clusters of barely a
47 few nanometres in diameter, after which, under favourable conditions, these particles can grow to
48 climate-relevant sizes. Here we present measurements from 2022 to 2024 of particle and ion number size
49 distributions from the Zeppelin Observatory (ZEP), an Arctic research station situated on the western edge of
50 Svalbard atop a mountain. NPF events begin in April and continue occurring into November. The events at the
51 start of the NPF season (i.e. April/May) are considerably stronger (i.e. a larger production of nucleation mode
52 particles) ~~than~~ compared with other months of the year. The peaks in NPF strength coincide with peaks in the
53 accumulated solar insolation experienced by arriving air masses. During the summer period NPF events occur
54 on 20-40% of days each month; however, there is a consistent decline starting in June. We show that the
55 combined influence of solar insolation and the surface area of pre-existing aerosols (i.e. condensation sink, CS)
56 ~~is a~~ ~~are~~ strong predictors for the likelihood of NPF. We develop a simplified predictive model which matches the
57 frequency of NPF events identified via the classification schemes used in this study; we show that the ratio of
58 solar insolation over CS corresponds well to the frequency of NPF events (R^2 of 0.78). We show that NPF events
59 occur during the polar night (i.e. when the Sun does not pass above the horizon), and that these events are linked
60 to high altitude air masses. Furthermore, we detail the likely geographic origins of nucleation mode particles as
61 measured at ZEP. We show that NPF events are considerably more likely to originate from marine regions
62 towards the west of Svalbard, particularly the Greenland Sea, which is the marine region with the greatest
63 likelihood of originating air masses linked to an NPF day. We also show that NPF events lead to an increase in
64 the number of Aitken mode particles, indicating that ~~potentially~~ a significant proportion of the Aitken mode
65 particles originate from NPF. Of the measured NPF events, 37% ~~of them~~ exhibited nucleation mode particles that
66 grew beyond 25nm; (a diameter representing the minimum activation diameter for particles to act as cloud
67 condensation nuclei). Overall, we present a concise picture of the life cycle of nucleation mode particles in the
68 Arctic, including the effect wet scavenging has in reducing the condensation sink, which in turn promotes ~~the~~
69 ~~occurrence of NPF events~~ ~~events to occur~~.

70

71

72 1. Introduction

73 New particle formation (NPF) describes the production of secondary aerosol particles from precursor gases to
74 clusters of molecules a few nanometres in diameter. Newly formed particles belong to the nucleation mode (i.e.
75 typically <10nm) and may grow through the continued coagulation and condensation of vapours into Aitken
76 mode particles (i.e. tens of nanometres in diameter). The phenomenon of NPF has been observed globally
77 (Kerminen et al., 2018), however, in each region the chemical characteristics and controlling source and sink
78 processes leading to NPF may vary.

79 Arctic aerosol seasonality is governed by a combination of atmospheric general circulation, meteorology, and
80 several source and sink processes including the direct emission of local primary particles, NPF, the long-range
81 transport of anthropogenic emissions from lower latitudes, and the removal of particles through wet scavenging
82 (Garrett et al., 2011; Schmale et al., 2022; Tunved et al., 2013; Willis et al., 2018). One distinctive feature of the
83 annual cycle is the Arctic haze in late winter and early spring, which is characterised by a sustained increase in
84 the number of accumulation mode aerosol (i.e. particles typically around 100nm) (e.g. Shaw (1995)). Reduced
85 wet scavenging and increased transport efficiency from anthropogenic source regions at lower latitudes during
86 this part of the year give rise to this phenomenon (Garrett et al., 2011). The Arctic summertime, by contrast, is
87 less influenced by long-range transported pollutants, and the increased precipitation effectively reduces the once
88 pronounced accumulation mode. Moreover, increased photochemistry and biological activity (e.g. phytoplankton
89 blooms) provide a source of precursor gases encouraging NPF and the growth of newly formed particles
90 (Tunved, Ström and Krejci, 2013; Croft et al., 2016; Price et al., 2023); as a result, the smaller Aitken and
91 ultrafine aerosol particles increase in number.

92 Changes in aerosol size, abundance and chemical composition can perturb the radiative balance either indirectly
93 by acting as cloud condensation nuclei (CCN), altering cloud formation, brightness, and lifetime or directly
94 through their interaction with both short- and long-wave radiation (Li et al., 2022). The Arctic region can at times
95 be CCN-limited, which means that small changes in CCN concentrations could potentially impact cloud
96 properties via cloud-mediated indirect aerosol effects, which in this and similar environments could result in
97 surface warming due to enhanced long-wave cloud forcing (Garrett et al., 2004; Garrett and Zhao, 2006;
98 Mauritsen et al., 2011). Thus, local-regional scale production of aerosols during the summertime and changes to
99 this production mechanism may have significant consequences for the Arctic (Kecorius et al., 2019). Still
100 however, the impact of newly formed particles on Arctic cloud properties remains poorly quantified.

101 Depending on certain environmental factors, for example the rate of particle nucleation, the production rate and
102 volatility of condensable vapours, and the condensation and coagulation sinks, to mention a few, these newly
103 formed particles can reach climate-relevant sizes; In the Arctic, studies have suggested that particles as small as
104 25nm can participate in cloud formation by acting as CCN (Leaitch et al., 2016; Karlsson et al., 2020, 2021;
105 Pöhlker et al., 2021; Gramlich et al., 2023; Motos et al., 2023) thus affecting cloud properties and climate (i.e.,
106 by influencing the radiative balance). Motos et al. (2023) predicted the smallest particle activation diameters in
107 the summertime to be 20nm, with a modal average of 60 nm, but suggested that the dominant summertime
108 Aitken mode originating from NPF may stay in the interstitial (inactivated) phase. Other studies have indicated
109 that a substantial fraction of CCN in the Arctic are produced via NPF (Gordon et al., 2017; Merikanto et al.,
110 2009).

111 Newly formed particles can reach climate-relevant sizes; however, this depends on certain factors, e.g. the
112 production and volatility of condensable vapours and the condensation and coagulation sinks. In the Arctic,
113 particles as small as 25 nm have been shown to act as CCN (Leaitch et al., 2016; Karlsson et al., 2020, 2021;
114 Pöhlker et al., 2021; Gramlich et al., 2023; Motos et al., 2023), thereby helping to form clouds and potentially
115 influencing their radiative properties and lifetime. Motos et al. (2023) predicted a similar lower limit for
116 activation diameters of 20 nm; however, they suggested that the summertime NPF-derived Aitken mode may
117 remain interstitial (inactivated). Other studies indicate that NPF produces a substantial fraction of Arctic CCN
118 (Gordon et al., 2017; Merikanto et al., 2009).

119 Air ions are an additional focus area of aerosol effects, as they can play a role in nucleation (Hirsikko et al.,
120 2011; Kirkby et al., 2011) and can influence aerosol particles through their formation and growth mechanism.
121 Both positive and negative ions contribute to the production of newly formed particles in the Arctic, via
122 ion-induced nucleation (see Beck et al., 2021 for negative ions and Jokinen et al. (2018) for positive ions) The
123 overall contribution of ions to NPF is disputed as quantifying their impact on nucleation rates in ambient
124 conditions remains difficult. However, studies suggest that the dynamics of sub-2nm clusters is dominated by
125 neutral clusters (Hirsikko et al., 2011; Kulmala et al., 2013) . Ions can help stabilise clusters which form initially
126 (Yu and Turco, 2001) , with most particles formed from ion-induced nucleation reaching neutrality after growing
127 beyond 2.5nm (Wagner et al., 2017).

128 Observations of ultrafine particles (~3-20nm) in the Arctic (onboard the Swedish icebreaker Oden, 70°N-85°N)
129 were reported by Covert et al. (1996) and Wiedensohler et al. (1996). At Zeppelin Observatory (ZEP), Ström et

al. (2003) and Tunved et al. (2013) presented measurements of small ($>10\text{nm}$) aerosol particles showing an increased concentration and frequency of formation events during the Arctic summer. Tunved et al. (2013) suggested that the summertime, coinciding with increased photochemical production of precursor gases, and the removal of larger pre-existing aerosol, generates favourable conditions in which NPF can occur. Other Arctic sites have been the focus of research into NPF, including Villum Research Station (hereafter Villum) (see Nguyen et al. (2016)) and Summit (see Ziemba et al. (2010)), both on Greenland, and Utqiagvik, Alaska (Kolesar et al., 2017a). Other studies, e.g. Brean et al. (2023) present NPF data from several locations, including Alert, Villum, Tiksi, ZEP, Gruebadet (Ny-Ålesund), and Utqiagvik; a commonality shared by all Arctic sites was the occurrence of NPF events during the summer, and also that events coincided with air masses generally influenced by marine regions, including at Utqiagvik; and Alert whereby a correlation was observed between biogenic methane sulphate (MSA-) and summertime particle number concentrations (Leaith et al., 2013; Quinn et al., 2002). Beck et al. (2021) through a comparison of measurements from both Ny-Ålesund and Villum, showed that different nucleation mechanisms can also occur at different times of the year at these two locations; at Villum iodic acid (IA, HIO_3) was found to be the primary driver of NPF events during springtime, and sulphuric acid (SA, H_2SO_4) -ammonium (AM, NH_4^+) driven events were shown to occur in the summer. At Ny-Ålesund, NPF was driven by SA-AM during the springtime and highly oxygenated molecules (HOMs) during the summer. There is no evidence that anthropogenic precursors contribute to NPF at ZEP (Schmale and Baccarini, 2021). Instead, Dimethyl Sulphide (DMS), produced by phytoplankton, can serve as a precursor gas contributing to NPF. Marine regions with strong chlorophyll a concentrations and DMS production capacity have been linked to increased concentrations of nanoparticles, which suggests that marine biogenic sources are an important source of nucleating and condensing material (Lee et al., 2020). Observed DMS concentrations vary with air mass origin, season and residence time over different marine source regions. For example, it has been shown that air masses exposed more to the Greenland Sea, compared with the Barents Sea bring about higher DMS concentrations (Park et al., 2018). In the Arctic, there are low concentrations of nucleating agents as opposed to continental rural and urban locations (Karl et al., 2012; Pirjola et al., 2000). However, in spring and summer increased solar radiation, a lower total surface area of pre-existing particles (i.e. reduced condensation sink, CS), and coinciding phytoplankton blooms are shown to give rise to precursor gas concentrations sufficient to sustain ion-induced nucleation, and growth of NPF-induced particles beyond 20nm (Beck et al., 2021).

Recent modelling and global aerosol-climate coupling studies both laboratory and field-based have increased our understanding of biogenic sulphur and iodine pathways. For sulphur pathways, Hoffmann et al. (2016) showed

that the aqueous-phase (i.e. cloud and aerosol water chemistry), and not just the gas-phase, play an important role in the oxidation products of DMS, demonstrating the complexity of atmospheric sulphur chemistry in part due to the multiple oxidation pathways, which in turn are not represented well in climate models. Beyond sulphur-derived precursors, Finkenzeller et al. (2023) shed light on the formation mechanism for iodic acid (HIO_3) under atmospheric conditions and helps to corroborate how iodine can be produced and therefore act as a strong nucleating vapour in polar regions. Iodine-assisted NPF is also directly supported by the experimental findings shown in Baccarini et al. (2020). When iodine chemistry is included iodine acts as a missing source of low-volatility vapours, and improves the modeled Aitken mode representation (Xavier et al., 2024).

In this study, we present two and a half years' worth of measurements of particle and ion number size distributions, including three summer periods (in total data covers the period April 2022 – October 2024). Continuous measurements of sub-3nm up to ~850nm particles were performed at the ZEP, Svalbard, providing unique information of the initial stages of nucleation and the subsequent growth. NPF events were identified, formation rates were calculated, and subsequent growth was analysed. The seasonality and coinciding environmental parameters encouraging NPF events and subsequent growth have been explored in detail. In this study, we offer a simplified approach to predicting the occurrence of NPF events, using proxies for the source and sink of newly formed particles, in a region with minimal anthropogenic influence particularly in the summertime, and also in a region where extensive measurement set ups are not always feasible.

2. Materials and methods

2.1 Measurement site

Zeppelin observatory (ZEP) (78.90°N, 11.88°E, 474 m a.s.l.) is situated on the ridge of Mount Zeppelin, 2 km south from Ny-Ålesund research village on the western edge of the Norwegian Svalbard archipelago. ZEP represents regional background conditions in the Atlantic sector of the Arctic, as it is largely unaffected by local air pollution due to its location well above Ny-Ålesund and the prevailing wind patterns driven by Kongsfjorden's geography. ZEP is one of the most developed atmospheric observational sites in the Arctic where a broad set of parameters has been measured for decades. It is part of numerous regional and global monitoring networks, including the Global Atmosphere Watch (WMO/GAW), Integrated Carbon Observation System (ICOS), The Aerosol, Clouds, and Trace Gases Research Infrastructure (ACTRIS) and the co-operative programme for monitoring and evaluation of the long-range transmission of air pollutants in Europe (EMEP).

For a detailed description of the observatory, local meteorology and climatology, history and observational program see Platt et al. (2022).

2.2 Instrumentation

For this study, an array of instruments has been used to measure aerosol and ion size distributions, between 0.8 and 850 nm including a Neutral Cluster and Air Ion Spectrometer (NAIS5, Airl Ltd. Tartu, Estonia), a Nanoparticle Scanning Mobility Particle Sizer (Nano SMPS, TSI), and a Differential Mobility Particle Sizer (DMPS) system. Particle and ion number size distributions and concentrations form the core data used in this study. Several other supporting data sets were used for data analysis and interpretation and are described below. This study covers the period from 2022-04-17 to 2024-10-04.

2.2.1 Differential Mobility Particle Sizer (DMPS) system

Particle number size distributions, from 5 to 850nm, were measured using a custom-made twin-DMPS system with a closed loop sheath circulation and composed of two Hauke-type Differential Mobility Analysers (DMA). DMPS-1, with a short (5.3 cm) DMA, measured aerosol size distribution from 5 to 57 nm using sample-to-sheath air ratio close to 1:10. DMPS-2, with medium length DMA (28 cm), measured size distribution from 20 to 850 nm using sample-to-sheath air ratio close to 1:5. The aerosol particles were counted using condensational particle counters (CPCs) at a frequency of 1 Hz with the two CPCs. Particles > 3 nm were measured using the Ultrafine CPC (UCPC) TSI model 3776 (cut-off of 2.5nm) coupled to the DMPS-1. Particles > 10 nm were measured with the CPC TSI model 3010 (cut-off of 10nm) coupled to the DMPS-2. The two DMAs were run in a stepwise mode, providing two particle number size distribution cycles approximately every 27 minutes. The overlap between the size ranges measured by both DMPSs was used to check consistency and to merge both size distributions into one. Additional information can be found in Karlsson et al. (2021). The DMPS measurements were corrected for diffusional losses using the Particle Loss Calculator (PLC) according to Von Der Weiden et al. (2009) (see Fig. S1 in the supplement for details).

2.2.2 Neutral Cluster and Air Ion Spectrometer (NAIS)

NAIS was utilised to measure concentrations of ions (charged particles and cluster ions) of both polarities between 0.8 and 40 nm, and particles between 2.5 and 40 nm. The NAIS is an aerosol mobility spectrometer; when the sample is left unmodified it detects naturally-charged ions and particles, and when the corona charger is used to charge the particle population it measures all particles including the uncharged (Manninen et al., 2011; Mirme et al., 2007; Mirme and Mirme, 2013). It consists of two mobility analyser columns, with a total of 25

electrometers per column. Its lower size thresholds and high temporal resolution make it highly suitable for observing the early stages of NPF. The NAIS provides very accurate size information, regardless of the choice of the inversion algorithm used (in this case, the 25-channel V14.1 inverter algorithm) (Wagner et al., 2016). The lowest detection limit for the NAIS in particle mode is set based on the diameter of the corona charger ions, and the inability of the electrical filter to remove all naturally charged particles (Manninen et al., 2011), however, there is a broad consensus that measurements of concentrations of particles below 2.5nm are inaccurate.

The NAIS at ZEP was installed with a very short metallic inlet, less than 1m long, to minimise diffusional losses. The inlet pointed slightly downward, and a metallic rain shield was attached at the end, to ensure that rain did not enter into the instrument. The NAIS inlet was heated to prevent ice blocking the inlet, and also to limit water condensing inside the tubing. The NAIS was serviced and cleaned once a year. Additional aspects of the NAIS observations, including the apparent overestimation of particle number concentrations when compared to DMPS measurements (see also Kangasluoma et al., 2020) and the loss of negative cluster ions can be found in section S2.6 of the supplementary material.

2.2.3 Complementary observations

A Nanoparticle Scanning Mobility Particle Sizer (Nano SMPS, TSI) comprising of a nano-differential mobility analyser (nano-DMA, TSI 3085, USA) and an ultrafine CPC (TSI 3776, USA), provided size distributions of nanoparticles (3-60nm). See Lee et al. (2020) for further details. Atmospheric Dimethyl Sulphide (DMS) was measured using an analytical system in which DMS is trapped and eluted, before being quantified using gas chromatography equipped with a pulsed-flame photometric detector. The detection limit is approximately 1.5 pptv given air samples of around 6 litres. For more details, see Jang et al. (2016) and Park et al. (2018). Meteorological parameters including visibility, relative humidity, wind speed and direction, ambient temperature and pressure were also further utilised throughout the study.

2.3 Data Analysis of aerosol measurements

For processing of the NAIS data the *nais-processor* was used (<https://github.com/jlpl/nais-processor>, v0.0.30). The *nais-processor* Python package corrected for diffusional losses in the inlet (Gormley and Kennedy, 1948), applied an ion mode calibration (Wagner et al., 2016), converted ion mobility to particle diameter, remapped the distribution to a new size grid, and adjusted the data to standard conditions (273.15 K, 101325 Pa).

Further information on the removal of artefacts can be found in Sect. S2 of the supplementary material. In addition, a comparison between the particle concentrations measured by NAIS and the DMPS-system can be

found in Sect. S2.5; we report a significant overestimation of particle number concentrations as measured by the NAIS in comparison to the DMPS-system, particularly at the smallest diameters (see Fig. S9-S11). It should be noted that the particle number size distributions presented in this study represent that of dry aerosol particle sizes. The measured particle size distributions for both the NAIS and DMPS were not corrected for hygroscopic growth, and thus do not reflect ambient conditions.

When assessing the qualitative and quantitative nature of NPF events, it is important to distinguish between on the one hand the production of newly formed particles (~2-4nm) and on the other hand subsequent particle growth. To qualify as an event, not only does formation of ultrafine particles need to be present, but also the subsequent growth of the particles. Growth can last for several hours and in some cases multiple days, with the latter timescale suggesting that the NPF events occur on a regional scale (Kecorius et al., 2019; Ström et al., 2009). In this study, we distinguish between (1) *on-site formation*, defined as occasions when the concentration of 2-4nm particles clearly rises above the background level; and (2) *formation and subsequent growth*, defined as a combination of *on-site formation* followed by subsequent growth. The latter is identified by visual inspection of the behaviour of the growing nucleation mode during the time it is present until when it disappears.

2.3.1 NPF Classification

New particle formation events were classified using schemes described by Dal Maso et al. (2005) and Aliaga et al. (2023). The classification schemes are used complementary. The classification by Dal Maso et al. (2005) focuses more on formation and growth, and is perhaps more intuitive to understand; however, it struggles with a large number of undefined cases. On the other hand, the nano-ranking analysis developed by Aliaga et al. (2023) can be used to explore the NPF intensity and focuses on the formation of nucleation mode particles as opposed to the subsequent growth. The Aliaga-method classifies NPF events into 3 groups based on the overall ranking of NPF intensity, avoiding the problem with unidentified event days (see Fig. S16 for the groups). The two classification methods can be compared with one another (see Fig. S14). We used the method for the intensity parameter and the diameter range that showed the best comparison with the Dal Maso et al. (2005) i.e. the daily maximum for the total number concentration of 2.8 nm and 5 nm particles ($\Delta N_{2.8-5 \text{ nm}}$).

Dal Maso classification system

In the Dal Maso et al. (2005) scheme an NPF event is defined if there exists a nucleation mode (3-25nm) which prevails for several hours and shows signs of growth; these events are further subdivided into three separate classes, namely, Class Ia and Ib, and Class II. Class I and Class II events are separated on the basis of whether

the growth and formation rates, of the well-behaved size distribution, can be determined with good confidence. Class I is further divided into Class Ia and Class Ib depending on whether there are pre-existing particles obscuring the newly formed mode (Dal Maso et al., 2005). Undefined events are days where there are sporadic occurrences of nucleation-mode particles, and this classification is used to separate clear events from clear non-events. For examples of the different classifications and their respective averages see Fig. S12 and S13 respectively in the supplement.

In this study, undefined days include days where the occurrence of nucleation mode particles (3-25nm) is linked to windblown snow and situations when ZEP is enveloped by clouds. For more details regarding these sampling conditions see Sect. S2.1 and S2.2 in the supplement for in-cloud-sampling and windblown snow events respectively. As a result, days classified as undefined are ones which experience bursts of nucleation mode particles, either from windblown snow (see Fig. S4), in-cloud conditions (see Fig. S2), from non-growing NPF or a combination of all three. There is a definite increase in the concentration of particles with diameters around 5nm when the NAIS measures in conditions of low visibility and this is not observed in other instruments (see Fig. S3). For windblown snow events, particles with diameters less than 20nm observed considerable increases in concentrations when the wind speed passed roughly 10ms⁻¹ (see Fig. S5). Non-events are days that display no nucleation mode (3-25nm). It is important to note that this scheme assumes that particle formation occurs over a geographically wide area more or less simultaneously. Identification and classification were performed using daily surface plots of particle number size distribution based solely on NAIS data, that is, 2.5-40nm.

It should be noted that there is a degree of subjectivity to this classification approach, as to what is considered a certain class. The addition of the NAIS, in combination with the DMPS-system, made it easier to detect events using surface plots, especially when the growing mode struggled to surpass ~6nm. However, we were not able to identify any NPF events with the NAIS that we could not identify with the DMPS-system, as close inspection of the number concentration of the lowest size bins of the DMPS-system was sufficient to recognise NPF events.

Nanoparticle Ranking Analysis

The Nanoparticle Ranking Analysis method developed by Aliaga et al. (2023) provides a more continuous method to characterise atmospheric NPF, allowing users to gauge the strength of NPF. The nanoparticle ranking analysis was utilised using the daily maximum concentration of 2.8-5nm particles ($\Delta N_{\text{max}, 2.8-5}$), from the negative channel of the NAIS, as the intensity parameter (note that no background value was used, unlike in Aliaga et al. (2023), due to the low concentrations measured at ZEP). In this study, periods which experienced windblown

snow or in-cloud events were removed before applying the Nanoparticle Ranking method (see Figs. S7 and S8), as opposed to the Dal Maso et al. (2005) approach which classified them as undefined days. Special care was taken to clean the data before utilising the nanoparticle ranking analysis method, however it may still be the case that not all in-cloud and windblown snow events were removed successfully (see Sect. S2.1 and S2.2 in the supplement). Windblown and in-cloud events can influence the daily maximum without necessarily reflecting NPF activity. The strength of NPF activity was split into 3 groups depending on the intensity (i.e., $g1$, $g2$, $g3$), where $g3$ is the most intense event, followed by $g2$ and then $g1$; see Aliaga et al. (2023) for more details (see Fig. S15 & Fig. S16).

2.3.2 Growth rates

The growth rate (GR) describes the rate of change of the modal diameter during an NPF event. For the GR estimation, NAIS and DMPS size distributions were merged at the maximum diameter of the NAIS size range (i.e. 40nm). The NAIS and DMPS data were both interpolated to 15-minute arithmetic means using linear interpolation. The start and end of each NPF event including their sequential growth (i.e. (2) *formation and subsequent growth*), were estimated by visual inspection. The growing nucleation mode was isolated, and the GR was estimated for the full duration of the NPF event (see Fig. S17). For each size bin, we applied a Gaussian filter ($\sigma = 1$) and found the time in which the concentration of a particular size bin experienced the highest maximum rate of change. Essentially, the evolution of the nucleation mode was tracked using the timings for each identified peak in the rate of change of the growing nucleation mode diameter. The GR was calculated using a simple linear ordinary least squares (OLS) method.

It should be noted that the aforementioned issues concerning the overestimation of particle number concentrations for the NAIS measurements do not impact the calculation of GR as it only examines the changes to the diameter of the nucleation mode. Very few events grew beyond the limit of the NAIS (i.e. 40nm), hence the impact of a diameter jump from NAIS to DMPS was not considered. We tried to measure the GR of all NPF events (e.g. Class 1a, 1b and II), even though by definition Class II events lack the strength and consistency, making it difficult to calculate GRs (Dal Maso et al., 2005). The GRs for some Class II events were not estimated due to uncertain tracking of the nucleation growth.

2.3.3 Particle Formation Rates

The formation rates namely J_{3-7} , J_{7-25} , and J_{3-25} ($\text{cm}^{-3}\text{s}^{-1}$) were estimated using the following equation:

$$J_{i-j} = \frac{dN_{i-j}}{dt} + \text{CoagS} \times N_{i-j} + \frac{GR}{\Delta d_p} \times N_{i-j}, \quad (1)$$

where J_{i-j} is the formation rate of particles between diameters i - j nm, CoagS is the coagulation sink of the pre-existing particle population, GR is the growth rate between i - j nm, Δd_p is the difference in diameter between i and j in nm and N_{i-j} is the total concentration of particles between i and j nm (Fig. S18 represents an example for calculating formation rates for particles between 3-7nm).

The Python GUI, *npf-event-analyzer*, was used to provide additional complementary estimations of the GRs and formation rates for the NPF events (i.e. in addition to the OLS fitting method). When using the *npf-event-analyzer* the max concentration method was used for the estimations of GR and J.

2.3.4. Condensation sink

The condensation sink (CS) is defined as the rate at which non-volatile vapours condense onto pre-existing particles (Kulmala et al., 2012). The twin DMPS system (5-850nm) was used to calculate CS, for SA vapour. The method for the determination of CS is described by Dal Maso et al. (2002), and uses the following equation:

$$CS = 2\pi D \int_i^{d_{p,i} \max} d_{p,i} \beta(d_{p,i}) N(d_{p,i}) dd_{p,i}, \quad (2)$$

where $d_{p,i}$ is the diameter of a particle in size class i , N_i is the particle number concentration in the respective size class, and D is the diffusion coefficient of the condensing vapour (in this case condensing vapours were assumed to have H_2SO_4 diffusion properties). We used the transition regime correction factor β from Fuchs and Sutugin (1970).

2.3.5 Concentration of condensable vapours and source strength

The concentration of vapours (C_v) required to sustain the calculated GRs and their respective source rates (Q) were estimated using the following equations, and assuming the condensation of H_2SO_4 .

$$C_v = A \times dD_p / dt. \quad (3)$$

where A is the constant $1.37 \cdot 10^{-7} \text{ h cm}^{-3} \text{ nm}^{-1}$ representative for the molecular properties of SA and dD_p/dt is the rate of change of the nucleation mode i.e. GR from the start of the NPF until it stops growing (see Sect. 3.1.5 below for further details). See Dal Maso et al. (2005) for more details. The constant A incorporates the probability that sulfuric acid molecules will ‘stick’ to particles upon collision, the change in volume which occurs after collision and condensation, and also the frequency of collisions (Fiedler et al., 2005).

356 2.4 Airmass analysis

357 2.4.1 Transport model

358 Airmass back trajectory analysis was performed using the Hybrid Single-Particle Lagrangian Integrated
359 Trajectory model (HYSPLIT V5.2.1) (Draxler et al., 1998; Stein et al., 2015). Ensemble back trajectories were
360 initialised every hour starting at the latitude and longitude of ZEP, and at a height of 250m above ground level
361 (AGL). The ensemble was generated by offsetting the meteorological grid point by one in the horizontal (i.e. $dx = 1.0$, $dy = 1.0$) and by the default offset of 0.01 sigma units in the vertical ($\sim 250\text{m}$), thus generating 27 back
362 trajectories for all possible offsets in X, Y and Z. The starting height of 250m ensured that the starting location
363 of all the ensemble members was at the surface or above. The back trajectories were calculated for 5 days back
364 in time. Global Data Assimilation System (GDAS) $1^\circ \times 1^\circ$ archive data (<http://ready.arl.noaa.gov/archives.php>,
365 last access: 08-08-2024) was used as the meteorological fields. No averaging of the back trajectories was
366 performed, reducing them to single back trajectories. HYSPLIT was run from the Desktop using the Python
367 package *pysplit* (<https://github.com/mscross/pysplit>).

369 The accumulated solar flux in the most recent 6 hours prior to arrival was used for the solar insolation term used
370 throughout this study. The solar flux was taken from the original GDAS-derived HYSPLIT output. Furthermore,
371 only endpoints within the mixed-layer were used as this is when air masses are influenced by the surface. The
372 last 6 hours were used, as opposed to a longer duration, to place more significance on solar radiation closer to the
373 receptor; an accumulated solar flux consisting for a longer duration could lead to misleading results.

374 2.4.2 Chlorophyll a data

375 Daily mapped chlorophyll a data derived from satellite observations (Aqua MODIS) (last accessed: 2024-10-04,
376 downloaded from <https://oceancolor.gsfc.nasa.gov/l3/>) were utilised along with the HYSPLIT output to estimate
377 the chlorophyll a exposure. Exposure of the air masses to chlorophyll a was calculated based on the equation
378 developed by Park et al. (2018), which linked chlorophyll a with DMS measurements on site at ZEP. Chlorophyll
379 a exposure was essentially used as a proxy for DMS emissions. DMS in turn can form H_2SO_4 via SO_2 , although
380 this was not explicitly addressed by Park et al. (2018). It should be noted that DMS is not only a precursor to
381 H_2SO_4 , but also MSA. Park et al. (2018) utilised the relationship between DMS and air mass exposure to
382 chlorophyll a to account for the differences in the species-specific phytoplankton population, between the
383 Greenland Sea and the Barents Sea. For more details on the calculation of the chlorophyll exposure see Sect.
384 12.

385 2.4.3 Nucleation site estimation

386 The method utilised here is similar to the NanoMap analysis developed by Kristensson et al. (2014) in which the
387 geographic position of where new particle formation took place was estimated based on the particle number size
388 distribution measurements during NPF events and HYSPLIT back trajectories. In this study, the duration of the
389 nucleation mode was calculated. For every hour of the NPF event, back trajectories were initialised. The length
390 at which the back trajectories were initialised depended on how long the event had lasted. In short, i hours after
391 the event began, the back trajectory will have been initialised to run i hours backwards. So, for every hour
392 except the start time the back trajectories were initialised for the following length of time: *back trajectory length*
393 = *duration* - (*end time* - t), where $t \in$ (end to start times). Hence, for an NPF event that lasts 7 hours, 6 back
394 trajectory ensembles were calculated for varying lengths from 7 hours to 1 hour long.

395 2.4.4 Air mass marine regions

396 The endpoints for every 5-day back trajectory ensemble were assigned to a marine region e.g. Arctic Ocean,
397 Greenland Sea, Barents Sea, Norwegian Sea and North Atlantic Ocean (see Fig. S21 in supplement for defined
398 regions). Days were assigned to a specific marine region based on the cumulative residence time in the specific
399 sector. A threshold of 0.5 (50%) was used as a qualifier. Days without clear source preference, i.e. the airmasses
400 that did not reach the 50% threshold for either of the pre-defined source regions, were classified as mixed.

401 3. Results & Discussion

402 3.1 Seasonality and interannual variability

403 3.1.1 Annual cycle of NPF events

404 The 848 days of valid NAIS measurements (2022-04-17 to 2024-10-04) are classified according to the Dal Maso
405 et al. (2005) classification system. 157 days are classified as NPF events, either Class Ia events (28), Class Ib
406 (19) or Class II (110). A total of 438 days are classified as Undefined, and the remaining 271 days are considered
407 to be either Non-event days (253) or consisting of too little data (i.e. defined as <12 hours of data per day) (18)
408 (see Fig. 1). Overall, this meant that 19% of days with valid NAIS measurements experienced an NPF event (i.e.
409 Class Ia, Ib or II). The majority of days (52%) are classified as undefined due to the appearance of a
410 non-growing nucleation mode that persisted for some hours. The undefined days are further sub-divided into
411 those containing windblown snow events (19% of all data), in-cloud events (8% of all data) (see Fig. S6 in the
412 supplement). Of the 438 Undefined days, 72 are a result of non-growing bursts (i.e. 17% of the Undefined days).

413 The monthly frequency of NPF events (i.e. class Ia, Ib or II) shows events first appearing in April (see Fig. 1),
 414 reaching a peak, in terms of the occurrences per month, in May to July with approximately 30 - 40% of days
 415 experiencing NPF. The daily occurrence decreases during the month of June compared with the preceding and
 416 succeeding months, a consistent finding throughout the three years of data. After July, the frequency of events
 417 declines towards the end of summer and into autumn, where NPF events cease to be present beyond November.
 418 NPF events were observed as late in the year as November, which is interesting given that this is during the polar
 419 night when the Sun stays below the horizon (as witnessed by ZEP) and thus ZEP experiences negligible solar
 420 radiation. Multiple NPF events were observed during the polar night period (see Sect. 3.1.3 for more details).
 421 April 2022 appeared anomalous perhaps given that the measurements for that month were not an entire month as
 422 the measurements started on 17th April 2022.
 423 Prior studies, at ZEP, which used slightly different event classification schemes, reported the occurrence of NPF
 424 events to be within the range of 18%-23% (Dallósto et al., 2017; Heintzenberg and Leck, 2012; Lee et al., 2020).
 425 Nieminen et al. (2018) reported NPF day frequencies of 14% (Mar-May), 34% (Jun-Aug) and 6.6% (Sep-Nov)
 426 and 0% (Dec-Feb). Lee et al. (2020) observed the highest frequency in terms of the formation and growth of
 427 N_{3-25} particles in June (>40%), with the NPF frequency in May and July being ~40% and <40%. Dome-C, in
 428 Antarctica and Alert, on Greenland are two other polar sites with annual NPF frequencies between 10-20%
 429 (Nieminen et al., 2018), comparable to ZEP.




Fig 1: Number of days per month classified as either Class Ia (red), Ib (orange), II (yellow), Undefined (green) and Non-event (grey). The left-hand axis is the number of days classified based on the total number of valid measurement days (i.e. Total). The right-hand side is the fraction of events (i.e. Class Ia, Ib, and II) over the total number of valid measurement days (i.e. Class Ia, Ib, II, Undefined and Non-event). The fraction is given by Event/Total. Almost three years of Neutral cluster and Air Ion Spectrometer (NAIS) particle number size distribution data is classified using the Dal Maso et al. (2005) classification system.

430

431 3.1.2 Interannual variability of NPF intensity, solar insolation, DMS and CS

432 The interannual variability of the following main parameters, namely NPF intensity, solar insolation, DMS
 433 concentration and the CS, are explored in figures 2a-b. A proxy for the intensity of events, as opposed to the
 434 frequency, was determined by utilising elements of the nanoparticle ranking analysis developed by Aliaga et al.
 435 (2023). The maximum daily concentration in 2.8 – 5nm particles ($\Delta N_{max\ 2.8-5}$), was used to represent the strength
 436 of NPF activity (shown to correlate well with calculated J) i.e. the NPF intensity. See Sect. 2.4.1 in the Methods
 437 for more details concerning the nanoranking method and about the definition of solar insolation.




Fig. 2: Time series of (a) the NPF intensity parameter ($\Delta N_{max,2.8-5}$; black dots), defined as the daily maximum particle number concentration between 2.8 and 5 nm. The black line shows the centred 4-day running mean. Orange bars indicate daily solar insolation accumulated over the preceding 6 h for air masses arriving at ZEP within the mixed layer (Sect. 2.4.1); faint yellow bars show the same for all air masses. The red shaded region is there for comparison to the yellow and orange bars and represents six times the average hourly clear-sky global horizontal irradiance (GHI) at ZEP. (b) Valid DMS measurements (green) and the condensation sink (CS; black dots, daily means) with its centred 4-day running mean (black line). Blue dashed lines indicate polar night periods.

438

439 $\Delta N_{max\ 2.8-5}$ shows various peaks throughout the annual cycle, however, the one striking feature is that every year
 440 $\Delta N_{max\ 2.8-5}$ has its first and highest peak occurring in late spring and the beginning of summer (i.e. end of April in
 441 2022, beginning of June in 2023, and towards the end of May in 2024) (see the red curve in Fig. 2a)). In terms
 442 of the annual cycle in NPF event frequency (see Fig. 1), the first peak in NPF event occurrences matches the
 443 April-June peak in increased $\Delta N_{max\ 2.8-5}$ (Fig. 2a)). However, the succeeding peak in NPF event frequency (i.e. in
 444 July, in Fig. 1) coincides with reduced intensity ($\Delta N_{max\ 2.8-5}$). The seasonality of the $\Delta N_{max\ 2.8-5}$ value follows that of
 445 the solar insolation, and for the years 2023 and 2024, the maximum NPF strength/intensity (i.e. $\Delta N_{max\ 2.8-5}$)

446 coincides with the greatest mean solar insolation; a much better relationship is observed when selecting for air
 447 masses within the mixed layer as this is when arriving air masses are influenced by the surface (i.e. the orange
 448 bars in Fig. 2a)). Averaging the intensity parameter and correlating it to the solar insolation calculated based on
 449 the entire air mass and the parts within the ML shows that selection of air masses within the ML leads to a better
 450 correlation (see Fig. S27 in supplement). The annual cycle for the calculated CS exhibits a sustained maximum
 451 in late winter and early spring due to the Arctic Haze (i.e. Feb. - May), and the CS stays high into the summer
 452 before reaching a minimum in autumn, typically the cleanest part of the year (Tunved et al., 2013) (see Fig. 2b)).
 453 During the Arctic Haze period, long-range transported air masses from Eurasia bring aged aerosol leading to the
 454 higher CS. At the same time this period is characterised by higher sulphate and SO₂ concentrations, mainly of
 455 anthropogenic origin (Platt et al., 2022), and hence there are NPF events occurring and high increased NPF
 456 intensity. One additional aspect is that it is clear that increases in $\Delta N_{max\ 2.8-5}$ occur during periods where there was
 457 a slight decrease in CS, below that of the seasonal CS average. For example, the peak in $\Delta N_{max\ 2.8-5}$ in April in
 458 2022 (see Fig. 2b)), which occurs slightly prior to the maximum solar insolation, can potentially be explained by
 459 the reduction in the CS during that period.

460 The solar intensity is a controlling factor in the production of SA (i.e. H₂SO₄) via the oxidation of SO₂. In the
 461 Arctic, DMS acts as a significant source of sulphate, however other sulphur compounds can be oxidised e.g.,
 462 carbon disulphide and hydrogen sulphide (Pei et al., 2021). The coagulation sink (CoagS) acts as a sink for the
 463 newly-formed particles, and the CS acts as a sink for nucleating vapours, thus high CoagS and CS can inhibit
 464 NPF. NPF days with greater intensity (i.e., $\Delta N_{max\ 2.8-5}$) are skewed towards the end of spring and beginning of
 465 summer. Similarly, solar insolation is also shifted towards the start of summer, even though the theoretical solar
 466 maximum has a symmetrical distribution centred around the middle of summer (see red shaded region in Fig.
 467 2a)). The skewed pattern in both the solar insolation and NPF intensity ($\Delta N_{max\ 2.8-5}$) can perhaps be explained by
 468 annual changes in cloud cover (see Maturilli and Ebell (2018) for the frequency occurrence of cloudy sky,
 469 increasing to ~80% during the summer from an April minimum of ~40-60%). Late autumn experiences
 470 numerous, but relatively weaker, NPF events, with peaks in $\Delta N_{max\ 2.8-5}$ (~100-500cm⁻³) every year at the end of the
 471 summer despite the relatively small solar insolation. The transition period between summer and haze (i.e. Oct. –
 472 Jan.) is typically the cleanest part of the year, and thus exhibits the lowest CS (Tunved et al., 2013). During this
 473 part of the year the small amount of solar insolation and vapours present are enough to trigger NPF, possibly as a
 474 result of the low CS.

494 Tiksi, Russia, also observe NPF events without the sites experiencing any solar radiation (Asmi et al., 2016;
495 Kolesar et al., 2017b). However, in these two cases the suggestion is that anthropogenic emissions of
496 semi-volatile gases from nearby oil fields or the build-up of anthropogenic emissions during the haze period can
497 explain these events; an interpretation which is not applicable for ZEP.

498 The vertical component of the arriving back trajectories associated with these polar night events shows that air
499 masses originated from higher altitudes (Figure S17, in the supplement), and that these air masses descend
500 before arriving at ZEP. One possible explanation for these types of events could be that precursor gases,
501 originally emitted from the surface, are transported to high altitudes where the CS is sufficiently low enough to
502 allow them to survive long enough to be nucleated on descent to ZEP. The air masses travel further south and
503 higher in altitude, where the small amount of solar radiation potentially allows for photochemical production of
504 nucleating vapours to build up.

505 3.1.4 Start times and duration throughout the year

506 The median duration of NPF events, including their subsequent growth, is approximately ~ 10 hours and seems
507 to be fairly consistent across the annual cycle, with a decrease towards the autumn (see Fig. 3a)). Of all events,
508 75% > 7 hours, and 90% > 4 hours. During the sunlit period, the starting time of events is typically around 09:00
509 (UTC) for the period April to August. From September onwards, the onset of the observed NPF shifts to later in
510 the day (Fig. 3b)). For the period August to October, onset of NPF is approximately 8 hours after sunrise at Ny
511 Ålesund (see the gradient of the two slopes in Fig. 3b) remains constant but the start time is shifted by 8 hours).
512 This could reflect that during the NPF season there is a delay of approximately 8 hours from the minimum solar
513 intensity to the onset of NPF, and this could be related to the time it takes for precursor gases to accumulate.

514 Lee et al. (2020) suggested that the duration of NPF at ZEP was approximately 6-7 hours, with the longest
515 duration in summer. In addition, NPF start times were 13:00 – 14:00 (local time, which is 11:00 – 12:00 UTC in
516 the summer time). The start times observed at ZEP, here in this study, were typically 2 hours earlier, perhaps as a
517 result of the slightly lower minimum particle diameter detection and not the same definition of the onset of
518 nucleation.




Fig 3: (a) the duration of the entire growth of newly formed particles from the first measurable size bin i.e. 2.5nm to the largest size bin (b) the average start time of the observed new particle formation (NPF) (monthly mean displayed by the cross and median by the red line), along with a shaded region representing the 25th and 75th percentiles. Along with the estimated time the global horizontal irradiance (GHI) goes beyond zero (i.e. daybreak). The average time of day break for the months April until July is not applicable, since the sun is constantly above the horizon (as experienced by ZEP). Time is given in terms of coordinated universal time (UTC).

3.1.5 Seasonality of Growth Rates and formation rates

The average, mean (median), GRs for all the events is 2.2 (1.5) nmhr⁻¹. For class Ia, Ib, and II events the average GRs are 1.4 (1.0), 1.7 (1.2), 2.6 (1.2) nmhr⁻¹ respectively. The largest GRs occur during July 4.0 (2.1) nmhr⁻¹. The polar night events exhibit a significantly reduced GR of 0.43 (0.35) nmhr⁻¹ (see Fig. 4 a)). Using the *npf-event-analyzer* for the Class Ia events the GRs are 0.9 (1.1), 2.1 (2.6), 4.1 (4.4) nmhr⁻¹ for 3-7, 3-25 and 7-25 nm diameter ranges.

The GRs presented here agree well with earlier estimates of GRs by Nieminen et al. (2018) reporting a GR_{10-25nm} at ZEP within the range of 1.2 -1.6 nmhr⁻¹. Kerminen et al. (2018) report a median GR for the Arctic of 2.3 (0.23-4.1, 5th and 95th percentiles) nmhr⁻¹, the site type (i.e. globally) with the smallest GRs.





Fig 4: a) the calculated growth rates (GRs) b) calculated median formation rates for particles between 3 and 7nm ($J_{med, 3-7}$) throughout the measurement period are represented by box-whiskers, which display the monthly growth rate averages and include the median and 25th - 75th percentiles. The green box-whiskers refer to the average GRs/ J_{3-7} s from the start of the NPF until it stops growing. The black line cross represents the mean, and the grey lines join the median values. Diamonds represent outliers.



528

529 Formation rates for $J_{\text{median}, 3-7\text{nm}}$ are $0.14 \pm 0.23 \text{ cm}^{-3}\text{s}^{-1}$ (mean $\pm$ std) for all events in which calculations of J and
 530 GR could be ascertained (see Fig. 4 b) for the seasonality). Formation rates are generally greater in May with
 531 2023 not exhibiting this maximum in May. The formation rates for $J_{3-7\text{nm}}, J_{3-25\text{nm}}, J_{7-25\text{nm}}$ are estimated to be $0.28 \pm$
 532 $0.33, 0.40 \pm 0.35$ and $0.31 \pm 0.28 \text{ cm}^{-3}\text{s}^{-1}$ (mean $\pm$ std) for class Ia events (using the max concentration method
 533 and *npf-event-analyzer* see Sect. 2.3.3 in the Methods).

534 Previous studies also estimate J for various diameters at Ny-Ålesund and ZEP. Beck et al. (2021) reports a $J_{1.5}$
 535 equal to $0.27 \text{ cm}^{-3}\text{s}^{-1}$ (Beck et al., 2021). Nieminen et al. (2018) presents the seasonal medians for J_{nuc} of
 536 nucleation-mode particles (10-25nm) varying from $0.08 \text{ cm}^{-3}\text{s}^{-1}$ in spring, to $0.032 \text{ cm}^{-3}\text{s}^{-1}$ in summer and to
 537 $0.0066 \text{ cm}^{-3}\text{s}^{-1}$ in autumn. Lee et al. (2020) gives $J_{3-7}, J_{7-25}, J_{3-25}$ values ranging between 0.001-0.54, 0.003-0.5,
 538 0.007 - $0.61 \text{ cm}^{-3}\text{s}^{-1}$ respectively, and the averages for $J_{3-7}, J_{7-25}, J_{3-25}$ are 0.04, 0.09 and $0.12 \text{ cm}^{-3}\text{s}^{-1}$. Kerminen et al.
 539 (2018) report a median Arctic J (i.e. varying size ranges) of $0.51 \text{ cm}^{-3}\text{s}^{-1}$.

540 In comparison to GRs and Js at other sites (e.g. Kerminen et al. (2018); Nieminen et al., (2018)), we see that
 541 processes within the Arctic, as observed at ZEP, are significantly slower. In urban environments, compared with
 542 more remote ones, estimations for J are significantly larger due to the larger coagulation sink that needs to be
 543 overcome (Cai and Jiang, 2017); here, the coagulation sink is much lower than in urban environments.

544 3.1.6 Seasonality of Condensable Vapours

545 The concentration of condensable vapours (C_v) required to sustain the GRs is estimated using equation (3) and
 546 the calculated growth rates for the various classes of events (i.e., Class Ia, Ib and II). We show that C_v increases
 547 until July before decreasing during the autumn and into the winter months. It is striking to note that the seasonal
 548 variation is similar to that of the sum of the measured concentrations of potential aerosol precursor gases
 549 measured in Ny-Ålesund during a 2017 campaign (see Fig. 5). The estimated C_v is also of a similar magnitude
 550 to the combined sum of measured HOMs, sulphuric acid (H_2SO_4), methanesulphonic acid (MSA) and iodic acid

551 (HIO₃). For Ny-Ålesund, Beck et al. (2021) report increasing IA, SA and MSA concentrations from the start of
 552 the sunlight period, and high concentrations prevailing before decreasing around mid-June. HOMs
 553 concentrations averaged around $\sim 1.8 \cdot 10^6 \text{ cm}^{-3}$ during the spring before increasing to $2.3 \cdot 10^6 \text{ cm}^{-3}$ in June. As
 554 a ratio of the total available precursor vapours HOMs account for approximately 0.78 of the entire concentration
 555 of precursor vapours after June and prior to June this fraction approximately 0.24 (see Fig. S24). It can therefore
 556 be argued that the composition of condensable vapours undergoes a dramatic transformation around the start of
 557 June, and the composition of newly formed particles during the initial spring peak is different to that of the
 558 subsequent peaks after June (see Fig. 1 & 2). Boyer et al. (2024) measured the concentrations of condensable
 559 vapours in the Arctic, however at higher latitudes ($>80^\circ\text{N}$). The estimated Cv was on average 5.9 times greater
 560 than the combined concentrations of the measured precursor gases given in Boyer et al. (2024) (see Fig. S23 in
 561 supplement). At higher latitudes ($>80^\circ\text{N}$), there is potentially less photochemistry and biological activity,
 562 moreover, the potential sources of precursor gases are more limited (Brean et al., 2023), and may be more
 563 influenced by the sea ice cover, further reducing an available and giving rise to the difference between the
 564 estimated Cv at ZEP and total concentrations during MOSiAC.




Fig 5: Estimated concentration of condensable vapours (Cv, green) compared with measured concentrations: the monthly means (crosses) and medians (thick line), along with the 25th and 75th percentiles (shaded region) for both Cv and Ny-Ålesund measurements. Arctic precursor vapours namely sulphuric acid (H₂SO₄, red), methanesulphonic acid (MSA, gold) and iodic acid (HIO₃, purple) and combined (grey) from Beck et al. (2021) are displayed. The estimated Cv measurements cover the years 2022 to 2024, whilst the Svalbard measurements from Beck et al. (2021) cover the period 2017-03-22 to 2017-08-23.

566 3.2 Positive and negative ions during NPF

567 The polarity of air ions, which aid in nucleation, provides information about the composition of the newly
 568 formed aerosol particles. Cluster air ions are typically defined between 0.8-1.7nm, whilst intermediate ions range
 569 from 1.6-7.4nm. Intermediate ions are created in the initial stage of nucleation and are typically formed when
 570 neutral nanoparticles encounter cluster ions and acquire their charge (Tammet et al., 2014). Negative ions can be
 571 linked to IA and SA to name a couple of compounds. Unfortunately, due to an instrumental problem related to a
 572 consistent loss of the negative cluster ions experienced during the measurement campaign (see Sect. S2.6 in the
 573 supplement) it was not possible to measure the concentration of negative ions less than 2nm.

574 The total number of positive (N_{2-4nm}^+) and negative (N_{2-4nm}^-) small intermediate ions (2-4nm) within the growing
 575 mode are calculated for each NPF event. In May, at the start of the NPF season, we observe the smallest fraction
 576 of positive small intermediate ions compared to the total number of small positive intermediate ions (i.e. the
 577 mean and median ratio, $N_{2-4nm}^+ / (N_{2-4nm}^- + N_{2-4nm}^+) \approx 0.2$). The low N_{2-4nm}^+ /total ratio in May perhaps relates to
 578 negatively charged molecules dominating the initial stages of ion-induced nucleation (i.e., compared with
 579 positive ions). Moreover, the low ratio in May could be related to the greater NPF intensity (i.e. larger
 580 production of negative nucleation mode particles in May). However, towards the end of the NPF season (i.e. Sep.
 581 - Nov.) there is a shift towards a growing nucleation mode (e.g., 2-4nm) more dominated by positive small
 582 intermediate ions, as opposed to negative small intermediate ions (see Fig. 6). The tendency for small
 583 intermediate ions (2-4nm) to exhibit a positive charge appears from September onwards, which is also when
 584 there begins to be limited solar insolation and the occurrence of polar night events (i.e. late Oct. and Nov.), it is
 585 also, and perhaps more importantly, when the composition of condensable vapours changes to being dominated
 586 by HOMs (see Sect. 3.1.6.). However, it needs to be noted that there are few events during September and
 587 beyond (29 in total). Despite the lack of information about the cluster ions, the shift in the overall charge of the
 588 small intermediate ions, from more negative to more positive, potentially hints at a seasonal change in the
 589 vapours contributing to the initial formation of critical clusters.




Fig 6: For all new particle formation (NPF) events the monthly ratio of the total number concentration of positively charged small intermediate ions (2-4nm) over the total concentration of both positively and negatively charged small intermediate ions (2-4nm) ($N_{2-4nm}^+ / (N_{2-4nm}^- + N_{2-4nm}^+)$). N_{2-4nm}^- and N_{2-4nm}^+ includes only ion concentrations within the identified growing mode between 2 and 4nm. The mean monthly $N_{2-4nm}^+ / \text{total}$ is presented as a cross, and the median is displayed by coloured circles, blue for an overall negative charge (ratio < 0.5) and red for an overall positive charge (ratio > 0.5). The grey shading displays the 25th – 75th percentiles. The dashed black line represents an overall neutral charge of small intermediate ions. It should be noted that there may be some instrumental biases between the two polarities, however, the background concentrations (during no-event periods) display $N_{2-4nm}^- / N_{2-4nm}^+$ ratios close to unity, ~1.02.

590

591 As mentioned above, Beck et al. (2021) reports higher concentrations of precursor gases with low proton affinity
 592 (e.g., MSA, SA, HSO_4^-) during the month of May, while highly oxygenated organic molecules (HOMs) display
 593 higher concentrations than SA and MSA during April and from June towards September. Beck et al. (2021)
 594 suggest that the events in May were likely initiated by negative ion-induced NPF and relate to sulphuric acid and
 595 ammonia nucleation.

596 Ammonia (NH_4^+) is a possible positively charged cluster ion which could play a role in the shift in polarity
 597 (Kirkby et al., 2016), however, it needs to be stressed that further work is needed to ascertain what is the reason
 598 for this shift in the polarity. We were unable to compare the overall polarity charges for small intermediate ions
 599 with concentrations of NH_4^+ . However, the shift from more negative to more positive small intermediate ions
 600 seems to coincide with the increase in concentrations of HOMs relative to the total concentration of other
 601 condensable vapours as also indicated above (see Fig. S24).

602 3.3 Source regions and geo-spatial extent

603 NPF events are assigned to various marine regions using the method described in Sect. 2.4.5. Air masses that
 604 traverse over the Arctic Ocean coincide with the highest number of NPF events (57 in total or 37% of NPF
 605 events; see Table 1). 37% of air masses are classified as originating from the Arctic Ocean, the largest marine
 606 region in terms of contribution. The air masses that mainly originate over the Greenland Sea result in 29 NPF

events and show the highest probability of an NPF event occurring, with 27% of the Greenland Sea air masses being linked to an NPF day. Air masses originating over the Arctic Ocean and Barents Sea have a 19% and 13% likelihood, respectively.

Table 2: The frequency of occurrence of air masses from different sectors compared with the number of events. Both numbers are given in days, followed by a ratio of the two (i.e. Likelihood). Although not shown in table, back trajectories traversed over the Kara Sea on two occasions. The average CS, solar insolation (both mixed-layer and all), maximum daily increase in 2.8 – 5nm particles ($\Delta N_{\max, 2.8-5}$) and proportion of back trajectory within the mixed layer are given for each marine sector. The averages for these variables are also given during the NPF days and presented in brackets.

Air mass	Region's frequency (days)	Number of NPF events (days)	Likelihood	CS [s^{-1}]	Solar insolation [$h W m^{-2}$]	$\Delta N_{\max, 2.8-5}$ [cm^{-3}]	Proportion in ML
Arctic Ocean	313	57	0.19	0.20 (0.19)	185 (378) 368 (753)	239 (776)	0.63 (0.60)
Greenland Sea	109	29	0.27	0.25 (0.25)	327 (442) 666 (991)	254 (665)	0.60 (0.54)
Barents Sea	69	9	0.13	0.36 (0.29)	118 (211) 597 (1100)	93 (204)	0.32 (0.28)
Norwegian Sea	44	7	0.16	0.25 (0.43)	232 (418) 618 (1330)	136 (382)	0.48 (0.34)
Mixed	303	51	0.17	0.23	184	147	0.54 (0.50)
Total	840	153	0.18	0.23	200	190	0.57 (0.53)

The analysis is expanded, using the method outlined in Sect. 2.4.4, by estimating the location of the formation of newly formed particles. The most common location for nucleation to occur is off the western coast of Svalbard, i.e., in the Greenland Sea and Arctic Ocean (see Fig. 7). Furthermore, the Greenland Sea, south-west of ZEP, as opposed to the Barents Sea, is home to more nucleation sites. It is also clear that the vast majority of nucleation sites are estimated to occur over the ocean. The estimated nucleation sites are almost all within a $1 \cdot 10^6 km$ radius of ZEP, and the vast majority are fairly close to ZEP (see Fig. 7), a finding which can also be inferred from the observation that the majority of events begin growing from 2.5nm. Furthermore, from the air mass back trajectory analysis, we gather that air masses typically reside between sea level and the altitude of ZEP (i.e. 0-474m, see Fig. S20). There is little difference between the altitudes of the arriving air masses for the NPF events compared to all days.




Fig 7: Total number of estimated nucleation sites per grid cell. The unit is based on the number of back trajectories initialised and does not reflect a physical value. Sites of nucleation for class Ia, Ib and II events are estimated by using HYSPLIT to trace the likely origin of particles produced during NPF events, and initialising the back trajectories based on the duration of the NPF event (see Sect. 2.4.4).

626

627 DMS emissions are closely linked to phytoplankton dynamics and abundance, and typically occur following
628 phytoplankton biomass maxima (Galí and Simão, 2015). Numerous studies have tried to link chlorophyll a, which
629 is used as a proxy for phytoplankton biomass, with DMS emissions to the atmosphere (Jang et al., 2019). Park
630 et al. (2018) note a strong correlation between DMS concentrations measured at ZEP and air masses traversing
631 over the surrounding ocean with high chlorophyll a. They find that the relationship between DMS concentrations
632 and air masses likely exposed to high DMS emissions are stronger for air masses traversing over the Greenland
633 Sea, compared with the Barents Sea. Therefore, suggesting that the Greenland Sea has a DMS production
634 capacity three times greater than the Barents Sea. By replicating the method detailed in Park et al. (2018), we
635 also find good correlations between chlorophyll a exposure and DMS, however, not for all years, and not for the
636 years related to this study (i.e. 2022 and 2023). In the study by Park et al. (2018) the years 2010, 2014 and 2015
637 were included, however, 2022 and 2023 did not yield such strong correlations. We were unable to replicate the
638 findings presented in Park et al. (2018) for the years 2022 and 2023 (see Fig. S30-37), however, we could
639 replicate the findings for 2010 (see Fig. S30). We can only speculate about the reasons for the interannual
640 variation between the relationship between chlorophyll a and DMS. We argue that because of the large spatial

641 and temporal interannual variability, possible geo-spatial changes in the DMSP-rich phytoplankton species in the
642 neighbouring seas, a wide range of DMS oxidation timescales depending on photochemistry and available solar
643 insolation (Ghahreman et al., 2019) can all potentially influence this relationship. Furthermore, the influence of
644 various other meteorological and environment factors that affect DMS oxidation and the lifetime of precursor
645 gases, and sea ice and cloud cover could lead to inaccuracies in HYSPLIT outputs and chlorophyll a satellite
646 retrievals. In addition, direct correlations between atmospheric DMS and chlorophyll a exposure are not detected
647 around Iceland, however, a high atmospheric DMS to chlorophyll a exposure coincides with areas containing
648 DMSP-rich phytoplankton species (Lee et al., 2023).

649 The Greenland Sea is connected to a higher likelihood of NPF events; compared with the Barents Sea there is a
650 distinct difference between both seas in terms of the proportion of air masses coinciding with NPF events, and
651 the number of nucleation sites estimated to arise from each sea (see Fig. 7). The combined effect of a larger
652 number of air masses coming from the west and an increased NPF-likelihood linked to western air masses means
653 that we observe more NPF events off the western edge of Svalbard. This difference could be linked to the
654 finding that the DMS production capacity in the Greenland Sea is estimated to be three times greater than that of
655 the Barents Sea, due to more DMSP-containing phytoplankton species (e.g. *Phaeocystis*) (Park et al., 2018).
656 However, another distinct difference between these two source regions is that air masses, which traversed over
657 the Greenland Sea, experienced more solar insolation and a relatively lower CS average, compared to the air
658 masses coming over the Barents Sea (see Table 1). The combination of both high solar insolation and relatively
659 low CS, could also explain why NPF likelihood is higher for the Greenland Sea as opposed to Barents Sea. One
660 further difference is that the Barents Sea has a much larger sea ice coverage compared with the Greenland Sea,
661 and thus there is potentially less open ocean in contact with the lower atmosphere during the NPF season. The
662 reduced availability of DMS given the greater sea ice coverage could additionally help explain the differences in
663 NPF likelihood of the Greenland and Barents Seas. The Arctic Ocean is another NPF source region (see Fig. 7
664 and Table 1.). Considering that the Arctic Ocean sector is considerably more covered by sea ice then the NPF
665 precursors are potentially being emitted from sea ice, e.g. iodine and sympagic algae. The Arctic Ocean
666 contributes significantly, in terms of the portion of arriving air masses, between March to May and September to
667 October (see Fig. S22). HIO_3 and HOMs are both significant contributors prior to May and HOMs increase in
668 concentration after June.

669 4.1 Predicting NPF likelihood with condensation sink and solar radiation

670 In Sect. 3.1.2, we show that NPF events and increases in NPF intensity, typically occur during periods of high
671 solar insolation. High solar insolation seems to be a prerequisite for NPF, as it leads to the photochemical
672 production of nucleating compounds. NPF events also seem to require the total surface area of pre-existing
673 aerosol particles (i.e., CS) to decrease below seasonal averages (see Fig. 2). The distribution of solar insolation
674 is statistically significantly different for periods experiencing events as opposed to non-events; or CS it depends
675 on the classification scheme used (as demonstrated using the Mann-Whitney U test, see Fig. S29 a) – b) in
676 supplement). NPF events require the concentration of nucleating vapours to exceed a critical concentration, and
677 as the CS acts as a sink of those highly condensable vapours, low CS values promote the potential for nucleation.
678 In this section, we argue that solar insolation and CS can be used as the two central parameters to determine with
679 high certainty the likelihood that a given day will experience an NPF event. For the definition of the term solar
680 insolation refer to Sect. 2.4.1 in the methods.

681 The majority of measurements feature the combination of both relatively low solar insolation and low CS (<1000
682 hW m^{-2} and $<0.5 \cdot 10^{-3} \text{ s}^{-1}$ see bottom-left of Fig. S25 b) in supplement)), and on very few occasions there are
683 measurements with a combination of high solar insolation ($>1000 \text{ hW m}^{-2}$) and low CS ($<0.5 \cdot 10^{-3}$) (see top-left
684 of Fig. S25 b)). Nonetheless, we show that when the combination of high solar insolation ($>1000 \text{ hW m}^{-2}$) and
685 relatively low CS ($<0.5 \cdot 10^{-3} \text{ s}^{-1}$ (i.e. top-left of Fig. S25 a) & b)) arises, an NPF event that day is likely to occur
686 (as defined by Aliaga et al. (2023), see Fig. 8). We find that periods exhibiting high solar insolation ($>1000 \text{ hW}$
687 m^{-2}) and relatively low CS ($<0.5 \cdot 10^{-3} \text{ s}^{-1}$) are linked to more NPF days. Overall, the combinations of CS and
688 solar insolation which promote NPF are relatively rare, compared to the entire set of measurements. The
689 probability of occurrence is derived from the ratio of the number of measurements for each combination of CS
690 and solar insolation during an NPF day to the number of valid data points for each of these combinations (where
691 both sets of data are composed of hourly measurements) (see Sect. S10.1 in the supplement for more details on
692 the normalisation approach)).




Fig. 8: Grid mesh where the data is binned by accumulated solar insolation and condensation sink (CS) (CS between $0 - 10^{-3} \text{ s}^{-1}$ every $2 \cdot 10^{-5} \text{ s}^{-1}$ and solar insolation between $0 - 4 \cdot 10^4 \text{ h W m}^{-2}$ every 200 h W m^{-2}). Each grid box displays the likelihood, calculated by normalising the data set for the event days (defined using Aliaga et al. (2023)) by the total count for each given bin (i.e. see Fig, S25 a) and b) in supplement)). Likelihood is the fraction of measurement combinations linked to an NPF day. The red line defines the region of interest (ROI) whereby data on the left of it is considered to have the ideal conditions for NPF i.e. high accumulated solar flux and relatively low CS. The equation for the red line is: $\text{Solar insolation} = 5 \cdot 10^6 \cdot \text{CS} + 500$. The grids required a count greater than 1.

693

694 We define a region of interest (ROI) (represented using the red line, separating the top left-hand corner in figure
 695 8). The ROI is defined using a simple straight line to separate ideal and non-ideal NPF conditions from each
 696 other. It should be stressed that it should not be used as an empirical relationship, defining the occurrence of
 697 NPF, instead it should be used as a guideline to highlight the general conditions most suitable for the promotion
 698 of NPF. In the ROI, the combination of low CS and plentiful solar insolation provide the most favourable
 699 conditions for NPF days to occur, and on the contrary, outside of the ROI, NPF days are much less likely to
 700 occur (i.e. bottom-right of Fig. 8).

701 Moreover, the region outside of the ROI with low solar insolation and low CS might be associated with
 702 increased cloudiness as the incident radiation is reduced by the presence of clouds, and large accumulation mode
 703 particles are reduced due to the effect of wet scavenging. The region with low solar insolation and high CS
 704 could be related to Arctic Haze conditions.

705 The ROI defined in Figure 8 is statistically significantly different from the non-ROI region on the probability
 706 that an NPF day occurs given such solar insolation and CS measurements occurring; the probabilities which arise
 707 when normalisation is done via the nanoranking classification scheme, as opposed to the Dal Maso et al. (2005)
 708 scheme, are more distinct between the ROI and non-ROI regions (see Fig. 9). Here, the statistical significance

709 between the two regions is conducted using the non-parametric Mann-Whitney U test (where H_0 representing
 710 two identical distributions is rejected). In addition, the ROI and non-ROI represented 23% and 77% of the data
 711 respectively; for the ROI 62.5% of the hourly data were linked to NPF days, as opposed to 23.8% in the non-ROI
 712 (The total number of Nano_{events} is 229 (g2=178, g3=51, see Fig.S16 in supplement)).




Fig 9: Violin plots representing the distribution of the probabilities as defined by the normalisation plot (see Fig. 8). The red violins on the left-hand-side represent the probability distributions related to the Dal Maso et al. (2005) classification and the green violin on the right-hand side represent the probability distributions according to the Nano-ranking approach.

714 The data points that lie within the ROI and when they occur can be used to try and estimate the likelihood of a
 715 day experiencing an NPF event. A time series representing the daily likelihood of NPF occurrence (i.e. $P_{NPF, D}$)
 716 can be generated from the ROI data (see Fig. 10) by labelling hourly data points as either 1 (i.e. within the ROI)
 717 or 0 (i.e. outside of the ROI) and then taking daily averages before applying a rolling average to create the daily
 718 likelihood, $P_{NPF, D}$. $P_{NPF, D}$ generally, matches the frequency of NPF events as defined by Aliaga et al. (2023) or
 719 Dal Maso et al. (2005) (see Fig. 10), however, there are some major discrepancies. Generally, the predicted
 720 frequency of occurrence (i.e., defined by the ROI) is significantly lower than the observations (can also be
 721 observed in the correlation figure Fig. S28 in the supplement). The spring and autumn peaks in both $P_{NPF, D}$ and
 722 the event frequencies generally line up well in terms of the number of peaks, however, 2023 displays only one
 723 main peak in $P_{NPF, D}$. $P_{NPF, D}$ nicely replicates the distinctive dips in the NPF frequency before July in the years
 724 2022 and 2024. However, in July 2023, the dip in NPF events in the observations is followed by an increase (i.e.
 725 two more sequential peaks in the observations) which is not captured by $P_{NPF, D}$. Still, it is striking that the two

parameters alone (CS and solar insolation) are able to serve as predictors for NPF probability, by representing the balance between the production and removal of NPF-precursors. The solar insolation reflects the photochemical production potential (likely via the production of hydroxyl radicals). However, the parameter combination does not account for the reactants required to produce the nucleating vapours (which may be SO₂ (likely via DMS) and/or iodine compounds). Hence, deviations like those observed during July 2023 are expected as the conceptual model is partly incomplete due to a lack of adequate information regarding the original reactants. This said, however, the NPF frequency and likelihood largely seem limited by the balance between photochemical production potential and CS, and this in turn is suggestive of, to some degree, a self-regulating mechanism regarding the number of particles present in the nucleation and Aitken modes. There exist more sophisticated models to predict NPF (Kuang et al., 2010).




Fig 10: The red line is $P_{NPF, D}$, the estimated daily likelihood of measurements occurring in the region of interest (ROI). The ROI is based on the relation between CS and 6-hour solar accumulation and is defined in Fig. 8). The green and blue lines represent the rolling averages of the NPF daily event frequencies using the Dal Maso et al. (2005) and Aliaga et al. (2023) classification respectively. The green and blue curves are present here to be able to compare to $P_{NPF, D}$ (daily prob. ROI).

736

737 4.1.1 Ratio of source and sink: ratio of solar irradiance

738 In section 4.1 we defined a region of interest, whereby measurements within this ROI provide an indication of
 739 the conditions likely to lead to an NPF day. Observing when these conditions occur offers one approach,
 740 however, the ability to predict the occurrence of NPF days can be improved by simply using the proxies for a
 741 source and a sink, and taking their respective ratio (i.e., solar insolation/CS) (see Fig. 11). For a rolling average
 742 window of 30 days the R^2 , comparing solar insolation/CS and the nano-ranking frequency, is 0.78 (see Fig. S26
 743 in supplement).




Fig 11: The solar insolation divided by the condensation sink (Solar insolation/CS, SUN FLUX/CS) with a 30-day rolling mean, is denoted by the red line. The intensity parameter ($\Delta N_{2.82-5nm}$) is shown in black. The green and blue lines represent the rolling averages of the NPF daily event frequencies using the Dal Maso et al. (2005) and Aliaga et al. (2023) classification schemes respectively. The green and blue curves are displayed to be able to compare to the red line, the proxy for NPF activity i.e. the present to be able to compare to the red line representing a proxy for the balance between sources and sinks.

744

745 Figure 11 demonstrates that by using a simplified parameter i.e., solar insolation/CS, albeit with a large rolling
 746 averaging window, we achieve a good agreement with both event frequencies (i.e., classified using the Dal Maso
 747 et al. (2005) or Aliaga et al. (2023) approach). There are some slight discrepancies between the solar
 748 insolation/CS parameter and the event frequencies, particularly at the start and end of the measurement period;
 749 the discrepancies could be due to the NPF seasons having started already or not having ended. The relationship
 750 between Solar insolation/CS and event frequency varies depending on the averaging window utilised, whether
 751 the ML is selected and also the type of classification used (see Fig. 12). The coefficient of determination (R^2)
 752 comparing solar insolation/CS and the daily event frequency (i.e., rolling average applied) increases and begins
 753 to plateau after a window of 2 weeks is applied (14 days, $R^2=0.65$) (see Fig. 12). Considering that a degree of
 754 averaging needs to be performed as the daily event frequencies are given as 1s (event) and 0s (non-event), then it
 755 seems justified that the solar insolation/CS parameter offers a robust predictive capability once a modest amount
 756 of smoothing is performed.




Fig. 12: Correlation between the event frequency (either defined using the Dal Maso or Nano ranking method) and the accumulated solar insolation (SUN parameter) with a rolling average applied. The coefficient of determination (R^2) is shown as a function of the rolling window size (number of days). The first 6, 12, and 24 hours of the back trajectory are selected (shown in slightly different shades). The curve labelled ML represents back trajectories that traversed only through the mixed layer, while the curves labelled all alts. represent back trajectories at all altitudes.

757

758 4.2 Dynamics driving NPF and growth

759 During NPF events there is an excess availability of condensable precursors, e.g. SA, which contribute to the
760 growth of newly formed and pre-existing particles, thus increasing CS. The increased CS enhances the removal
761 of condensable and nucleating vapours. As a result, a typical day with sustained NPF can lead to a relatively
762 high CS ($>0.5 \cdot 10^{-3} \text{ s}^{-1}$), although the events tend to begin at low CS values ($\sim 0.1 \cdot 10^{-3} \text{ s}^{-1}$) (see Fig. 13). It is
763 worth noting, that during an event, not only do the nucleation mode particles grow in size, but also all other
764 existing particles grow as well. Practically, this means that the sink is doubled during the NPF-event.
765 Consequently, and assuming steady state, this would mean that the source rate of the condensing species must
766 double to sustain the same concentration of condensing material.




Fig. 13: The calculated condensation sink (CS) during NPF events. It should be noted that the average NPF typically lasts for 10 hours; however, in some cases events can persist for days until the newly formed mode is removed via wet scavenging or advected away from ZEP. The black line is the mean and the blue is the median. The blue shaded region represents 25th-75th percentiles. The dashed blue lines display CS values of 0.1, 0.2 and 0.4 • 10⁻³ s⁻¹ and are there to guide the reader.

767

768 The effect of this additional increase in CS during NPF events can be illustrated by using the ROI. The increase
 769 in CS shifts the ambient conditions away from the defined ROI ($>0.5 \cdot 10^{-3} \text{ s}^{-1}$) (see Fig. 8), thus potentially
 770 suppressing any subsequent NPF events. This mechanism will prevent further NPF, if the increase in CS during
 771 the NPF event is large enough to limit the availability of precursors. This mechanism also helps to explain why
 772 the combination of low CS and high solar insolation measurements are seemingly rare occurrences, as these
 773 conditions are soon followed by NPF, and increasing CS.

774 During the Arctic Haze period in late winter and early spring, high concentrations of accumulation mode aerosol
 775 dominate the contribution to the CS. Reduction in accumulation mode aerosol at the end of the Arctic Haze
 776 season (i.e., end of May) should result in decreased CS, but ZEP continues to experience relatively high CS. We
 777 argue that this relatively high sustained CS is the result of particle formation and growth within the Arctic; the
 778 increasing frequency of NPF events during the summer months increases CS, and maintains the comparatively
 779 high CS throughout the summer and the most intense part of the NPF season (see Fig. 2b)), additionally Tunved
 780 et al. (2013)). Towards the late summer and autumn, the solar insolation has decreased substantially, and both
 781 marine surface water productivity and photochemical production potential are declining. Together with low
 782 transport efficiency from anthropogenic sources at lower latitudes and efficient wet removal, the aerosol reaches
 783 a minimum with respect to CS and number during this part of the year.




Fig 14: Air mass history for a) percentage of back trajectory endpoints within an ensemble with a relative humidity above 85, and b) percentage of endpoints within ensemble which experienced a rainfall event (i.e. rainfall > 0mm). Y-axis is the number of hours backwards before the air masses arrive at ZEP, while the x-axis is the number of hours into the event e.g. if one event started at 6am and another at 9am, their start times would both be 0 hours. The black line is a 1:1 line which represents the start of the NPF event such that the grid points towards the top-left represent endpoints prior to the NPF event, whilst grid points towards the bottom-right represent endpoints during the NPF event. The x-axis has been sliced based on the typical duration of events (most events are <14 hours). This figure consists of the air mass history of Class Ia and Ib NPF events.

784

785 In the Arctic environment, wet removal is a key factor responsible for the rapid reduction in CS (Garrett et al.,
 786 2011). For class Ia and Ib NPF events, we can observe that prior to the start of an NPF event there is an
 787 increased likelihood of cloudy conditions ($RH > 85\%$) and rainfall events (see Figs 14 a-b). Figures 14 a-b
 788 demonstrate how wet scavenging helps to effectively reduce the CS and provide more suitable conditions for
 789 NPF events to occur. The changes in the likelihood of cloudy conditions and rainfall prior to and post the start of
 790 an NPF event is slight, but significantly different, especially considering the duration prior to the event is longer
 791 and the likelihood of any rainfall should be compounded.

792 Any reduction in CS will shift the system towards the ROI, and as a result the system will rapidly respond to this
 793 reduction by producing new particles and growing the already existing ones. If suitable gaseous precursors and
 794 solar insolation are present, NPF will take place and the cycle will be reinitialised. This conceptual model, whilst
 795 highly simplified, offers substantial predictive value regarding the timing and frequency of NPF events (see Fig.
 796 10). It does not, however, account for the original source of condensable and nucleating species nor does it
 797 consider cloud activation and the in-cloud chemistry of Aitken mode particles in non-precipitating clouds.

798 4.3 NPF contribution to Aitken and potential CCN

799 Here, we argue that new particle formation is the main contributor to the Aitken mode in the Arctic; through
800 frequent NPF events and the subsequent growth of nucleation mode particles, NPF activity is able to provide a
801 significant proportion of the overall concentration of Aitken mode particles. We show in Sect 3.1.2 that at ZEP
802 the production of nucleation mode particles is largest at the end of spring, however, the production of newly
803 formed particles persists throughout the summer months and into the autumn (see Fig. 2a)). During NPF events,
804 including formation and subsequent growth the total number concentration of Aitken mode particles (5-70nm) is
805 3.6 times the amount compared with pre-event conditions (see Fig. 16). These events are relatively frequent (30
806 - 40% of summer days experience NPF, (see Fig. 1). Fundamentally, though, it is difficult to use in-situ
807 measurements at a single location to provide an estimation of the fraction of Aitken mode particles which are
808 produced via NPF; for the Aitken mode particles at ZEP which are not directly linked to NPF activity, we cannot
809 rule out the possibility that they are derived from NPF events outside of the vicinity of ZEP and then transported
810 downwind.

811 At ZEP, NPF and the subsequent growth that follows, results in particles large enough to potentially be activated
812 and become CCN; on numerous occasions (57 times) we observed newly formed particles able to grow beyond
813 25nm while in the same regional event (see Fig. 15b), before being scavenged. Here, we use 25nm as the lowest
814 bound for particles to act as CCN, however, it should be stated that this limit requires strong updrafts and
815 extremely hygroscopic particles for particles to be activated. We observe that after ~30 hours of growth, we
816 reach potential CCN-sizes at ZEP (see Fig. 15a)). In addition, on 11, 8 and 3 respective occasions
817 NPF-produced particles grew beyond 60, 70 and 80nm were observed (see Fig. 15b)). It is important to note that
818 once produced these particles most likely continue to grow downwind of ZEP by condensation and in-cloud
819 processing, so a large fraction of the Aitken mode should inevitably reach CCN size in due time, as long as they
820 are not otherwise scavenged.




Fig 15: Change in diameter during all class Ia, Ib and II NPF events. The blue shaded region covers 25nm – 60nm. (a) shows the average change in the diameter of the nucleation mode over time. The black and blue lines signify the mean and median diameters respectively. The blue shaded region displays the distance between the 25th and 75th percentiles respectively (b) displays the number of NPF events against the maximum diameter reached during growth. The horizontal dashed line marks 25nm whilst the vertical dashed line marks the number of NPF events which reached 25nm.

821

822 It is difficult to ascertain the proportion of NPF-derived Aitken mode particles that actually become CCN before
823 being scavenged or deposited, and estimating an exact fraction is beyond the scope of this study. However,
824 previous modelling and measurement studies can help to shed some light on the likelihood that NPF events as
825 measured from ZEP can contribute to the CCN budget. Jung et al. (2018) demonstrate that at supersaturations
826 (ss) greater than 0.4% there is a considerable number of CCN measured during the summer months at ZEP.
827 Moreover, the changes in sources of CCN-sized aerosol particles, from mainly anthropogenic to NPF-derived,
828 during the Haze to summer transition can be seen in the reduction of the geometric mean diameter. For summer,
829 Jung et al. (2018) suggest a significant number of particles >10nm are activated into cloud droplets, as the
830 activation ratio during summer is around 0.5 for fairly high ss, i.e. >0.6%. In addition, Karlsson et al. (2021)
831 show that Aitken mode-sized cloud residuals are fairly common in the summer time (i.e. July onwards); the
832 transition from accumulation mode-dominated to Aitken mode-dominated aerosol did not lead to a significant
833 reduction in the concentration of cloud residuals, indicating that particles as small as 20nm can be activated.
834 Motos et al. (2023) suggest that particles >20nm are activated in the summer, with a modal activation diameter
835 of 60nm showing that Aitken mode particles act as CCN.




Fig 16: The particle number size distributions (DMPS) for periods before the NPF events (1-2 hours before event started) (pre-event, light blue), during NPF and growth (black) and on-site nucleation (production of 2-4nm particles, red, *on-site formation*).

836

837 5. Conclusion

838 In this study, we explore the annual cycle of atmospheric new particle formation (NPF) events from almost
839 3-years' worth of measurements (2022-2024). This study is the first to utilise the Nano-ranking method in the
840 Arctic. It should be stated that special care needs to be taken in treating wind-blown events separately from NPF
841 events. Furthermore, this study also highlights potential problems when sampling lines are submerged in dense
842 clouds. It is recommended that relative humidity, visibility and/or web cam footage is used to help screen for
843 these in-cloud events when sampling at high altitudes. In keeping with previous studies, e.g., Tunved et al.
844 (2013) and Ström et al. (2009), the Arctic summer and more specifically the end of spring and start of summer
845 represent a peak in NPF intensity as observed at Zeppelin Observatory (ZEP). We find that this peak in the
846 strength of NPF events coincides roughly with the maximum solar insolation (defined as the last 6 hours of
847 accumulated solar radiation along HYSPLIT back trajectories); solar insolation allows for the increased
848 availability of precursors by encouraging oxidation. The NPF frequency during the summer remains fairly
849 consistent, with an estimated 20-30% of days experiencing an NPF event. Towards the end of summer, the
850 decline in the amount of solar insolation reduces the NPF intensity. In addition, it is also apparent that the
851 condensation sink (CS i.e. linked to the total surface area of pre-existing aerosol) plays an important role, as the
852 NPF intensity is more pronounced when there are reductions in CS. For NPF events, the minimum CS was

853 observed at the start of the events and signifies that a reduced condensation sink is required for events to begin.
854 ~~This study is the first to utilise the Nano-ranking method in the Arctic. It should be stated that special care needs~~
855 ~~to be taken in treating wind-blown events separately from NPF events. Furthermore, this study also highlights~~
856 ~~potential problems when sampling lines are submerged in dense clouds. It is recommended that relative~~
857 ~~humidity, visibility and/or web cam footage is used to help screen for these in-cloud events when sampling at~~
858 ~~high altitudes.~~

859 Typically, as the Arctic transitions from the Haze season to summer, cloud coverage and precipitation increase,
860 which in turn, via wet scavenging, reduces the once predominant accumulation mode. However, the increased
861 cloud cover, not only reduces the CS, but also reduces the solar insolation. Hence, the transition to summertime
862 both promotes and discourages NPF. The latter stages of the summer and the autumn provide less solar
863 insolation; the peak in solar insolation is skewed more towards the start of summer. These competing effects can
864 explain why we observe such a reduction in NPF intensity as the summer progresses. The dip in frequency and
865 intensity that occurs around July coincides with a decrease in brightness (solar insolation over potential solar
866 radiation). It is the interplay between both solar insolation and condensation sink that can potentially give rise to
867 NPF events, and thus cloudiness will also play a role in NPF activity given interplay with CS and solar radiation.
868 We can only speculate that a warming climate may appear more similar to the late Arctic summer, being both
869 warmer and wetter. Some studies indicate that cloud cover in the Arctic could potentially increase (Barton and
870 Veron, 2012) or has (Eastman and Warren, 2010; Francis et al., 2009; Kay and Gettelman, 2009; Palm et al.,
871 2010; Vihma et al., 2008) with periods of rapid sea ice loss.

872 One main finding from this study is that we can, with rather good accuracy, predict the occurrence of an NPF
873 day (Fig. S26 in the supplement) by using CS and solar insolation as the two main predictive parameters.
874 Periods exhibiting low CS and high solar insolation provide the most ideal conditions for NPF events and the
875 occurrence of these periods can reproduce the observed frequency of NPF events. It should be noted, though,
876 that other factors play a role. The sources to the precursors of the condensable vapours, as well as additional
877 meteorological and environmental parameters such as sea-ice cover, chlorophyll a concentration and the varying
878 contributions from different source regions all can influence the occurrence and strength of NPF. However, from
879 this study we argue that both solar insolation and CS can be used to create a rather simple predictive model,
880 simply by taking the ratio of these two terms. The Arctic perhaps offers a unique environment, where the
881 relationship between solar insolation and NPF is potentially stronger, due to the typically lower CS and lower
882 overall solar radiation intensity (Bousiotis et al., 2021).

Quantifying the contribution of CCN from NPF on low-level clouds is beyond the scope of this study. However, we are able to demonstrate that periods of formation and subsequent growth lead to a significant increase in the proportion of the total number of Aitken mode particles. In combination with this, we show that on many occasions during NPF events, the nucleation mode particles grow to potential CCN sizes once they are formed (57 NPF events lead to diameters $>25\text{nm}$). With the correct conditions, i.e. a relatively strong updraft, the formation of particles $>25\text{nm}$ could lead to droplet formation and in turn alter cloud properties.

Given the high frequency of onsite NPF and the large potential area for formation, it is not unlikely that a majority of Aitken mode particles originate from NPF within or close to the Arctic region. Provided sufficient time for growth up to CCN-sizes, it seems clear that NPF events could constitute a significant source of CCN in the Arctic. For a more detailed picture of the aerosol budget in the Arctic and a more quantitative account of the NPF-contribution to the overall Arctic CCN-population, large scale modelling evaluated against observations and combined with process studies are required.

Code availability

Code will be made available on GitHub when published.

Data availability

Data will be on the Bolin Centre database when published (<https://bolin.su.se/data/>).

Author Contributions

Dominic Heslin-Rees (DHR), Radovan Krejci (RK), and Peter Tunved (PT) worked together to develop the research questions and design the study. DHR performed data analysis and wrote the paper together with assistance from RK and PT. There was good input and advice along the way from Ilona Riipinen (IR), Annica Ekman (AK), Diego Aliaga (DA) and Martina Mazzini (MMi). Janne Lampilahti (JL) developed many of the NAIS specific Python packages and showed DHR how to clean and service the NAIS. Stefania Gilardoni (SG) provided comments. SG, Mikko Sipilä (MS), Mauro Mazzola (MMA) and Roseline Thakur (RT) helped with the Gruvabadet work. Ki-Tae Park (KTP) and Kitack Lee (KT) maintained and operated the Zeppelin DMS analyzer as part of a collaborative effort between KOPRI and POSTECH research team. Kihong Park (KP) and Young Jun Yoon (YJY) provided the Nano-SMPS data.

910 Acknowledgements

911 This study would not be possible without the work done by the research engineers Birgitta Noone, Tabea
912 Henning, and Ondrej Tesar from ACES and the staff from the Norwegian Polar Institute (NPI). NPI provides
913 invaluable on-site support including attending to the NAIS. NPI is also acknowledged for substantial long-term
914 support in maintaining the measurements at Zeppelin Observatory (ZEP). We would also like to thank Kai
915 Rosman for developing the software we used for the instrumentation at ZEP. We would like to thank Ove
916 Hermansen from the Norwegian Institute for Air Research (NILU) for providing the ambient meteorological
917 data.

918 Financial Support

919 Aerosol size distribution measurements are supported by the Swedish EPA. Swedish participation was supported
920 by the Swedish Environmental Protection Agency (Naturvårdsverket), Knut and Alice Wallenberg foundation
921 (KWA) and FORMAS funding agency.

922 Knut-and-Alice-Wallenberg Foundation within the Arctic Climate Across Scales (Project No. \,2016.0024), the
923 Swedish EPA's (Naturvårdsverket) Environmental monitoring program (Miljöövervakning), the Swedish
924 Research Council FORMAS (Project "Interplay between water, clouds and Aerosols in the Arctic," \#
925 2016-01427)

926 The aerosol observations at the Zeppelin station have been supported by the KAW Stiftelse (grant
927 no. 2016.0024), the Swedish Environmental Protection agency (Naturvårdsverket), and the ACTRIS Sweden
928 project supported by the Swedish Research Council

929 The KAW project CLIVE and VR financed infrastructure project ACTRIS-Sweden.

930

931

932

933 References

- 934 Aliaga, D., Tuovinen, S., Zhang, T., Lampilahti, J., Li, X., Ahonen, L., Kokkonen, T., Nieminen, T., Hakala, S.,
935 Paasonen, P., Bianchi, F., Worsnop, D., Kerminen, V.-M., and Kulmala, M.: Nanoparticle ranking analysis:
936 determining new particle formation (NPF) event occurrence and intensity based on the concentration spectrum of
937 formed (sub-5 nm) particles, *Aerosol Research*, 1, 81–92, <https://doi.org/10.5194/ar-1-81-2023>, 2023.
- 938 Asmi, E., Kondratyev, V., Brus, D., Laurila, T., Lihavainen, H., Backman, J., Vakkari, V., Aurela, M., Hatakka,
939 J., Viisanen, Y., Uttal, T., Ivakhov, V., and Makshtas, A.: Aerosol size distribution seasonal characteristics
940 measured in Tiksi, Russian Arctic, *Atmos Chem Phys*, 16, 1271–1287,
941 <https://doi.org/10.5194/acp-16-1271-2016>, 2016.
- 942 Barton, N. and Veron, D.: Response of clouds and surface energy fluxes to changes in sea-ice cover over the
943 Laptev Sea (Arctic Ocean), *Clim Res*, 54, 69–84, <https://doi.org/10.3354/cr01101>, 2012.
- 944 Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä,
945 O., Wimmer, D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, X. C., Kontkanen, J., Wolf,
946 K. K. E., Cappelletti, D., Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling,
947 A., Nøjgaard, J. K., Krejci, R., Karlsson, L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur, R.
948 C., Leino, K., Kangasluoma, J., Duplissy, E. M., Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A., Petäjä,
949 T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V. M., and Sipilä, M.: Differing Mechanisms of
950 New Particle Formation at Two Arctic Sites, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2020GL091334>,
951 2021a.
- 952 Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä,
953 O., Wimmer, D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, X. C., Kontkanen, J., Wolf,
954 K. K. E., Cappelletti, D., Mazzola, M., Traversi, R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling,
955 A., Nøjgaard, J. K., Krejci, R., Karlsson, L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur, R.
956 C., Leino, K., Kangasluoma, J., Duplissy, E. M., Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez, A., Petäjä,
957 T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V. M., and Sipilä, M.: Differing Mechanisms of
958 New Particle Formation at Two Arctic Sites, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2020GL091334>,
959 2021b.
- 960 Bousiotis, D., Brean, J., Pope, F. D., Dall'Osto, M., Querol, X., Alastuey, A., Perez, N., Petäjä, T., Massling, A.,
961 Klenø Nøjgaard, J., Nordstrøm, C., Kouvarakis, G., Vratolis, S., Eleftheriadis, K., Niemi, J. V., Portin, H.,

962 Wiedensohler, A., Weinhold, K., Merkel, M., Tuch, T., and Harrison, R. M.: The effect of meteorological
 963 conditions and atmospheric composition in the occurrence and development of new particle formation (NPF)
 964 events in Europe, *Atmos Chem Phys*, 21, 3345–3370, <https://doi.org/10.5194/acp-21-3345-2021>, 2021.

965 Boyer, M., Aliaga, D., Quéléver, L. L. J., Bucci, S., Angot, H., Dada, L., Heutte, B., Beck, L., Duetsch, M.,
 966 Stohl, A., Beck, I., Laurila, T., Sarnela, N., Thakur, R. C., Miljevic, B., Kulmala, M., Petäjä, T., Sipilä, M.,
 967 Schmale, J., and Jokinen, T.: The annual cycle and sources of relevant aerosol precursor vapors in the central
 968 Arctic during the MOSAiC expedition, *Atmos Chem Phys*, 24, 12595–12621,
 969 <https://doi.org/10.5194/acp-24-12595-2024>, 2024.

970 Brean, J., Beddows, D. C. S., Harrison, R. M., Song, C., Tunved, P., Ström, J., Krejci, R., Freud, E., Massling,
 971 A., Skov, H., Asmi, E., Lupi, A., and Dall’osto, M.: Collective geographical ecoregions and precursor sources
 972 driving Arctic new particle formation, *Atmos Chem Phys*, 23, 2183–2198,
 973 <https://doi.org/10.5194/acp-23-2183-2023>, 2023.

974 Cai, R. and Jiang, J.: A new balance formula to estimate new particle formation rate: Reevaluating the effect of
 975 coagulation scavenging, *Atmos Chem Phys*, 17, 12659–12675, <https://doi.org/10.5194/acp-17-12659-2017>,
 976 2017.

977 Covert, D. S., Wiedensohler, A., Aalto, P., Heintzenberg, J., Mcmurry, P. H., and Leck, C.: Aerosol number size
 978 distributions from 3 to 500 nm diameter in the arctic marine boundary layer during summer and autumn, *Tellus*
 979 *B Chem Phys Meteorol*, 48, 197–212, <https://doi.org/10.3402/tellusb.v48i2.15886>, 1996.

980 Croft, B., Martin, R. V., Richard Leitch, W., Tunved, P., Breider, T. J., D’Andrea, S. D., and Pierce, J. R.:
 981 Processes controlling the annual cycle of Arctic aerosol number and size distributions, *Atmos Chem Phys*, 16,
 982 3665–3682, <https://doi.org/10.5194/acp-16-3665-2016>, 2016.

983 Dada, L., Chellapermal, R., Buenrostro Mazon, S., Paasonen, P., Lampilahti, J., E Manninen, H., Junninen, H.,
 984 Petäjä, T., Kerminen, V. M., and Kulmala, M.: Refined classification and characterization of atmospheric
 985 new-particle formation events using air ions, *Atmos Chem Phys*, 18, 17883–17893,
 986 <https://doi.org/10.5194/acp-18-17883-2018>, 2018.

987 Dall’osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Ström, J., Hansson, H. C., Yoon, Y. J., Park, K. T.,
 988 Becagli, S., Udisti, R., Onasch, T., Ódowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to
 989 atmospheric new particle formation, *Sci Rep*, 7, <https://doi.org/10.1038/s41598-017-03328-1>, 2017.

990 Dal Maso, M., Kulmala, M., Lehtinen, K. E. J., Mäkelä, J. M., Aalto, P., and O'Dowd, C. D.: Condensation and
 991 coagulation sinks and formation of nucleation mode particles in coastal and boreal forest boundary layers,
 992 *Journal of Geophysical Research Atmospheres*, 107, <https://doi.org/10.1029/2001JD001053>, 2002.

993 Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P., Lehtinen, K. E. J., Maso, D.,
 994 Kulmala, M., Riipinen, M., Wagner, I., Hussein, R., Aalto, T., and Lehtinen, P. P. &: Formation and growth of
 995 fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II, Hyytiälä, Finland,
 996 2005.

997 Draxler, R. R., Spring, S., Maryland, U. S. A., and Hess, G. D.: An Overview of the HYSPLIT_4 Modelling
 998 System for Trajectories, Dispersion, and Deposition, *Australian Meteorological Magazine*, 295–308 pp., 1998.

999 Eastman, R. and Warren, S. G.: Interannual variations of arctic cloud types in relation to sea ice, *J Clim*, 23,
 1000 4216–4232, <https://doi.org/10.1175/2010JCLI3492.1>, 2010.

1001 Fiedler, V., Dal Maso, M., Boy, M., Aufmhoff, H., Hoffmann, J., Schuck, T., Birmili, W., Hanke, M., Uecker, J.,
 1002 Arnold, F., and Kulmala, M.: The contribution of sulphuric acid to atmospheric particle formation and growth: a
 1003 comparison between boundary layers in Northern and Central Europe, *Atmos. Chem. Phys.*, 5, 1773–1785,
 1004 <https://doi.org/10.5194/acp-5-1773-2005>, 2005.

1005 Finkenzeller, H., Iyer, S., He, X.C. et al. The gas-phase formation mechanism of iodic acid as an atmospheric
 1006 aerosol source. *Nat. Chem.* 15, 129–135 (2023). <https://doi.org/10.1038/s41557-022-01067-z>

1007 Francis, J. A., Chan, W., Leathers, D. J., Miller, J. R., and Veron, D. E.: Winter Northern Hemisphere weather
 1008 patterns remember summer Arctic sea-ice extent, *Geophys Res Lett*, 36, <https://doi.org/10.1029/2009GL037274>,
 1009 2009.

1010 Fuchs, N. A. and Sutugin, A. G.: *Highly Dispersed Aerosols*, Ann Arbor Science Publishers, Michigan, 1970.

1011 Galí, M. and Simão, R.: A meta-analysis of oceanic DMS and DMSP cycling processes: Disentangling the
 1012 summer paradox, *Global Biogeochem Cycles*, 29, 496–515, <https://doi.org/10.1002/2014GB004940>, 2015.

1013 Garrett, T. J. and Zhao, C.: Increased Arctic cloud longwave emissivity associated with pollution from
 1014 mid-latitudes, *Nature*, 440, 787–789, <https://doi.org/10.1038/nature04636>, 2006.

1015 Garrett, T. J., Zhao, C., Dong, X., Mace, G. G., and Hobbs, P. V.: Effects of varying aerosol regimes on low-level
 1016 Arctic stratus, *Geophys Res Lett*, 31, <https://doi.org/10.1029/2004GL019928>, 2004.

1017 Garrett, T. J., Brattström, S., Sharma, S., Worthy, D. E. J., and Novelli, P.: The role of scavenging in the seasonal
 1018 transport of black carbon and sulfate to the Arctic, *Geophys Res Lett*, 38,
 1019 <https://doi.org/10.1029/2011GL048221>, 2011.

1020 Ghahreman, R., Gong, W., Galí, M., Norman, A.-L., Beagley, S. R., Ayodeji, A., Zheng, Q., Lupu, A., Lizotte,
 1021 M., Levasseur, M., and Leaitch, W. R.: Dimethyl sulfide and its role in aerosol formation and growth in the
 1022 Arctic summer – a modelling study, *Atmos Chem Phys*, 19, 14455–14476,
 1023 <https://doi.org/10.5194/acp-19-14455-2019>, 2019.

1024 Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J.,
 1025 Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle,
 1026 C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozkhov, Y.,
 1027 Tröstl, J., Tsagkogeorgas, G., Wagner, R., Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K.
 1028 S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *Journal of*
 1029 *Geophysical Research: Atmospheres*, 122, 8739–8760, <https://doi.org/10.1002/2017JD026844>, 2017.

1030 Gormley, P. G. and Kennedy, M.: Diffusion from a Stream Flowing through a Cylindrical Tube, Source:
 1031 *Proceedings of the Royal Irish Academy. Section A: Mathematical and Physical Sciences*, 163–169 pp., 1948.

1032 Gramlich, Y., Siegel, K., Haslett, S. L., Freitas, G., Krejci, R., Zieger, P., and Mohr, C.: Revealing the chemical
 1033 characteristics of Arctic low-level cloud residuals - in situ observations from a mountain site, *Atmos Chem Phys*,
 1034 23, 6813–6834, <https://doi.org/10.5194/acp-23-6813-2023>, 2023.

1035 Heintzenberg, J. and Leck, C.: The summer aerosol in the central Arctic 1991-2008: Did it change or not?,
 1036 *Atmos Chem Phys*, 12, 3969–3983, <https://doi.org/10.5194/acp-12-3969-2012>, 2012.

1037 Hirsikko, A., Nieminen, T., Gagné, S., Lehtipalo, K., Manninen, H. E., Ehn, M., Hörrak, U., Kerminen, V. M.,
 1038 Laakso, L., McMurry, P. H., Mirme, A., Mirme, S., Petäjä, T., Tammet, H., Vakkari, V., Vana, M., and Kulmala,
 1039 M.: Atmospheric ions and nucleation: A review of observations, <https://doi.org/10.5194/acp-11-767-2011>, 2011.

1040 Hoffmann, E. H., Tilgner, A., Schrödner, R., Bräuer, P., Wolke, R., and Herrmann, H.: An advanced modeling
 1041 study on the impacts and atmospheric implications of multiphase dimethyl sulfide chemistry, *P. Natl. Acad. Sci.*
 1042 *USA*, 113, 11776–11781, <https://doi.org/10.1073/pnas.1606320113>, 2016. Jang, E., Park, K. T., Jun Yoon, Y.,
 1043 Kim, T. W., Hong, S. B., Becagli, S., Traversi, R., Kim, J., and Gim, Y.: New particle formation events observed

1044 at the King Sejong Station, Antarctic Peninsula - Part 2: Link with the oceanic biological activities, *Atmos Chem*
 1045 *Phys*, 19, 7595–7608, <https://doi.org/10.5194/acp-19-7595-2019>, 2019.

1046 Jang, S., Park, K. T., Lee, K., and Suh, Y. S.: An analytical system enabling consistent and long-term
 1047 measurement of atmospheric dimethyl sulfide, *Atmos Environ*, 134, 217–223,
 1048 <https://doi.org/10.1016/j.atmosenv.2016.03.041>, 2016.

1049 Jokinen, T., Sipilä, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E.-M., Junninen, H., Kangasluoma, J.,
 1050 Manninen, H. E., Petäjä, T., Kulmala, M., Worsnop, D. R., Kirkby, J., Virkkula, A., and Kerminen, V.-M.:
 1051 Ion-induced sulfuric acid-ammonia nucleation drives particle formation in coastal Antarctica, *Sci. Adv*, 2018.

1052 Jung, C. H., Yoon, Y. J., Kang, H. J., Gim, Y., Lee, B. Y., Ström, J., Krejci, R., and Tunved, P.: The seasonal
 1053 characteristics of cloud condensation nuclei (CCN) in the arctic lower troposphere, *Tellus B Chem Phys*
 1054 *Meteorol*, 70, 1–13, <https://doi.org/10.1080/16000889.2018.1513291>, 2018.

1055 Kangasluoma, J., Cai, R., Jiang, J., Deng, C., Stolzenburg, D., Ahonen, L. R., Chan, T., Fu, Y., Kim, C., Laurila,
 1056 T. M., Zhou, Y., Dada, L., Sulo, J., Flagan, R. C., Kulmala, M., Petäjä, T., and Lehtipalo, K.: Overview of
 1057 measurements and current instrumentation for 1–10 nm aerosol particle number size distributions,
 1058 <https://doi.org/10.1016/j.jaerosci.2020.105584>, 1 October 2020.

1059 Karl, M., Leck, C., Gross, A., and Pirjola, L.: A study of new particle formation in the marine boundary layer
 1060 over the central Arctic Ocean using a flexible multicomponent aerosol dynamic model, *Tellus B Chem Phys*
 1061 *Meteorol*, 64, <https://doi.org/10.3402/tellusb.v64i0.17158>, 2012.

1062 Karlsson, L., Krejci, R., Koike, M., Ebell, K., and Zieger, P.: The role of nanoparticles in Arctic cloud formation,
 1063 <https://doi.org/10.5194/acp-2020-417>, 29 May 2020.

1064 Karlsson, L., Krejci, R., Koike, M., Ebell, K., and Zieger, P.: A long-term study of cloud residuals from
 1065 low-level Arctic clouds, *Atmos Chem Phys*, 21, 8933–8959, <https://doi.org/10.5194/acp-21-8933-2021>, 2021.

1066 Kay, J. E. and Gettelman, A.: Cloud influence on and response to seasonal Arctic sea ice loss, *Journal of*
 1067 *Geophysical Research Atmospheres*, 114, <https://doi.org/10.1029/2009JD011773>, 2009.

1068 Kecorius, S., Vogl, T., Paasonen, P., Lampilahti, J., Rothenberg, D., Wex, H., Zeppenfeld, S., Van Pinxteren, M.,
 1069 Hartmann, M., Henning, S., Gong, X., Welti, A., Kulmala, M., Stratmann, F., Herrmann, H., and Wiedensohler,
 1070 A.: New particle formation and its effect on cloud condensation nuclei abundance in the summer Arctic: A case

1071 study in the Fram Strait and Barents Sea, *Atmos Chem Phys*, 19, 14339–14364,
1072 <https://doi.org/10.5194/acp-19-14339-2019>, 2019.

1073 Kerminen, V. M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi, F.: Atmospheric new particle
1074 formation and growth: Review of field observations, <https://doi.org/10.1088/1748-9326/aadf3c>, 27 September
1075 2018.

1076 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L.,
1077 Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer,
1078 D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C.,
1079 Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K.,
1080 Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela,
1081 A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A.,
1082 Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M.,
1083 Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and
1084 galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–435,
1085 <https://doi.org/10.1038/nature10343>, 2011.

1086 Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C.,
1087 Almeida, J., Trostl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A. K.,
1088 Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin,
1089 A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf,
1090 M., Kurten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Perakyla, O.,
1091 Piel, F., Petaja, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo,
1092 L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipila, M., Steiner, G., Stozhkov, Y., Stratmann,
1093 F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P.
1094 M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala,
1095 M., Carslaw, K. S., and Curtius, J.: Ion-induced nucleation of pure biogenic particles, *Nature*, 533, 521–526,
1096 <https://doi.org/10.1038/nature17953>, 2016.

1097 Kolesar, K. R., Cellini, J., Peterson, P. K., Jefferson, A., Tuch, T., Birmili, W., Wiedensohler, A., and Pratt, K. A.:
1098 Effect of Prudhoe Bay emissions on atmospheric aerosol growth events observed in Utqiagvik (Barrow), Alaska,
1099 *Atmos Environ*, 152, 146–155, <https://doi.org/10.1016/j.atmosenv.2016.12.019>, 2017a.

1100 Kolesar, K. R., Cellini, J., Peterson, P. K., Jefferson, A., Tuch, T., Birmili, W., Wiedensohler, A., and Pratt, K. A.:
 1101 Effect of Prudhoe Bay emissions on atmospheric aerosol growth events observed in Utqiaġvik (Barrow), Alaska,
 1102 *Atmos Environ*, 152, 146–155, <https://doi.org/10.1016/j.atmosenv.2016.12.019>, 2017b.

1103 Kristensson, adam, Johansson, martin, swietlicki, erik, Kivekäs, niku, hussein, tareq, nieminen, tuomo, Kulmala,
 1104 markku, and Dal maso, miikka: nanomap: Geographical mapping of atmospheric new-particle formation through
 1105 analysis of particle number size distribution and trajectory data, 2014.

1106 Kuang, C., Riipinen, I., Sihto, S. L., Kulmala, M., McCormick, A. V., and McMurry, P. H.: An improved
 1107 criterion for new particle formation in diverse atmospheric environments, *Atmos Chem Phys*, 10, 8469–8480,
 1108 <https://doi.org/10.5194/acp-10-8469-2010>, 2010.

1109 Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P.,
 1110 Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V. M.: Measurement
 1111 of the nucleation of atmospheric aerosol particles, *Nat Protoc*, 7, 1651–1667,
 1112 <https://doi.org/10.1038/nprot.2012.091>, 2012.

1113 Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., and Nieminen, T.: Direct
 1114 Observations of Atmospheric Aerosol Nucleation, *Science* (1979), 339, 943–946,
 1115 <https://doi.org/10.1126/science.1227385>, 2013.

1116 Leaitch, W. R., Sharma, S., Huang, L., Toom-Sauntry, D., Chivulescu, A., Macdonald, A. M., Von Salzen, K.,
 1117 Pierce, J. R., Bertram, A. K., Schroder, J. C., Shantz, N. C., Chang, R. Y. W., and Norman, A. L.: Dimethyl
 1118 sulfide control of the clean summertime Arctic aerosol and cloud, *Elementa*, 1,
 1119 <https://doi.org/10.12952/journal.elementa.000017>, 2013.

1120 Leaitch, W. R., Korolev, A., Aliabadi, A. A., Burkart, J., Willis, M. D., Abbatt, J. P. D., Bozem, H., Hoor, P.,
 1121 Köllner, F., Schneider, J., Herber, A., Konrad, C., and Brauner, R.: Effects of 20-100nm particles on liquid
 1122 clouds in the clean summertime Arctic, *Atmos Chem Phys*, 16, 11107–11124,
 1123 <https://doi.org/10.5194/acp-16-11107-2016>, 2016.

1124 Lee, H., Lee, K., Rene Lunder, C., Krejci, R., Aas, W., Park, J., Park, K. T., Yong Lee, B., Yoon, Y. J., and Park,
 1125 K.: Atmospheric new particle formation characteristics in the Arctic as measured at Mount Zeppelin, Svalbard,
 1126 from 2016 to 2018, *Atmos Chem Phys*, 20, 13425–13441, <https://doi.org/10.5194/acp-20-13425-2020>, 2020.

1127 Lee, K., Kim, J. S., Park, K. T., Park, M. J., Jang, E., Gudmundsson, K., Olafsdottir, S. R., Olafsson, J., Yoon, Y.
 1128 J., Lee, B. Y., Kwon, S. Y., and Kam, J.: Observational evidence linking ocean sulfur compounds to atmospheric
 1129 dimethyl sulfide during Icelandic Sea phytoplankton blooms, *Science of the Total Environment*, 879,
 1130 <https://doi.org/10.1016/j.scitotenv.2023.163020>, 2023.

1131 Li, J., Carlson, B. E., Yung, Y. L., Lv, D., Hansen, J., Penner, J. E., Liao, H., Ramaswamy, V., Kahn, R. A.,
 1132 Zhang, P., Dubovik, O., Ding, A., Lacis, A. A., Zhang, L., and Dong, Y.: Scattering and absorbing aerosols in the
 1133 climate system, <https://doi.org/10.1038/s43017-022-00296-7>, 1 June 2022.

1134 Manninen, H. E., Franchin, A., Schobesberger, S., Hirsikko, A., Hakala, J., Skromulis, A., Kangasluoma, J., Ehn,
 1135 M., Junninen, H., Mirme, A., Mirme, S., Sipilä, M., Petäjä, T., Worsnop, D. R., and Kulmala, M.:
 1136 Characterisation of corona-generated ions used in a Neutral cluster and Air Ion Spectrometer (NAIS), *Atmos*
 1137 *Meas Tech*, 4, 2767–2776, <https://doi.org/10.5194/amt-4-2767-2011>, 2011.

1138 Maturilli, M. and Ebell, K.: Twenty-five years of cloud base height measurements by ceilometer in Ny-Ålesund,
 1139 Svalbard, *Earth Syst Sci Data*, 10, 1451–1456, <https://doi.org/10.5194/essd-10-1451-2018>, 2018.

1140 Mauritsen, T., Sedlar, J., Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O.
 1141 G., Brooks, I. M., and Swietlicki, E.: An Arctic CCN-limited cloud-aerosol regime, *Atmos Chem Phys*, 11,
 1142 165–173, <https://doi.org/10.5194/acp-11-165-2011>, 2011.

1143 Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Atmospheric Chemistry and
 1144 Physics Impact of nucleation on global CCN, *Atmos. Chem. Phys*, 8601–8616 pp., 2009.

1145 Mirme, A., Tamm, E., Mordas, G., Vana, M., Uin, J., Mirme, S., Bernotas, T., Laakso, L., Kulmala, M., and
 1146 Hirsikko, A.: A wide-range multi-channel air ion spectrometer, *Boreal Environment Research*, 12, 247–264,
 1147 2007.

1148 Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS - A spectrometer for the
 1149 measurement of cluster ion and nanometer aerosol size distributions, *Atmos Meas Tech*, 6, 1061–1071,
 1150 <https://doi.org/10.5194/amt-6-1061-2013>, 2013.

1151 Motos, G., Freitas, G., Georgakaki, P., Wieder, J., Li, G., Aas, W., Lunder, C., Krejci, R., Pasquier, J. T.,
 1152 Henneberger, J., David, R. O., Ritter, C., Mohr, C., Zieger, P., and Nenes, A.: Aerosol and dynamical
 1153 contributions to cloud droplet formation in Arctic low-level clouds, *Atmos Chem Phys*, 23, 13941–13956,
 1154 <https://doi.org/10.5194/acp-23-13941-2023>, 2023a.

1155 Motos, G., Freitas, G., Georgakaki, P., Wieder, J., Li, G., Aas, W., Lunder, C., Krejci, R., Pasquier, J. T.,
 1156 Henneberger, J., David, R. O., Ritter, C., Mohr, C., Zieger, P., and Nenes, A.: Aerosol and dynamical
 1157 contributions to cloud droplet formation in Arctic low-level clouds, *Atmos Chem Phys*, 23, 13941–13956,
 1158 <https://doi.org/10.5194/acp-23-13941-2023>, 2023b.

1159 Nguyen, Q. T., Glasius, M., Sørensen, L. L., Jensen, B., Skov, H., Birmili, W., Wiedensohler, A., Kristensson,
 1160 A., Nøjgaard, J. K., and Massling, A.: Seasonal variation of atmospheric particle number concentrations, new
 1161 particle formation and atmospheric oxidation capacity at the high Arctic site Villum Research Station, Station
 1162 Nord, *Atmos Chem Phys*, 16, 11319–11336, <https://doi.org/10.5194/acp-16-11319-2016>, 2016.

1163 Nieminen, T., Kerminen, V. M., Petäjä, T., Aalto, P. P., Arshinov, M., Asmi, E., Baltensperger, U., Beddows, D.
 1164 C. S., Paul Beukes, J., Collins, D., Ding, A., Harrison, R. M., Henzing, B., Hooda, R., Hu, M., Hörrak, U.,
 1165 Kivekäs, N., Komsaare, K., Krejci, R., Kristensson, A., Laakso, L., Laaksonen, A., Richard Leaitch, W.,
 1166 Lihavainen, H., Mihalopoulos, N., Németh, Z., Nie, W., O’Dowd, C., Salma, I., Sellegri, K., Svenningsson, B.,
 1167 Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler, A., Wu, Z., Virtanen, A., and
 1168 Kulmala, M.: Global analysis of continental boundary layer new particle formation based on long-term
 1169 measurements, *Atmos Chem Phys*, 18, 14737–14756, <https://doi.org/10.5194/acp-18-14737-2018>, 2018.

1170 Palm, S. P., Strey, S. T., Spinhirne, J., and Markus, T.: Influence of Arctic sea ice extent on polar cloud fraction
 1171 and vertical structure and implications for regional climate, *Journal of Geophysical Research Atmospheres*, 115,
 1172 <https://doi.org/10.1029/2010JD013900>, 2010.

1173 Park, K. T., Lee, K., Kim, T. W., Yoon, Y. J., Jang, E. H., Jang, S., Lee, B. Y., and Hermansen, O.: Atmospheric
 1174 DMS in the Arctic Ocean and Its Relation to Phytoplankton Biomass, *Global Biogeochem Cycles*, 32, 351–359,
 1175 <https://doi.org/10.1002/2017GB005805>, 2018a.

1176 Park, K. T., Lee, K., Kim, T. W., Yoon, Y. J., Jang, E. H., Jang, S., Lee, B. Y., and Hermansen, O.: Atmospheric
 1177 DMS in the Arctic Ocean and Its Relation to Phytoplankton Biomass, *Global Biogeochem Cycles*, 32, 351–359,
 1178 <https://doi.org/10.1002/2017GB005805>, 2018b.

1179 Pei, Q., Saikawa, E., Kaspari, S., Widory, D., Zhao, C., Wu, G., Loewen, M., Wan, X., Kang, S., Wang, X.,
 1180 Zhang, Y. L., and Cong, Z.: Sulfur aerosols in the Arctic, Antarctic, and Tibetan Plateau: Current knowledge and
 1181 future perspectives, <https://doi.org/10.1016/j.earscirev.2021.103753>, 1 September 2021.

1182 Pirjola, L., O'Dowd, C. D., Brooks, I. M., and Kulmala, M.: Can new particle formation occur in the clean
 1183 marine boundary layer?, *Journal of Geophysical Research: Atmospheres*, 105, 26531–26546,
 1184 <https://doi.org/10.1029/2000JD900310>, 2000.

1185 Platt, S. M., Hov, Ø., Berg, T., Breivik, K., Eckhardt, S., Eleftheriadis, K., Evangeliou, N., Fiebig, M., Fisher, R.,
 1186 Hansen, G., Hansson, H. C., Heintzenberg, J., Hermansen, O., Heslin-Rees, D., Holmén, K., Hudson, S.,
 1187 Kallenborn, R., Krejci, R., Krognes, T., Larssen, S., Lowry, D., Myhre, C. L., Lunder, C., Nisbet, E., Nizzetto, P.
 1188 B., Park, K. T., Pedersen, C. A., Pfaffhuber, K. A., Röckmann, T., Schmidbauer, N., Solberg, S., Stohl, A.,
 1189 Ström, J., Svendby, T., Tunved, P., Tørnkvist, K., Van Der Veen, C., Vratolis, S., Yoon, Y. J., Yttri, K. E., Zieger,
 1190 P., Aas, W., and Tørseth, K.: Atmospheric composition in the European Arctic and 30 years of the Zeppelin
 1191 Observatory, Ny-Ålesund, *Atmos Chem Phys*, 22, 3321–3369, <https://doi.org/10.5194/acp-22-3321-2022>, 2022.

1192 Pöhlker, M. L., Zhang, M., Campos Braga, R., Krüger, O. O., Pöschl, U., and Ervens, B.: Aitken mode particles
 1193 as CCN in aerosol- And updraft-sensitive regimes of cloud droplet formation, *Atmos Chem Phys*, 21,
 1194 11723–11740, <https://doi.org/10.5194/acp-21-11723-2021>, 2021.

1195 Price, R., Baccarini, A., Schmale, J., Zieger, P., Brooks, I. M., Field, P., and Carslaw, K. S.: Late summer
 1196 transition from a free-Tropospheric to boundary layer source of Aitken mode aerosol in the high Arctic, *Atmos*
 1197 *Chem Phys*, 23, 2927–2961, <https://doi.org/10.5194/acp-23-2927-2023>, 2023.

1198 Quinn, P. K., Miller, T. L., Bates, T. S., Ogren, J. A., Andrews, E., and Shaw, G. E.: A 3-year record of
 1199 simultaneously measured aerosol chemical and optical properties at Barrow, Alaska, *Journal of Geophysical*
 1200 *Research: Atmospheres*, 107, <https://doi.org/10.1029/2001jd001248>, 2002.

1201 Schmale, J. and Baccarini, A.: Progress in Unraveling Atmospheric New Particle Formation and Growth Across
 1202 the Arctic, *Geophys Res Lett*, 48, <https://doi.org/10.1029/2021GL094198>, 2021.

1203 Schmale, J., Sharma, S., Decesari, S., Pernov, J., Massling, A., Hansson, H. C., Von Salzen, K., Skov, H.,
 1204 Andrews, E., Quinn, P. K., Upchurch, L. M., Eleftheriadis, K., Traversi, R., Gilardoni, S., Mazzola, M., Laing, J.,
 1205 and Hopke, P.: Pan-Arctic seasonal cycles and long-term trends of aerosol properties from 10 observatories,
 1206 *Atmos Chem Phys*, 22, 3067–3096, <https://doi.org/10.5194/acp-22-3067-2022>, 2022.

1207 Shaw, G. E.: The Arctic Haze Phenomenon, 2403–2414 pp.,
 1208 [https://doi.org/https://doi.org/10.1175/1520-0477\(1995\)076<2403:TAHP>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0477(1995)076<2403:TAHP>2.0.CO;2), 1995.

1209 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT
1210 atmospheric transport and dispersion modeling system, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 1
1211 December 2015.

1212 Ström, J., Umegård, J., Tørseth, K., Tunved, P., Hansson, H. C., Holmén, K., Wismann, V., Herber, A., and
1213 König-Langlo, G.: One year of particle size distribution and aerosol chemical composition measurements at the
1214 Zeppelin Station, Svalbard, March 2000–March 2001, *Physics and Chemistry of the Earth*, 28, 1181–1190,
1215 <https://doi.org/10.1016/j.pce.2003.08.058>, 2003.

1216 Ström, J., Engvall, A. C., Delbart, F., Krejci, R., and Treffeisen, R.: On small particles in the Arctic summer
1217 boundary layer: Observations at two different heights near Ny-Ålesund, Svalbard, *Tellus B Chem Phys*
1218 *Meteorol*, 61 B, 473–482, <https://doi.org/10.1111/j.1600-0889.2008.00412.x>, 2009.

1219 Tammet, H., Komsaare, K., and Hörrak, U.: Intermediate ions in the atmosphere, *Atmos Res*, 135–136, 263–273,
1220 <https://doi.org/10.1016/j.atmosres.2012.09.009>, 2014.

1221 Tunved, P., Ström, J., and Krejci, R.: Arctic aerosol life cycle: Linking aerosol size distributions observed
1222 between 2000 and 2010 with air mass transport and precipitation at Zeppelin station, Ny-Ålesund, Svalbard,
1223 *Atmos Chem Phys*, 13, 3643–3660, <https://doi.org/10.5194/acp-13-3643-2013>, 2013.

1224 Vihma, T., Jaagus, J., Jakobson, E., and Palo, T.: Meteorological conditions in the Arctic Ocean in spring and
1225 summer 2007 as recorded on the drifting ice station Tara, *Geophys Res Lett*, 35,
1226 <https://doi.org/10.1029/2008GL034681>, 2008.

1227 Wagner, R., Manninen, H. E., Franchin, A., Lehtipalo, K., Mirme, S., Steiner, G., Petäjä, T., and Kulmala, M.:
1228 On the accuracy of ion measurements using a Neutral cluster and Air Ion Spectrometer, 2016.

1229 Wagner, R., Yan, C., Lehtipalo, K., Duplissy, J., Nieminen, T., Kangasluoma, J., Ahonen, L. R., Dada, L.,
1230 Kontkanen, J., Manninen, H. E., Dias, A., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A. K., Bianchi, F.,
1231 Brilke, S., Buenrostro Mazon, S., Chen, X., Draper, D. C., Fischer, L., Frege, C., Fuchs, C., Garmash, O.,
1232 Gordon, H., Hakala, J., Heikkinen, L., Heinritzi, M., Hofbauer, V., Hoyle, C. R., Kirkby, J., Kürten, A.,
1233 Kvashnin, A. N., Laurila, T., Lawler, M. J., Mai, H., Makhmutov, V., Mauldin, R., Molteni, U., Nie, W.,
1234 Nie, W., Ojdanic, A., Onnela, A., Piel, F., Quéléver, L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S.,
1235 Sengupta, K., Simon, M., Stolzenburg, D., Stozhkov, Y., Tröstl, J., Viisanen, Y., Vogel, A. L., Wagner, A. C.,
1236 Xiao, M., Ye, P., Baltensperger, U., Curtius, J., Donahue, N. M., Flagan, R. C., Gallagher, M., Hansel, A., Smith,

- 1237 J. N., Tomé, A., Winkler, P. M., Worsnop, D., Ehn, M., Sipilä, M., Kerminen, V. M., Petäjä, T., and Kulmala, M.:
 1238 The role of ions in new particle formation in the CLOUD chamber, *Atmos Chem Phys*, 17, 15181–15197,
 1239 <https://doi.org/10.5194/acp-17-15181-2017>, 2017.
- 1240 Von Der Weiden, S.-L., Drewnick, F., and Borrmann, S.: Atmospheric Measurement Techniques Particle Loss
 1241 Calculator-a new software tool for the assessment of the performance of aerosol inlet systems, *Atmos. Meas.*
 1242 *Tech*, 479–494 pp., 2009.
- 1243 Wiedensohler, A., Covert, D. S., Swietlicki, E., Aalto, P., Heintzenberg, J., and Leck, C.: Occurrence of an
 1244 ultrafine particle mode less than 20 nm in diameter in the marine boundary layer during Arctic summer and
 1245 autumn, *Tellus B Chem Phys Meteorol*, 48, 213–222, <https://doi.org/10.3402/tellusb.v48i2.15887>, 1996.
- 1246 Willis, M. D., Leaitch, W. R., and Abbatt, J. P. D.: Processes Controlling the Composition and Abundance of
 1247 Arctic Aerosol, <https://doi.org/10.1029/2018RG000602>, 1 December 2018.
- 1248 Xavier C., Wollesen de jonge R., Jokinen T., Beck L., Sipilä M., Olenius T., and Roldin P.: Role of
 1249 Iodine-Assisted Aerosol Particle Formation in Antarctica, *Environmental Science & Technology*, 58 (17),
 1250 7314–7324, DOI: 10.1021/acs.est.3c09103, 2024.
- 1251 Yu, F. and Turco, R. P.: From molecular clusters to nanoparticles: Role of ambient ionization in tropospheric
 1252 aerosol formation, *Journal of Geophysical Research Atmospheres*, 106, 4797–4814,
 1253 <https://doi.org/10.1029/2000JD900539>, 2001.
- 1254 Ziemba, L. D., Dibb, J. E., Griffin, R. J., Huey, L. G., and Beckman, P.: Observations of particle growth at a
 1255 remote, Arctic site, *Atmos Environ*, 44, 1649–1657, <https://doi.org/10.1016/j.atmosenv.2010.01.032>, 2010.
- 1256

