

Long-term PM trends at boreal forest site in southern Finland from three different measurement techniques

Ilona Ylivinkka^{1,2}, Helmi-Marja Keskinen¹, Lauri R. Ahonen^{1,2}, Liine Heikkinen^{1,3,4}, Pasi P. Aalto¹, Tuomo Nieminen^{1,5}, Katrianne Lehtipalo^{1,6}, Krista Luoma^{1,6}, Sujai Banerji¹, Juho Aalto^{2,7}, Janne Levula¹, Jutta Kesti^{1,6}, Ekaterina Ezhova¹, Markku Kulmala¹, and Tuukka Petäjä¹

¹Institute for Atmospheric and Earth System Research (INAR) / Physics, Faculty of Science, University of Helsinki, Helsinki, 00014, Finland

²Station for Measuring Ecosystem - Atmosphere Relations II (SMEAR II), University of Helsinki, Korkeakoski, 35500, Finland

³Department of Environmental Science, Stockholm University, Sweden

⁴Bolin Centre for Climate Research, Stockholm University, Sweden

⁵Department of Physics, Faculty of Science, University of Helsinki, Finland

⁶Finnish Meteorological Institute, 00560 Helsinki, Finland

⁷Institute for Atmospheric and Earth System Research (INAR) / Forest Sciences, Faculty of Agriculture and Forestry, University of Helsinki, Helsinki, 00014, Finland

Correspondence to: Ilona Ylivinkka (ilona.ylivinkka@helsinki.fi)

Abstract. Three independent particulate matter (PM) mass concentration measurements and their long-term (2005–2020) trends were compared at the Station for Measuring Ecosystem–Atmosphere Relations (SMEAR II, Hyytiälä, Finland). The different methods (gravimetric method with a cascade impactor, Synchronized Hybrid Ambient Real-time Particulate Monitor (SHARP; only PM₁₀), and calculated PM concentration from combined Differential Mobility Particle Sizer (DMPS) and Aerosol Particle Sizer (APS) particle number size distribution data) showed good correlation (Pearson’s correlation coefficient approximately 0.8) in all size classes (PM₁, PM_{2.5} and PM₁₀). The mass concentrations in all PM classes were the highest in summer and the lowest in autumn and winter. Statistically significant (Mann–Kendall test) declining annual trends were observed in DMPS+APS and impactor data in all size classes, ranging from -0.021 to -0.036 $\mu\text{g m}^{-3} \text{y}^{-1}$. While DMPS+APS method indicated statistically significant decline also in all seasons, the decline in impactor data was statistically significant only in spring and winter. SHARP data could not be used for trend estimation due to the change in inlet heating temperature, affecting the measured PM₁₀

concentrations. Seasonally, the decline was smallest in summer, which follows the trends observed also in SO₂ and NO_x concentrations. The results underline both the summertime dominance of biogenic sources for the aerosol mass concentration in the rural boreal forest environment and the reduction of anthropogenic pollution due to the EU level restrictions for improved air quality.

1 Introduction

Particulate matter (PM) concentrations are monitored worldwide, because they are connected to health effects, such as asthma and cardiovascular diseases, and premature deaths (Pope et al., 2003; Shiraiwa et al., 2017; WHO, 2021). The increased knowledge regarding the relationship between air pollution and mortality have resulted in air pollution regulations, which additionally aim to decrease inequality related to air pollution exposure (Wang et al., 2017; WHO, 2021). Besides the adverse health effects, aerosol particles can also scatter or absorb radiation and participate in cloud formation and processing, thus affecting the Earth's climate (IPCC, 2021). While the overall effect of aerosol particles on climate is considered to be cooling, **radiative forcing due to aerosol particles and especially due to aerosol–cloud–radiation interactions is uncertain** (IPCC, 2021).

PM measurements are divided into size classes based on the aerodynamic diameter of the particles: PM₁, PM_{2.5}, and PM₁₀ with upper maximum diameters of particles 1 µm, 2.5 µm, and 10 µm, respectively. **The PM mass concentration in these classes is** the total mass of particles below the limiting size. The size of aerosol particles is a critical parameter, both in terms of their climate (e.g., Pöschl, 2005; Dusek et al., 2006) and health effects (Schraufnagel, 2020). In principle, the smaller the particles are, the deeper they can penetrate in the human respiratory system and thus end up also in other organs than lungs (Pope et al., 2003; Maynard & Kuempel, 2005). The smallest particles have only a minor contribution to the aerosol mass concentration, but they dominate the particle number concentration. In climate perspective, the most relevant particles **are larger** than about 50–100 nm, since **those** can act as cloud condensation nuclei **as well as** scatter or absorb radiation (IPCC, 2021).

Aerosol particles have both natural and anthropogenic sources. Additionally, particles can be transported over hundreds or thousands of kilometers, since the lifetime of PM in the atmosphere is days to weeks, depending on the size, composition, and source region of the particles (Seinfeld and Pandis, 2006; Manavi et al., 2025). Primary aerosol particles consist mostly of particles from traffic and industry (e.g., black carbon (BC)), or from natural sources (e.g., volcanic ash, sea-spray, dust, and pollen), and they contribute to all PM classes. Secondary aerosol particles are formed in the atmosphere from gas-phase precursor vapors (e.g. Kulmala et al., 2013). These particles eventually grow to larger sizes, contributing particularly to the accumulation mode, and thereby to PM₁.

At SMEAR II, organic aerosol (OA) from oxidized biogenic volatile organic compounds (VOCs), most importantly monoterpenes from the surrounding forest (Rinne et al., 2005), is the most abundant PM₁ component (Jimenez et al., 2009; Heikkinen et al., 2020). The emission rates of monoterpenes are boosted by warm temperatures (Guenther et al., 1993), which is also observed in the OA mass concentrations (Heikkinen et al., 2021; Yli-Juuti et al., 2021). Sulfate, another key PM₁ component at SMEAR II and globally, is formed, e.g., upon oxidation from sulfur dioxide (SO₂), mostly emitted by industry (Seinfeld and Pandis, 2006). Nitrate aerosol mass concentrations, mostly prevalent in agricultural or urban environments, are therefore less abundant at SMEAR II (Makkonen et al., 2014).

The European Union has regulated the exposure on air pollution since 2005 via air quality directives (https://environment.ec.europa.eu/topics/air/air-quality/eu-air-quality-standards_en; accessed: 31 Jul 2025). The air quality directives concern basic pollutants: PM, trace gases (SO₂, NO₂, O₃, CO, benzene, and polyaromatic hydrocarbons) as well as heavy metals (Pb, As, Ni, and Cd). Originally legislation on PM concerned only PM₁₀ concentration (yearly average concentration was limited to 40 µg m⁻³), but in 2010 target value was set for PM_{2.5} concentration (25 µg m⁻³ and limited to 20 µg m⁻³ in 2020).

In the end on 2024, The Ambient Air Quality Directive was revised (2024/2881), forcing further reductions for targets values of many pollutants, including PM₁₀, PM_{2.5}, O₃, SO₂, CO, and benzene. Additionally, the new air quality directive introduces advanced measurement parameters, such as aerosol number concentration, aerosol size distribution, BC, and oxidative potential, to address i.a. the harmfulness of small aerosol particles. Air quality supersite concept was implemented as well (Kuula et al., 2021), aiming to compare the health

impacts of the ultrafine particles and PM_{2.5} in urban and rural supersites. In addition, the EU Commission mandates measurements of ultrafine (defined as particles between 10 to 100 nm in diameter) and BC concentrations in the vicinity of air pollution hotspots.

Several studies have already reported declining PM concentrations in Europe (Barmpadimos et al., 2011; 2012; Cusack et al., 2012; Pandolfi et al., 2016), ranging approximately from -0.008 PM_{2.5} trend in Po Valley, Italy (Bigi & Ghermandi, 2016) to -4.11 µg m⁻³ yr⁻¹ in (sub)urban Germany (Guerreiro et al., 2014). In some cases, also increasing trends have also been measured in relation to increased emissions from, for example, household combustion and agriculture (Guerreiro et al., 2014). Declining trends are related to the legislation on air quality as well as improved technology in industry, traffic, and heating (Spindler et al., 2004; Anttila & Tuovinen, 2010; Barmpadimos et al., 2011; Cusack et al., 2012; Keuken et al., 2012). In Finland, the PM concentrations have been declining during the past decades and generally are well below the limit values (Laakso et al., 2003; Anttila, 2020).

Techniques for measuring aerosol mass concentrations have improved remarkably during the last decades (Van Dingenen et al., 2004; Occhipinti & Oluwasanya, 2017; Shukla & Aggarwal, 2022). Most of the PM measurements have traditionally been done by offline gravimetric analyses where particle size classes are separated, e.g., by impactor (Laakso et al., 2003) or special high-volume samplers (Barmpadimos et al., 2011). The offline methods are quite laborious as their sampling time is up to few days and weighing is done manually. Thus, PM concentrations are nowadays more commonly measured with online techniques, such as tapered element oscillating microbalance (TEOM) with the Continuous Ambient Particulate Monitor and Synchronized Hybrid Ambient Real-time Particulate monitor (SHARP) (Laakso et al., 2008; Chen et al., 2018, Waldén et al., 2010). Besides the direct mass measurements, the particle mass can be calculated from the particle number size distribution with assumptions regarding particles' shape and density (Neusüß et al., 2000).

The aim of this work is threefold. First, we compare the PM concentrations obtained from gravimetric impactor, online mass analyzer SHARP and from the particle number size distribution to explore their applicability for continuous PM measurements. Second, we report for the first time long-term (2005–2020) measurements of PM₁₀, PM_{2.5} and PM₁ at SMEAR II, Finland, and explore the overall concentration levels as well as selected specific episodes. Third, we estimate the trends of the PM concentrations separately for each season

and the impact of the EU legislation on the PM trends. Quality controlled data on aerosol particle mass concentration in a boreal background station enable us to explore the role of local, regional and global phenomena controlling the aerosol mass concentration in the region. This work continues the analysis presented in Keskinen et al. (2020) with updated datasets and revised analysis methods.

2 Methods

2.1 Measurement station

The measurements were performed at SMEAR II, located in Hyytiälä in southern Finland (61°51'N, 24°17'E; 181 m a.s.l.; Fig. 1a). Hyytiälä is a rural background measurement site with low local anthropogenic emissions (Hari and Kulmala, 2005). A photo of the homogeneous 60-year-old Scots pine stand surrounding SMEAR II is presented in Fig. 1c. The nearest cities are Tampere (50 km southwest; 249 000 inhabitants) and Jyväskylä (90 km northeast; 146 000 inhabitants).

The station is equipped with instruments for continuous and comprehensive measurements of interactions between the forest ecosystem and atmosphere (Hari and Kulmala et al., 2005). SMEAR II is part of the European Aerosols, Clouds, and Trace gases Research Infrastructure (ACTRIS; Laj et al. 2024; <https://www.actris.eu/>, accessed 07 Aug 2025). The presented measurements are conducted inside the canopy with total suspended particulates (TSP) or PM₁₀ design inlets for the different aerosol measurements on the roof of the aerosol cottage (Fig. 1b). Winter at SMEAR II is defined to be from December to February (DJF), spring is from March to May (MAM), summer from June to August (JJA) and autumn from September to November (SON). Note that winter has January and February data from the following year. Due to the data availability, measurements start from spring 2005 and in the end of the measurement period winter includes only December 2020.




Figure 1: (a) The location of SMEAR II (© OpenStreetMap contributors 2020. Distributed under a Creative Commons BY-SA License), (b) cottage for aerosol instrumentation, and (c) a photo of the surrounding region around SMEAR II.

2.2 Weighing-based mass measurements with cascade impactor

PM measurements with gravimetric cascade impactor started in late 1990s at SMEAR II. The impactor has an unheated TSP inlet with stainless-steel tube, placed at 5 m height above the ground. The cascade impactor has three stages with impactor cut points at $10\ \mu\text{m}$ (PM_{10}), $2.5\ \mu\text{m}$ ($\text{PM}_{2.5}$) and $1\ \mu\text{m}$ (PM_1) (Dekati PM10 impactor) (Berner and Luerzer, 1980). The sample air flow rate during collection is $30\ \text{l min}^{-1}$. Collection substrates are 25 mm polycarbonate membranes (Nuclepore 800 203) without holes. At the last stage there is a 47 mm Teflon filter with $2\ \mu\text{m}$ pore size (R2P J047) from Pall Corporation. To prevent the bouncing back of the particles from the collection substrates, the membranes are greased with Apiezon L vacuum grease diluted in toluene. **The impactor samples are collected for two to three days, before the filters are taken to a clean laboratory room, where they are dried in**

laminar flow hood for at least two hours before weighting to get the mass distribution. The samples are stored in a freezer for occasional further analyses.

2.3 Online mass measurements with SHARP

The Synchronized Hybrid Ambient Real-time Particulate Monitor (SHARP, Thermo Scientific, Model 5030) is a real-time particulate monitor measuring at 1 s time resolution (Goohs et al., 2009). SHARP combines light scattering photometry and β -ray attenuation for continuous PM₁₀ measurement. In SHARP the light scattering signal (nephelometer) is automatically calibrated against the beta attenuation mass sensor. The sample line inlet is placed on the roof of the cottage at 6 m height above the ground level and its flow rate is 16.7 l min⁻¹. The sample line is heated to reduce the humidity of the sample air. The temperature was fixed to 45 °C until August 2016 and to 35 °C after that. The sensitivity of the instrument was calibrated regularly with a specific foil. Sampling with SHARP at SMEAR II started in 2012.

2.4 Aerosol mass derived from the particle size distribution

The aerosol mass concentration for different size classes PM₁₀, PM_{2.5} and PM₁ can also be estimated by combining the number size distributions measured with Differential Mobility Particle Sizer (DMPS) and Aerodynamic Particle Sizer (APS) and calculating the mass by assuming that the particles are spherical and have a constant density. The instrument set-ups for DMPS and APS are described in detail by Aalto et al. (2001). Briefly, the twin-DMPS consists of a long and a short Vienna type Differential Mobility Analyzers (DMA) and two condensation particle counters (CPC; TSI 3025 and TSI 3775). The DMPS inlet is placed on the roof of the cottage at 8 m height and APS inlet at 5 m above ground level. The DMPS and APS systems provide aerosol number size distribution with a 10 min time resolution. The APS inlet line is heated to 35 °C, similarly as the SHARP inlet line. In the DMPS system, the sheath flow is dried with a silica diffusion dryer. The relative humidity of the sheath flow was kept below 40 %. The calibrations of both instruments are checked regularly using polystyrene latex spheres.

At SMEAR II, the DMPS measures the aerosol number size distribution in the electrical mobility equivalent diameter range of 3–1000 nm (Aalto et al., 2001). The APS (TSI 3320) measures the aerodynamic particle size distribution of particles with aerodynamic diameter within the range of 0.5–20 μ m (Peters et al., 2006). To have comparable particle size

distributions, we **converted** the aerodynamic diameter (d_a) of the APS to mobility equivalent diameter (d_m):

209

$$d_m = \sqrt{\frac{\rho_0}{\rho_p}} d_a, \quad (1)$$

211

where ρ_p is the density of the particle and ρ_0 is the unit density of the particle (1 g cm^{-3}). The density of the particles is assumed to be 1.5 g cm^{-3} (Saarikoski et al., 2005; Kannosto et al., 2008), **but we additionally calculated the mass concentrations using 1.1 and 2.0 g cm^{-3} densities, which are the minimum and maximum densities of accumulation mode sized particles at SMEAR II (Kannosto et al. 2008) to understand the importance of constant density assumption to the particle mass.** The mass of the particles measured with DMPS+APS is calculated as:

219

$$m = \frac{1}{6} \rho_p \pi d_m^3. \quad (2)$$

221

The mass concentrations (PM_{10} , $\text{PM}_{2.5}$ and PM_{10}) were then calculated by integrating over the corresponding size range:

224

$$\text{PM}_i = \int_{0.6 \mu\text{m}}^{0.6 \mu\text{m}} N_{\text{DMPS}} \cdot m_{\text{DMPS}} dd_m + \int_{0.6 \mu\text{m}}^i N_{\text{APS}} \cdot m_{\text{APS}} dd_m \quad (3)$$

226

In practice, we utilized DMPS data from 0.003 to 0.6 μm and APS size distribution from 0.6 μm to 1 μm , 2.5 μm or 10 μm , depending on the mass fraction in question. Typical size distributions for different seasons are presented in Fig. 2.

230




Figure 2: Seasonal median number size distributions for 2005–2020 at SMEAR II measured with a combination of DMPS and APS with a constant density assumption. The dash-dotted line indicates the crossover size between the instrument data to determine integrated mass concentrations.

2.5 Ancillary data

SO₂ and NO_x were measured at 16.8 m height above ground level at SMEAR II with gas analyzers by Thermo Fisher Scientific Inc., USA. SO₂ was measured with pulsed fluorescence technique, using model TEI 43CTL until September 2010 and model TEI 43i-TLE after that. NO_x concentration was measured with TEI 42CTL (molybdenum converter) until February 2007, then with TEI 42CTL (photolytic converter) until April 2011, and after that with TEI 42iTL (photolytic converter).

Monoterpene concentration was measured with quadrupole Proton Transfer Reaction Mass Spectrometer (PTR-MS; Ionicon, Austria). We used concentration measured at 16.8 m height. The measurement setup is described in Taipale et al. (2008) and Rantala et al. (2014).

Equivalent black carbon (eBC) concentration was derived by filter-based optical instruments: Aethalometer (Magee Scientific, model AE31) in years 2006–2017 and Multi-Angle

Absorption Photometer (MAAP; Thermo Fisher Scientific, model 5012) in years 2013–2020. The correction procedure for AE31 data at SMEAR II as well as the measurement setup of AE31 and MAAP are described in detail in Luoma et al. (2021). The AE31 data were corrected by using a correction algorithm described in Virkkula et al. (2007) and using a multiple scattering correction factor of 3.14, which was derived by comparing the AE31 to MAAP. To derive the eBC concentration from the measured absorption coefficient, a mass absorption cross section values of 6.6 and 4.78 m² g⁻¹ were used for MAAP at wavelength 637 nm and AE31 at wavelength 880 nm, respectively.

Air mass origins were calculated using Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT) (Stein et al., 2015). The arrival height of the trajectories was 100 m, and they were calculated 96 h backwards in 1 h resolution. The trajectories were divided into three sectors as described in Rätty et al. (2023). Clean sector (Fig. S5e) represents area with minor anthropogenic contribution, while European and Eastern sectors represent more pollutant air mass source areas (Niemi et al., 2009; Riuttanen et al., 2013). Trajectory was classified into certain sector when it spent at least 90 % of the time in that sector, otherwise it was classified as mixed.

2.6 Correlations, bivariate fitting and long-term trend estimation

The Pearson's correlation coefficients between the mass concentrations from different instruments were calculated in Matlab, along with bivariate fitting (Cantrell, 2008). Before the analysis, we removed clear outliers that were further than 6 scaled median absolute deviations (MAD) away from the median using the Matlab built-in function *isoutlier*. The procedure was done for the whole dataset at once, i.e. without regarding for instance seasonal dynamics, but separately for each instrument and PM size. **The limit was determined using visual inspection.** About 1.5 % of the data were removed. When comparing DMPS+APS and SHARP with the impactor data, we calculated 2–3 days' cumulative aerosol mass concentration to make DMPS+APS and SHARP measurements comparable to the impactor data time resolution.

The statistical significance of long-term trends in linear scale were calculated using the *mannkendall* function for Matlab (v1.1.0, 10.5281/zenodo.4495589). We applied the seasonal 3PW method, which utilizes three pre-whitening methods for the trend estimation (Hirsch et

al., 1982). Pre-whitening methods by Kulkarni and von Storch (1995) and Yue et al. (2002) remove lag-1 autocorrelation and autocorrelation on detrended data, enabling to determine the statistical significance of Mann-Kendall test reliably; of these the one with higher value is reported. Variance-corrected trend-free pre-whitening method by Wang et al. (2015) is used for calculation of Sen's slope, which leads to more accurate trend analysis (Collaud Coen et al., 2020).

3 Results and discussion

3.1 Comparison between the mass measurement methods

Here, we present the comparison between the different aerosol mass measurement techniques at SMEAR II (Fig. 3 and S1a). We found that the data from the different mass measurement techniques correlate well, with the correlation coefficients $R > 0.8$ for all the measurements except between SHARP and impactor for which $R = 0.74$ (Table 1). Thereby, the correlation was lower between the two direct mass measurements, SHARP and impactor, than between DMPS+APS derived and impactor or SHARP measurements, even though with the DMPS+APS method we had to assume constant density and spherical shape of the particles in the mass concentration calculations. In reality, the particle composition, density, and shape vary between different particles (Kannosto et al., 2008; Heikkinen et al., 2020), which could potentially lead to the higher uncertainty in the indirect DMPS+APS mass calculations.

Table 1: Correlation coefficients between different PM measurement techniques. Correlation coefficient between SHARP and DMPS+APS in PM_{10} was 0.84. In all cases P -value < 0.05 .

Method	Impactor, PM_{10}	Impactor, $PM_{2.5}$	Impactor, PM_1
DMPS+APS	0.84	0.86	0.88
SHARP	0.74	-	-

To estimate the impact of selected density in DMPS+APS method, we calculated the average mass concentrations using 1.1 and 2.0 $g\ cm^{-3}$ as lower and upper estimates of the particle densities (Kannosto et al., 2008). The average PM mass concentrations for 2005–2020 are presented in Table S1. The average mass concentrations calculated with 1.1 $g\ cm^{-3}$ particle density were 7–12 % smaller compared to the mass obtained with 1.5 $g\ cm^{-3}$ density for

PM₁₀, 12–20 % for PM_{2.5}, and 20–25 % for PM₁. Correspondingly, with 2.0 g cm⁻³ particle density, the calculated mass concentrations were 14–17 % larger for PM₁₀, 19–27 % for PM_{2.5}, and 29–30 % for PM₁.




Figure 3: Correlation between the different mass measuring methods against impactor measurements a) PM₁₀ from SHARP, b) PM₁₀ from DMPS+APS, c) PM_{2.5} from DMPS+APS, and d) PM₁ from DMPS+APS. Bivariate fit to the data is represented with a blue line and 1:1 line is black. Color is PM₁ to PM₁₀ ratio from impactor measurements and in (a) markers differentiate the inlet heating temperature of SHARP (circle = 45 °C and plus sign = 35 °C). The data are averaged based on the impactor time resolution (2–3 days).

Comparing Fig. 3 and Fig. S1a, it seems that the data points between SHARP and DMPS+APS are positioned more distinctly on the 1:1 line whereas the impactor data are scattered more towards higher concentrations in all size classes. After the inlet heating temperature reduction in SHARP from 45 to 35 °C, the PM₁₀ values measured by SHARP were more comparable to those measured by impactor, except for the lowest and highest PM₁₀ concentrations (Fig. S2a–b). When comparing to DMPS+APS data (Fig. S2c–d),

SHARP also showed slightly lower PM₁₀ concentrations when the inlet was heated to 45 °C, and mostly similar concentrations (within standard deviation) when inlet was heated to 35 °C. Again, with the lowest concentrations, SHARP showed higher variability in the measured PM₁₀ concentrations. When excluding the lowest mass concentrations (approximately below 1.5 µg m⁻³), with 45 °C inlet heating SHARP to impactor ratio was 0.65 and SHARP to DMPS+APS ratio 0.85. With 35 °C inlet heating, the ratios were 0.91 and 1.0, respectively. This indicates that the higher inlet heating temperature might have led to 15–25 % losses of semi-volatile compounds from the sample air of SHARP.

Color in Fig. 3 and S1a is PM₁ to PM₁₀ ratio from impactor measurements. In general, the correlations between instruments were rather independent of the fractions of different particle sizes, but in PM₁₀ correlation with impactor (Fig. 3a–b), the scattered data points have lower PM₁ to PM₁₀ ratio. This implies that the impactor PM₁₀ measurements were likely overestimated in these cases since in PM₁ and PM_{2.5}, as well as in the DMPS+APS correlation with SHARP, PM₁ to PM₁₀ ratio from impactor data seem to be more evenly distributed (Fig. 3c-d and S1a). We additionally plotted correlation between monthly median concentration of PM₁ and PM_{2.5} as well as PM_{2.5} and PM₁₀ from DMPS+APS and impactor measurements (Fig. S1b). The figure shows that while the data from DMPS+APS is rather well aligned with 1:1 line, the PM₁₀ against PM_{2.5} from impactor data has more scattered data points, further implying that the impactor data might be overestimated.


In Waldén et al. (2010) different PM analyzers were tested for air quality monitoring in Helsinki. They reported that the two tested SHARP instruments passed the equivalence tests for PM₁₀ monitoring while for the PM_{2.5} measurements a calibration correction factor had to be applied. In their instruments, inlet lines were heated to 35 °C. They also reported that while Dekati PM10 impactor was overall indicative measurement method for PM_{2.5} (other sizes were not measured), it overestimated the concentrations compared to the reference methods. For the impactor, they used 24 h sampling period with 30 l min⁻¹ flow rate.

The measurement methods used in this study differ considerably from each other, and hence they are subject to different kinds of issues in PM monitoring. The impactor data, for example, is sensitive to any disturbances related to the weighing of the filters or evaporation of semi-volatile material from the filters during the long sampling time. The impactor is,

however, the only purely weighing-based mass measurement at SMEAR II. Thus, in the next section, we compare all the other methods against the impactor data.

3.2 PM concentrations, seasonal variation, and emission events

We explore the time series of PM concentrations to observe overall concentration levels, seasonal differences, and specific emission events (Fig. 4 and S2). Mean values from 1991–2002 reported by Laakso et al. (2003) are also included in the figures to compare the results with the earlier values from SMEAR II. To enable the comparison with the values by Laakso et al. (2003), we present mean and median concentrations of shorter, approximately five-year periods (2005–2010, 2011–2015, and 2016–2020) in Table 2.






Figure 4: Seasonal median (a) PM_{10} , (b) $PM_{2.5}$ and (c) PM_1 concentrations measured with the impactor as well as (d) temperature and their 25 and 75 quartile ranges at SMEAR II. The tick marks on the x-axis are in the beginning of a year. Mean values for 1991–2002 are from Laakso et al. (2003).

Table 2: Average (mean / median) PM concentrations measured at SMEAR II. Values (mean) for 1999–2001 are from Laakso et al. (2003) and other values from this work. First number in each cell is the mean for the indicated period and following numbers are seasonal means. Unit is $\mu\text{g m}^{-3}$.

	1999–2001	2005–2010	2011–2015	2016–2020	2005–2020
PM_{10} , impactor	6.9	5.4 / 4.4	5.8 / 4.8	4.4 / 3.4	4.5 / 4.2
-Spring	7.4	5.9 / 4.7	5.3 / 4.2	4.4 / 3.3	4.2 / 4.1
-Summer	7.2	6.4 / 5.6	7.4 / 6.4	5.8 / 5.5	5.5 / 5.6

-Autumn	6.9	4.5 / 3.4	5.4 / 4.4	4.1 / 3.0	4.3 / 3.5
-Winter	5.9	4.7 / 3.9	5.0 / 4.4	3.3 / 2.8	3.9 / 3.6
PM ₁₀ , SHARP	-	-	4.2 / 3.6	4.7 / 4.0	5.2 / 3.8
-Spring			4.0 / 3.5	4.4 / 3.6	5.2 / 3.6
-Summer			4.9 / 4.4	6.0 / 5.5	6.5 / 5.1
-Autumn			3.8 / 3.1	4.7 / 3.8	4.7 / 3.7
-Winter			4.1 / 3.3	3.8 / 3.3	4.4 / 3.3
PM ₁₀ , DMPS+APS	-	5.5 / 4.8	4.8 / 4.0	4.2 / 3.4	4.9 / 4.1
-Spring		5.8 / 4.9	4.4 / 3.9	4.1 / 3.5	4.8 / 4.1
-Summer		6.2 / 5.9	5.9 / 5.0	5.5 / 4.9	5.9 / 5.3
-Autumn		4.8 / 3.9	4.2 / 3.3	3.8 / 2.8	4.3 / 3.4
-Winter		5.5 / 4.7	4.7 / 3.9	3.5 / 2.9	4.7 / 3.9
PM _{2.5} , impactor	5.8	4.6 / 3.7	4.7 / 3.8	3.5 / 2.8	4.3 / 3.4
-Spring	6.4	5.0 / 4.1	4.2 / 3.5	3.4 / 2.6	4.2 / 3.3
-Summer	5.9	5.2 / 4.6	5.9 / 5.0	4.5 / 3.7	5.2 / 4.4
-Autumn	5.7	3.6 / 2.7	4.2 / 3.4	3.3 / 2.3	3.7 / 2.8
-Winter	5.1	4.4 / 3.5	4.5 / 3.8	2.9 / 2.4	4.0 / 3.2
PM _{2.5} , DMPS+APS	-	4.7 / 4.0	4.1 / 3.4	3.6 / 3.0	4.2 / 3.6
-Spring		4.8 / 4.2	3.7 / 3.2	3.4 / 2.9	4.1 / 3.4
-Summer		5.1 / 4.8	4.9 / 4.2	4.6 / 4.2	4.9 / 4.5
-Autumn		3.9 / 3.2	3.5 / 2.7	3.3 / 2.4	3.6 / 2.8
-Winter		5.1 / 4.3	4.3 / 3.6	3.3 / 2.8	4.3 / 3.6
PM ₁ , impactor	4.3	3.8 / 3.0	3.8 / 2.9	2.7 / 2.1	3.4 / 2.7
-Spring	4.4	4.2 / 3.4	3.3 / 2.7	2.6 / 2.0	3.4 / 2.7
-Summer	5.6	4.4 / 3.7	4.9 / 4.0	3.5 / 3.0	4.2 / 3.5
-Autumn	3.6	2.8 / 2.1	3.3 / 2.3	2.4 / 1.6	2.8 / 2.0
-Winter	3.8	3.7 / 2.8	3.7 / 3.0	2.3 / 1.7	3.3 / 2.5
PM ₁ , DMPS+APS	-	3.8 / 3.3	3.3 / 2.6	3.0 / 2.4	3.4 / 2.8
-Spring		3.9 / 3.3	2.9 / 2.5	2.8 / 2.3	3.3 / 2.6
-Summer		4.2 / 3.9	4.1 / 3.6	3.8 / 3.3	4.0 / 3.6
-Autumn		3.0 / 2.3	2.7 / 1.9	2.6 / 1.8	2.8 / 2.1
-Winter		4.3 / 3.4	3.5 / 2.8	2.7 / 2.0	3.5 / 2.8

387

388 The PM concentrations in all size classes are typically highest in summer and lowest in
389 autumn (Table 2). In summertime, the surrounding boreal forest is a large source of organic
390 compounds (Fig. S4c), which contribute to the aerosol load as shown already in several
391 studies (e.g. Heikkinen et al., 2020; 2021; Yli-Juuti et al., 2021). Due to the temperature
392 dependent activity of the forest, warm spells and heatwaves increase the VOC emissions,
393 such as in 2018 (Fig. S4c; Neefjes et al., 2022), which is also evident in the PM data in all

size classes (Fig. 4). Furthermore, pollen and other biological particles contribute especially to coarse mode particle mass at SMEAR II from late spring to early autumn (Manninen et al., 2014).

Although PM mass concentrations are generally decreasing (Fig. 4), certain events associated with higher PM levels, such as wildfires and volcanic eruptions, can be detected. In 2006 springtime as well as in 2006 and 2010 summer forest fires in eastern Europe increased the measured PM concentrations at SMEAR II (Fig. 4) as seen also in Anttila et al. (2008) and Leino et al. (2014). The growing seasons of 2006 and 2011 were exceptionally warm at SMEAR II based on the analysis spanning years 1996–2017 (Pysarenko et al., 2022), which is also visible in PM concentrations (Fig. 4), but the relatively high PM concentrations in spring 2010 and 2011 can also be caused by the plume of ash and SO₂ from the erupted Eyjafjallajökull (Thomas et al., 2011; Gudmundsson et al., 2012; Flanner et al., 2014) and Grímsvötn (Cooke et al., 2011; Tesche et al., 2012) volcanoes in Iceland. The considerably higher PM concentrations in autumn 2014 were affected by eruptions of Bardarbunga (Gislason et al., 2015) and Holuhraun (Ilyinskaya et al., 2017) volcanoes in Iceland, which Heikkinen et al. (2020) also noticed in the sulfate aerosol and SO₂ concentrations.

During the coldest winters 2009–2010 and 2010–2011, the measured PM concentrations were high (Fig. 4 and S2). These years were also associated with high concentrations of SO₂, NO_x, and eBC (Fig. S4). Residential heating is known to be a source of particulate emissions as wood is burned for heating (Spindler et al., 2004; Viana et al., 2008; Barmpadimos et al., 2011). However, the coldest winter temperatures are typically measured in Finland when air is transported from the eastern continental areas (Sui et al., 2020). These, and particularly southeastern, areas are also a source of atmospheric pollutants (Niemi et al., 2009; Riuttanen et al., 2013). Hence, rather than being local, pollutants can also be advected to Finland. Air mass source area analysis shows that winters with higher fraction of easterly air masses (Fig. S5) were colder and had also higher PM levels, although we acknowledge that this analysis does not reveal the actual source of the measured PM. Further, boundary layer height dynamics affect the measured concentrations, because shallow boundary layer heights during cold winter days can concentrate the anthropogenic pollutants close to the surface (Stull, 1988; Sinclair et al., 2022).

Overall, the air quality at SMEAR II was very good during our measurement period from 2005 to 2020. The average concentrations measured by impactor were 4.5, 4.3, and 3.4 $\mu\text{g m}^{-3}$ for PM_{10} , $\text{PM}_{2.5}$, and PM_1 , which are in line with $\text{PM}_{2.5}$ concentrations measured in 2002 and 2010 at background stations in Sweden and Norway (average $\text{PM}_{2.5}$ ranging from 4.3 to 9.9 $\mu\text{g m}^{-3}$ (Cusack et al., 2012)) and generally lower than at other background stations in Europe (average $\text{PM}_{2.5}$ ranging from 5.5 to 26.2 $\mu\text{g m}^{-3}$ (Cusack et al., 2012)).

3.3 Long-term trends

The measured PM concentrations show a declining trend during the measurement period from 2005 to 2020 (Fig. 4). Compared to the values reported by Laakso et al. (2003) for 1999–2001 (6.9, 5.8, and 4.3 $\mu\text{g m}^{-3}$ for PM_{10} , $\text{PM}_{2.5}$, and PM_1 , respectively), the values in 2016–2020 are almost 40 % lower in all size classes (Table 2).

While the overall PM concentrations are decreasing, the different mass measurement methods gave slightly inconsistent results: DMPS+APS method shows constant decline in PM concentrations, whereas the impactor data shows slight increase in all PM sizes for 2011–2015 period for all seasons but spring (Table 2). The clearest difference between the impactor and DMPS+APS data seems to be the steadier decline in autumn concentrations in DMPS+APS data (Fig. 4 and Fig. S2b-c). Hence, despite the discrepancies the methods give generally comparable results.

SHARP data shows increased PM_{10} concentration between 2011–2015 and 2016–2020 for all other seasons except for winter, but this is likely explained by the decreased inlet heating temperature, changed between the two periods (Fig. S3). Hence, no further conclusions of the trend in SHARP data can be drawn, even though generally the concentrations measured by SHARP follow the concentrations measured by DMPS+APS method (Fig. S2a-b).

Long-term trends are shown seasonally for each size class PM_{10} , $\text{PM}_{2.5}$, and PM_1 using impactor (Fig. 5, S6, and S7), and DMPS+ APS data (Fig. S8–S10). Calculated seasonal and annual trends are presented in Table 3. On seasonal scale, decreasing trends, ranging from -0.007 to -0.066 $\mu\text{g m}^{-3} \text{y}^{-1}$, are observed in each measured size class, while on annual scale, the trends vary between -0.021 and -0.036 $\mu\text{g m}^{-3} \text{y}^{-1}$. In general, the largest decreases in all size classes are observed in winter, whereas the decrease is the lowest in autumn. For the

impactor method, the decline is statistically significant at 95 % level in spring and winter, but not in summer and autumn. However, when calculating the trends from DMPS+APS data using 6-hour averages, the Mann-Kendall test revealed a statistically significant decrease in all size classes and seasons (Fig. S8–S10).




Figure 5: PM₁ concentration with the impactor method in (a) spring, (b) summer, (c) autumn, and (d) winter. Red horizontal line represents the median, the distance between the box edges shows the interquartile range, and whiskers extend to 1.5 times the interquartile range. Outliers are not shown. Slope represents trend calculated using Sen's slope and statistical significance is calculated using Mann-Kendall test. The trends were statistically significant in spring and winter, but not in summer and autumn.

The seasonal differences in PM trends follow the trends observed also in SO₂ and NO_x concentrations (Fig. S11 and S12), as well as in eBC (Fig. S4d-e). Further, Luoma et al. (2019) showed that the relative decline of light absorbing aerosol is faster than the light scattering aerosol at SMEAR II, and that the decline was strongest in spring and winter. Hence, the results imply that the decrease in anthropogenic pollutants drive the decrease in PM, which has been seen also elsewhere (e.g. Yttri et al., 2021). On the other hand, the lower

summer and autumn time decline can also be explained by the high fraction (more than 50 %) of OA from the surrounding boreal forest in the PM mass concentration at SMEAR II (Heikkinen et al., 2020). Further, in Li et al. (2023) the concentrations of organic precursors have even been shown to have an increasing trend at SMEAR II.

Table 3: Annual and seasonal trends in 2005–2020 PM concentrations at SMEAR II. Results from the DMPS+APS method are calculated from 6 h averages while impactor data has original time resolution (averaged over 2–3 days). Results, which are not statistically significant in 95 % level, are marked with *. Unit is $\mu\text{g m}^{-3} \text{y}^{-1}$ except the second number in annual trends which is $\% \text{y}^{-1}$.

	Method	Spring	Summer	Autumn	Winter	Annual
PM ₁₀	Impactor	-0.053	-0.018*	-0.017*	-0.056	-0.035 / -0.59
	DMPS+APS	-0.038	-0.012	-0.016	-0.066	-0.032 / -0.56
PM _{2.5}	Impactor	-0.059	-0.020*	-0.012*	-0.057	-0.034 / -0.70
	DMPS+APS	-0.034	-0.013	-0.009	-0.061	-0.028 / -0.56
PM ₁	Impactor	-0.064	-0.018*	-0.013*	-0.059	-0.036 / -0.89
	DMPS+APS	-0.027	-0.013	-0.007	-0.042	-0.021 / -0.52

In winter, biogenic OA precursors have minima in their concentrations (Fig. S4c), as shown also in Heikkinen et al. (2020), and consequently the collected PM originates mostly from anthropogenic sources, such as traffic, industry, and different combustion processes (Forsberg et al., 2005; Anttila & Tuovinen, 2010) as is indicated by the winter maxima in eBC concentrations (Fig. S4d-e). Moreover, many gaseous pollutants, emitted from anthropogenic processes and contributing to atmospheric chemistry or aerosol processes, such as SO₂ and NO_x, have maxima in their seasonal cycle in spring and winter (Fig. S4a-b; Lyubovtseva et al., 2005; Anttila & Tuovinen, 2010; Riuttanen et al., 2013; Heikkinen et al., 2020), further affirming the contribution of anthropogenic pollution to the observed trends. Additionally, Banerji et al. (2025) showed that at SMEAR II, light absorbing aerosol peak in winter, being thus associated with e.g. black carbon from anthropogenic activities, while aerosol scattering peaks in summer and winter, being thus likely associated with organic aerosol in summer and sulfates in winter. They also found an increasing trend in single scattering albedo, indicating that the relative proportion of light absorbing aerosol decrease. Luoma et al. (2019), in turn, reported decreased light scattering and absorption with a

simultaneous increase in light backscattering fraction and scattering Ångström exponent at SMEAR II, indicating reduction in large particle concentration.

The seasonal difference in PM sources is visible also in the ratios between PM_1 to $PM_{2.5}$ and $PM_{2.5}$ to PM_{10} plotted against temperature bins (Fig. 6) as well as in monthly PM_1 to PM_{10} ratio (Fig. S13). The fraction of smaller particles increases in cold and warm temperatures (Fig. 6), which could be attributed to anthropogenic pollution during winters and secondary aerosol formation in summer, which is also visible in aerosol PM_1 to PM_{10} aerosol light scattering coefficient (Luoma et al., 2019). In winters, nearly 80 % of PM_{10} consists actually of PM_1 (Fig. S13).

The PM_1 to $PM_{2.5}$ and $PM_{2.5}$ to PM_{10} ratios exhibit small, but statistically significant at the 95 % confidence level, negative trends (Fig. S14). The decline is particularly attributed to the decline in PM_1 concentration due to decreasing anthropogenic precursor concentrations since the it is larger in PM_1 to $PM_{2.5}$ ratio. Also, the annual relative trends from the impactor data are largest for PM_1 (Table 3). Sweden at two regional background sites, the $PM_{2.5}$ to PM_{10} ratios in 1999–2001 were 0.77 and 0.8 (Forsberg et al., 2005), which is in line with the measurements from SMEAR II. In 1999–2001 dataset, the $PM_{2.5}$ to PM_{10} ratio at SMEAR II was 0.86 in winter and spring, and 0.82 in summer and autumn (Laakso et al., 2003) while in 2020, the ratios fell below 0.8 (Fig. S14) also highlighting the change in aerosol population at SMEAR II.




Figure 6: PM₁ to PM_{2.5} and PM_{2.5} to PM₁₀ ratios in different temperature bins using impactor data. Horizontal line represents the median, the distance between the box edges shows the interquartile range, whiskers extend to 1.5 times the interquartile range, and data points even further from the median are presented with circles.

Generally, the PM concentrations have been observed to decrease in Europe (Spindler et al., 2004; Barmpadimos et al., 2011; Keuken et al., 2012; Guerreiro et al., 2014). However, in Guerreiro et al. (2014) non-significant positive trends in PM₁₀ (2002–2011) and PM_{2.5} (2006–2011) were also observed at Finnish rural background sites. In Anttila & Tuovinen (2010) both increasing and decreasing trends were detected in a dataset from 1994 to 2007 from Finland, which they linked to different measurement environments (urban, suburban, and industrial). Table 4 lists trends observed at different measurement sites across Europe.

Table 4: Summary of PM trends in previous literature. Unit is $\mu\text{g m}^{-3} \text{y}^{-1}$ except in Anttila & Tuovinen (2010) where the unit of the first number is $\mu\text{g m}^{-3} \text{month}^{-1}$ and the second is $\% \text{yr}^{-1}$. Note that the trends are calculated using different methods.

PM ₁₀	PM _{2.5}	Time	Location	Author
-0.15 to -1.2		1991–2008	Switzerland, various sites	Barmpadimos et al., (2011)
-0.4	-0.4	1998–2010	Europe, various sites	Barmpadimos et al., (2012)
	-1.9	2002–2010	Spain, regional background site	Cusack et al., (2012)
	-1.8	2002–2010	Europe, regional background sites	Cusack et al., (2012)
3.42 to -1.95	2.30 to -0.62	2002–2011 (PM ₁₀) 2006–2011 (PM _{2.5})	Europe, rural background sites	Guerreiro et al., (2014)
2.29 to -4.11	1.19 to -1.91	2006–2011	Europe, (sub)urban background sites	Guerreiro et al., (2014)
	-0.008 to -1.717	From 2005–2008 to 2015	Italy, Po Valley, various sites	Bigi & Ghermandi, (2016)
-0.13 to -2.83	-0.26 to -2.03	2004–2014	Spain, various sites	Pandolfi et al., (2016)

0.024 to -0.054 / -0.3 to -2.8	1994–2007	Finland, various sites	Anttila & Tuovinen, (2010)
--------------------------------------	-----------	---------------------------	----------------------------------

The trends measured at SMEAR II (Table 3) are similar than reported previously for Finland, but lower than the trends observed elsewhere in Europe (Table 4). One reason for this is likely the overall lower PM concentrations at SMEAR II compared to other locations as well as the longer timespan. Moreover, the trends in different studies have been calculated using different methods, such as Sen's slope (this study; Cusack et al., (2012); Guerreiro et al., (2014); Paldolfi et al., (2016)), and generalized least squares regression (Anttila & Tuovinen, 2010; Bigi & Ghermandi, (2016)), and also using different data preprocessing, which can further affect the trend estimates.

The connection between decreasing gaseous pollutant emissions and secondary aerosol concentrations has already been noted previously (Anttila & Tuovinen, 2010; Cusack et al., 2012; Kyrö et al., 2014; Pandolfi et al., 2016; Li et al., 2023) and decreasing PM trends in Europe have been connected to modernization of industry and heating systems as well as technology development of vehicles (Spindler et al., 2004; Barmpadimos et al., 2011; Keuken et al., 2012). Hence, the observed decrease in PM concentrations at SMEAR II is in line with previous studies and can be connected to the emission reductions driven by air quality legislation.

4 Conclusions

In this paper, different long-term aerosol mass concentration (PM_{10} , $PM_{2.5}$, PM_1) measurement techniques were compared and reported for the years 2005–2020 from SMEAR II, Finland. The direct mass concentration measurements with a cascade impactor and SHARP were compared with the mass concentrations calculated from the combined aerosol number size distributions of DMPS and APS. The results obtained using different methods are well comparable with the correlation coefficients of about 0.8.

The lower correlation values were connected to sampling methodologies: e.g., reducing the inlet heating temperature of SHARP increased the correlation with the impactor.

Additionally, although impactor measurements are simple and purely based on weighing of

filters, the impactor data showed somewhat higher concentrations than the other two methods, especially in the PM_{10} size, which might stem from the difficulties related to weighing masses down to micrograms. Any disturbances or deposited dust particles can lead to overestimated mass concentration, which might be the reason why impactor data showed statistically insignificant trends in summer and autumn while DMPS+APS data with similar absolute values resulted in statistically significant decreasing trend in PM concentration. On annual scale, both methods indicated statistically significant decreasing trends, which were comparable with the trends observed elsewhere in Finland.

The measured masses were similar between all the methods, and hence we can conclude that all methods were applicable for long-term PM monitoring. Yet, we acknowledge that the comparison of PM concentrations measured with different techniques gives valuable information for data quality control purposes, as well as for validating the applicability of the different methods. Therefore, we encourage conducting extensive comparisons with different methods at each measurement site.

The PM concentrations at SMEAR II were generally low, mostly less than $5 \mu g m^{-3}$, which clearly fell below the $20 \mu g m^{-3}$ limit by the EU air quality legislation. The highest PM concentrations at SMEAR II were measured in summer, when organic compounds from the surrounding boreal forest contribute to the measured PM mass. Peaks observed in the PM data can be related to transported particles from regions with e.g., forest fires or on-going volcanic eruptions.

The measurements showed overall decreasing PM trends for all size classes and in all seasons, although the decline was faster in PM_1 size class, which can be attributed to the decrease in anthropogenic pollution due to legislation aiming for improved air quality. Importantly, the trends were weakest in summer when natural emissions of VOCs from the forest lead to the formation of OA. As these natural VOCs are projected to increase with increasing temperature, it is possible that summertime OA concentrations keep increasing in the future. Taken together with the declining anthropogenic emissions, the role of natural aerosol particles could be anticipated to signify in the future. Overall, the results emphasize the importance of the long-term measurements (Kulmala et al., 2023) for understanding atmospheric aerosol mass concentrations and factors controlling them. This is a requirement

to quantify the relative roles of natural and anthropogenic sources to PM concentrations and ultimately to their impacts on health and climate.

Data availability

Aerosol data used in this study are available through EBAS database operated by NILU: <https://ebas-data.nilu.no/> (accessed 05 May 2025), BC data is available at SmartSMEAR database at <https://smear.avaa.csc.fi/> (accessed 05 September 2025), and temperature and trace gas data are published by Aalto et al. (2023) at <https://doi.org/10.23729/23dd00b2-b9d7-467a-9cee-b4a122486039> (accessed: 05 May 2025). Air mass source area data is available upon request from the corresponding author.

Author contribution

The idea and design of the study were conceived by HMK, MK, and TP. IY, HMK, and KLu wrote the manuscript, and together with LA analyzed the data and provided the visualizations. LH, TN, KLe, EE, MK, and TP helped to interpret the results. IY, HMK, LA, PA, JA, JL, KLu, SJ, and JK were also providing measurement data. All authors also contributed to reviewing and commenting on the manuscript.

Competing interests

MK is member of the editorial board of Aerosol Research.

Acknowledgements

The authors acknowledge the SMEAR II and INAR RI infrastructure as well as ACTRIS for the data accessed from SmartSMEAR (<https://smear.avaa.csc.fi/>) and EBAS (<https://ebas.nilu.no>) databases. We thank also thank SMEAR II personnel and PIs for the long-term measurement data.

Financial support

We acknowledge the following projects: ACCC Flagship funded by the Research Council of Finland grant number 337549 (UH), Academy professorship funded by the Research Council of Finland (grant no. 302958), Research Council of Finland projects no. 1325656, 311932, 334792, 316114, 325647, 325681, 347782, “Quantifying carbon sink, CarbonSink+ and their interaction with air quality” INAR project funded by Jane and Aatos Erkko Foundation, “Gigacity” project funded by Wihuri foundation, European Research Council (ERC) project

635 ATM-GTP Contract No. 742206, European Union via Non-CO2 Forcers and their Climate,
636 Weather, Air Quality and Health Impacts (FOCI) and Horizon 2020 research and innovation
637 program under grant agreements No. 654109 and 739530 (ACTRIS).

638 **References**

639 Aalto, J., Aalto, P., Keronen, P., Kolari, P., Rantala, P., Taipale, R., Kajos, M., Patokoski, J.,
640 Rinne, J., Ruuskanen, T., Leskinen, M., Laakso, H., Levula, J., Pohja, T., Siivola, E.,
641 Kulmala, M., and Ylivinkka, I.: SMEAR II Hyytiälä forest meteorology, greenhouse gases,
642 air quality and soil, University of Helsinki, Institute for Atmospheric and Earth System
643 Research, <https://doi.org/10.23729/23dd00b2-b9d7-467a-9cee-b4a122486039>, 2023.

644

645 Aalto, P., Hämeri, K., Becker, E., Weber, R., Salm, J., Mäkelä, J. M., Hoell, C., O'Dowd, C.
646 D., Hansson, H.-C., Väkevä, M., Koponen, I. K., Buzorius, G., and Kulmala, M.: Physical
647 characterization of aerosol particles during nucleation events, *Tellus B: Chem. and Phys.*
648 *Meteorol.*, 53, 344-358, DOI:10.1034/j.1600-0889.2001.d01-25.x, 2001.

649

650 Anttila, P., Makkonen, U., Hellén, H., Kyllönen, K., Leppänen, S., Saari, H., and Hakola, H.:
651 Impact of the open biomass fires in spring and summer of 2006 on the chemical composition
652 of background air in south-eastern Finland, *Atmospheric Environment*, 42, 6472–6486,
653 <https://doi.org/10.1016/j.atmosenv.2008.04.020>, 2008.

654

655 Anttila, P. and Tuovinen J.-P.: Trends of primary and secondary pollutant concentrations
656 in Finland in 1994–2007, *Atmospheric Environment*, 44, 30-41,
657 doi:10.1016/j.atmosenv.2009.09.041, 2010.

658

659 Anttila, P.: Air Quality Trends in Finland, 1994–2018, Academic dissertation in physics,
660 Institute of Atmospheric and Earth System Research / Physics, Faculty of Science, University
661 of Helsinki, Helsinki, Finland, Finnish Meteorological Institute Contributions No. 163,
662 [https://helda.helsinki.fi/bitstream/handle/10138/320460/thesis-Pia-](https://helda.helsinki.fi/bitstream/handle/10138/320460/thesis-Pia-Anttilaweb.pdf?sequence=1&isAllowed=y)
663 [Anttilaweb.pdf?sequence=1&isAllowed=y](https://helda.helsinki.fi/bitstream/handle/10138/320460/thesis-Pia-Anttilaweb.pdf?sequence=1&isAllowed=y), 2020.

664

665 Banerji, S., Luoma, K., Ylivinkka, I., Ahonen, L., Kerminen, V.-M., and Petäjä, T.:
 666 Measurement Report: Optical properties of supermicron aerosol particles in a boreal
 667 environment, EGU sphere [preprint], <https://doi.org/10.5194/egusphere-2025-1776>, 2025.
 668 Barmpadimos, I., Hueglin, C., Keller, J., Henne, S., and Prévôt, A. S. H.: Influence of
 669 meteorology on PM₁₀ trends and variability in Switzerland from 1991 to 2008, *Atmos.*
 670 *Chem. Phys.*, 11, 1813-1835, DOI:10.5194/acp-11-1813-2011, 2011.
 671
 672 Barmpadimos, I., Hueglin, C., Keller, J., Henne, S., and Prévôt, A. S. H.: Influence of
 673 meteorology on PM₁₀ trends and variability in Switzerland from 1991 to 2008, *Atmos. Chem.*
 674 *Phys.*, 11, 1813–1835, <https://doi.org/10.5194/acp-11-1813-2011>, 2011.
 675
 676 Barmpadimos, I., Keller, J., Oderbolz, D., Hueglin, C., and Prévôt, A. S. H.: One decade of
 677 parallel fine (PM_{2.5}) and coarse (PM₁₀–PM_{2.5}) particulate matter measurements in Europe:
 678 trends and variability, *Atmos. Chem. Phys.*, 12, 3189–3203, [https://doi.org/10.5194/acp-12-](https://doi.org/10.5194/acp-12-3189-2012)
 679 [3189-2012](https://doi.org/10.5194/acp-12-3189-2012), 2012.
 680
 681 Berner, A. and Luerzer, C: Mass size distributions of traffic aerosols at Vienna, *J. Phys.*
 682 *Chem.*, 84, 2079-2083, DOI:10.1021/j100453a016, 1980.
 683
 684 Bigi, A. and Ghermandi, G.: Trends and variability of atmospheric PM_{2.5} and PM₁₀–
 685 _{2.5} concentration in the Po Valley, Italy, *Atmos. Chem. Phys.*, 16, 15777–15788,
 686 <https://doi.org/10.5194/acp-16-15777-2016>, 2016.
 687
 688 Chen, Y., Zang, L., Du, W., Xu, D., Shen, G., Zhang, Q., Zou, Q., Chen, J., Zhao, M., and
 689 Yao, D.: Ambient air pollution of particles and gas pollutants, and the predicted health risks
 690 from long-term exposure to PM(2.5) in Zhejiang province, China, *Environ. Sci. Pollut. Res.*
 691 *Int.*, 25, 23833-23844, DOI:10.1007/s11356-018-2420-5, 2018.
 692
 693 Cantrell, C. A.: Technical Note: Review of methods for linear least-squares fitting of data and
 694 application to atmospheric chemistry problems, *Atmospheric Chemistry and Physics*, 8(17),
 695 5477–5487, DOI:10.5194/acpd-8-6409-2008, 2008.
 696
 697 Collaud Coen, M., Andrews, E., Bigi, A., Martucci, G., Romanens, G., Vogt, F. P. A., and
 698 Vuilleumier, L.: Effects of the prewhitening method, the time granularity, and the time

699 segmentation on the Mann–Kendall trend detection and the associated Sen's slope, *Atmos.*
700 *Meas. Tech.*, 13, 6945–6964, <https://doi.org/10.5194/amt-13-6945-2020>, 2020.

701

702 Cooke, M.C., Francis, P.N., Millington, S., Saunders, R. and Witham, C.: Detection of the
703 Grímsvötn 2011 volcanic eruption plumes using infrared satellite measurements, *Atmos. Sci.*
704 *Lett.*, 15, 321–327, <https://doi.org/10.1002/asl2.506>, 2011.

705

706 Cusack, M., Alastuey, A., Pérez, N., Pey, J., and Querol, X.: Trends of particulate matter
707 (PM_{2.5}) and chemical composition at a regional background site in the Western
708 Mediterranean over the last nine years (2002–2010), *Atmos. Chem. Phys.*, 12, 8341–8357,
709 <https://doi.org/10.5194/acp-12-8341-2012>, 2012.

710

711 Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D.,
712 Drewnick, F., Hings, S., Jung, D., Borrmann, S., and Andreae, M. O.: Size Matters More
713 Than Chemistry for Cloud-Nucleating Ability of Aerosol Particles, *Science*, 312, 1375–1378,
714 DOI:10.1126/science.1125261, 2006.

715

716 Flanner, M. G., Gardner, A. S., Eckhardt, S., Stohl, A., and Perket J.: Aerosol radiative
717 forcing from the 2010 Eyjafjallajökull volcanic eruptions, *J. Geophys. Res. Atmos.*, 119,
718 9481–9491, doi:10.1002/2014JD021977, 2014.

719

720 Forsberg, B., Hansson, H.-C., Johansson, C., Areskoug, H., Söderlund, K., and Järholm, B.:
721 Comparative Health Impact Assessment of Local and Regional Particulate Air Pollutants in
722 Scandinavia, *Ambio*, 34, 11–19, 2005.

723

724 Gislason, S., Stefansdottir, G., Pfeffer, M. A., Barsotti, S., Jóhannsson, T., Galeczka, I., Bali,
725 E., Sigmarsson, O., Stefansson, A., Keller, N., Sigurdsson, A., Bergsson, B., Galle, B.,
726 Jacobo, V. C., Arellano, S., Aiuppa, A., Jonasdottir, E., Eiríksdottir, E., Jakobsson, S., and
727 Gudmundsson, M.: Environmental pressure from the 2014–15 eruption of Bárðarbunga
728 volcano, Iceland, *Geochem. Perspect. Lett.*, 1, 84–93, DOI:10.7185/geochemlet.1509, 2015.

729

730 Goohs, K., Lilienfeld, P., and Wilbertz, J.: A Synchronized Hybrid Real-Time Particulate
731 Monitor, EAC 2009 proceedings, 2009.

732

733 Gudmundsson, M. T., Thordarson, T., Höskuldsson, A., Larsen, G., Björnsson, H., Prata, F.
 734 J., Oddsson, B., Magnússon, E., Högnadóttir, T., Petersen, G. N., Hayward, C. L., Stevenson,
 735 J. A., and Jónsdóttir, I.: Ash generation and distribution from the April-May 2010 eruption of
 736 Eyjafjallajökull, Iceland, Scientific reports, 2, 572, <https://doi.org/10.1038/srep00572>, 2012.
 737
 738 Guenther, A. B., Zimmerman, P. R., Harley, P. C., Monson, R. K., and Fall, R.: Isoprene and
 739 monoterpene emission rate variability: Model evaluations and sensitivity analyses, Journal of
 740 Geophys. Res.: Atmos., 98, 12609-12617, DOI:10.1029/93JD00527, 1993.
 741
 742 Guerreiro, C.B.B., Foltescu, V., and de Leeuw, F.: Air quality status and trends in Europe,
 743 Atmospheric Environment, 98, 376-384, 2014.
 744
 745 Hari, P. and Kulmala, M.: Station for Measuring Ecosystem-Atmosphere Relations: (SMEAR
 746 II), Boreal Environ. Res., 10, 315-322, DOI:10.1007/978-94-007-5603-8_9, 2005.
 747
 748 Heikkinen, L., Äijälä, M., Riva, M., Luoma, K., Dällenbach, K., Aalto, J., Aalto, P., Aliaga,
 749 D., Aurela, M., Keskinen, H., Makkonen, U., Rantala, P., Kulmala, M., Petäjä, T., Worsnop,
 750 D., and Ehn, M.: Long-term sub-micrometer aerosol chemical composition in the boreal
 751 forest: inter- and intra-annual variability, Atmos. Chem. Phys., 20, 3151-3180,
 752 DOI:10.5194/acp-20-3151-2020, 2020.
 753
 754 Heikkinen, L., Äijälä, M., Daellenbach, K. R., Chen, G., Garmash, O., Aliaga, D., Graeffe,
 755 F., Rätty, M., Luoma, K., Aalto, P., Kulmala, M., Petäjä, T., Worsnop, D., and Ehn, M.: Eight
 756 years of sub-micrometre organic aerosol composition data from the boreal forest
 757 characterized using a machine-learning approach, Atmos. Chem. Phys., 21, 10081–10109,
 758 <https://doi.org/10.5194/acp-21-10081-2021>, 2021.
 759
 760 Hirsch, R. M., Slack, J. R., and Smith, R. A.: Techniques of trend analysis for monthly water
 761 quality data, Water Resour. Res., 18, 107–121, DOI:10.1029/WR018i001p00107, 1982.
 762
 763 Ilyinskaya, E., Schmidt, A., Mather, T. A., Pope, F. D., Witham, C., Baxter, P., Jóhannsson,
 764 T., Pfeffer, M., Barsotti, S., Singh, A., Sanderson, P., Bergsson, B., McCormick Kilbride, B.,
 765 Donovan, A., Peters, N., Oppenheimer, C., and Edmonds, M.: Understanding the
 766 environmental impacts of large fissure eruptions: Aerosol and gas emissions from the 2014–

767 2015 Holuhraun eruption (Iceland), *Earth Planet. Sci. Lett.*, 472, 309-322,
 768 DOI:10.1016/j.epsl.2017.05.025, 2017.
 769
 770 IPCC (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working*
 771 *Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.*
 772 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2021.
 773
 774 Kannosto, J., Virtanen, A., Lemmetty, M., Mäkelä, J. M., Keskinen, J., Junninen, H.,
 775 Hussein, T., Aalto, P., and Kulmala, M.: Mode resolved density of atmospheric aerosol
 776 particles, *Atmos. Chem. Phys.*, 8, 5327-5337, DOI:10.5194/acp-8-5327-2008, 2008.
 777
 778 Keskinen, H.-M., Ylivinkka, I., Heikkinen, L., Aalto, P. P., Nieminen, T., Lehtipalo, K.,
 779 Aalto, J., Levula, J., Kesti, J., Ahonen, L. R., Ezhova, E., Kulmala, M., and Petäjä, T.: Long-
 780 term aerosol mass concentrations in southern Finland: instrument validation, seasonal
 781 variation and trends, *Atmos. Meas. Tech. Discuss.* [preprint], [https://doi.org/10.5194/amt-](https://doi.org/10.5194/amt-2020-447)
 782 2020-447, 2020.
 783
 784 Keuken, M.P., Roemer, M.G.M., Zandveld, P., Verbeek, R.P., and Velders, G.J.M.: Trends in
 785 primary NO₂ and exhaust PM emissions from road traffic for the period 2000-2020 and
 786 implications for air quality and health in the Netherlands, *Atmospheric Environment*, 54,
 787 313-319, 2012.
 788
 789 Kulkarni, A., and von Storch, H.: Monte Carlo Experiments on the Effect of Serial
 790 Correlation on the Mann-Kendall Test of Trend Monte Carlo experiments on the effect,
 791 *Meteorologische Zeitschrift*, 82–85, DOI:10.1127/METZ/4/1992/82, 1995.
 792
 793 Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T.,
 794 Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E.,
 795 Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P., Mikkilä, J., Vanhanen, J.,
 796 Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., Duplissy, J.,
 797 Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M. V., Smith, J.
 798 N., Ehn, M., Mentel, T. F., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., and
 799 Worsnop, D. R.: Direct Observations of Atmospheric Aerosol Nucleation, *Science*, 339, 943-
 800 946, DOI: 10.1126/science.1227385, 2013.

801
802 Kulmala, M., Lintunen, A., Lappalainen, H., Virtanen, A., Yan, C., Ezhova, E., Nieminen, T.,
803 Riipinen, I., Makkonen, R., Tamminen, J., Sundström, A.-M., Arola, A., Hansel, A.,
804 Lehtinen, K., Vesala, T., Petäjä, T., Bäck, J., Kokkonen, T., and Kerminen, V.-M.: Opinion:
805 The strength of long-term comprehensive observations to meet multiple grand challenges in
806 different environments and in the atmosphere, *Atmos. Chem. Phys.*, 23, 14949–14971,
807 <https://doi.org/10.5194/acp-23-14949-2023>, 2023.

808
809 Kyrö, E.-M., Väänänen, R., Kerminen, V.-M., Virkkula, A., Petäjä, T., Asmi, A., Dal Maso,
810 M., Nieminen, T., Juhola, S., Shcherbinin, A., Riipinen, I., Lehtipalo, K., Keronen, P., Aalto,
811 P. P., Hari, P., and Kulmala, M.: Trends in new particle formation in eastern Lapland,
812 Finland: effect of decreasing sulfur emissions from Kola Peninsula, *Atmos. Chem. Phys.*,
813 14(9), 4383–4396, 2014.

814
815 Laakso, L., Hussein, T., Aarnio, P., Komppula, M., Hiltunen, V., Viisanen, Y., and Kulmala,
816 M.: Diurnal and annual characteristics of particle mass and number concentrations in urban,
817 rural and Arctic environments in Finland, *Atmos. Environ.*, 37, 2629–2641, 2003.

818
819 Laakso, L., Laakso, H., Aalto, P. P., Keronen, P., Petäjä, T., Nieminen, T., Pohja, T., Siivola,
820 E., Kulmala, M., Kgabi, N., Molefe, M., Mabaso, D., Phalatse, D., Pienaar, K., and
821 Kerminen, V. M.: Basic characteristics of atmospheric particles, trace gases and meteorology
822 in a relatively clean Southern African Savannah environment, *Atmos. Chem. Phys.*, 8, 4823–
823 4839, DOI:10.1016/S1352-2310(03)00206-1, 2008.

824
825 Laj, P., Lund Myhre, C., Riffault, V., Amiridis, V., Fuchs, H., Eleftheriadis, K., Petäjä, T.,
826 Salameh, T., Kivekäs, N., Juurola, E., Saponaro, G., Philippin, S., Cornacchia, C., Alados
827 Arboledas, L., Baars, H., Claude, A., De Mazière, M., Dils, B., Dufresne, M., Evangeliou, N.,
828 Favez, O., Fiebig, M., Haeffelin, M., Herrmann, H., Höhler, K., Illmann, N., Kreuter, A.,
829 Ludewig, E., Marinou, E., Möhler, O., Mona, L., Eder Murberg, L., Nicolae, D., Novelli, A.,
830 O'Connor, E., Ohneiser, K., Petracca Altieri, R. M., Picquet-Varrault, B., van Pinxteren, D.,
831 Pospichal, B., Putaud, J., Reimann, S., Siomos, N., Stachlewska, I., Tillmann, R., Voudouri,
832 K. A., Wandinger, U., Wiedensohler, A., Apituley, A., Comerón, A., Gysel-Beer, M.,
833 Mihalopoulos, N., Nikolova, N., Pietruczuk, A., Sauvage, S., Sciare, J., Skov, H., Svendby,
834 T., Swietlicki, E., Tonev, D., Vaughan, G., Zdimal, V., Baltensperger, U., Doussin, J.,

Kulmala, M., Pappalardo, G., Sorvari Sundet, S., and Vana, M.: Aerosol, Clouds and Trace Gases Research Infrastructure (ACTRIS): The European Research Infrastructure Supporting Atmospheric Science, *Bulletin of the American Meteorological Society*, 105(7), E1098-E1136, <https://doi.org/10.1175/BAMS-D-23-0064.1>, 2024.

Leino, K., Riuttanen, L., Nieminen, T., Dal Maso, M., Väänänen, R., Pohja, T., Keronen, P., Järvi, L., Aalto, P., Virkkula, A., Kerminen, V.-M., Petäjä, T., and Kulmala, M.: Biomass-burning smoke episodes in Finland from eastern European wildfires, *Boreal Environ. Res.*, 19, 275-292, <http://www.borenv.net/BER/pdfs/ber19/ber19B-275.pdf>, 2014.

Li X., Li H., Yao L., Stolzenburg D., Sarnela N., Vettikkat L., de Jonge R.W., Baalbaki R., Uusitalo H., Kontkanen J., Lehtipalo K., Daellenback K.R., Jokinen T., Aalto J., Keronen P., Schobesberger S., Nieminen T., Petäjä T., Kerminen V.-M., Bianchi F., Kulmala M., and Dada L.: Over 20 years of observations in the boreal forest reveal a decreasing trend of atmospheric new particle formation. *Boreal Env. Res.* 29, 25–52, 2024.

Luoma, K., Virkkula, A., Aalto, P., Petäjä, T., and Kulmala, M.: Over a 10-year record of aerosol optical properties at SMEAR II, *Atmos. Chem. Phys.*, 19, 11363–11382, <https://doi.org/10.5194/acp-19-11363-2019>, 2019.

Luoma, K., Virkkula, A., Aalto, P., Lehtipalo, K., Petäjä, T., and Kulmala, M.: Effects of different correction algorithms on absorption coefficient – a comparison of three optical absorption photometers at a boreal forest site, *Atmos. Meas. Tech.*, 14, 6419–6441, <https://doi.org/10.5194/amt-14-6419-2021>, 2021.

Lyubovtseva, Y. S., Sogacheva, L., Dal Maso, M., Bonn, B., Keronen, P., and Kulmala, M.: Seasonal variations of trace gases, meteorological parameters, and formation of aerosols in boreal forests. *Boreal Env. Res.* 10, 493–510, 2005.

Manavi, S.E.I., Aktypis, A., Siouti, E., Skyllakou, K., Myriokefalitakis, S., Kanakidou, M., Pandis, S.N.: Atmospheric aerosol spatial variability: Impacts on air quality and climate change, *One Earth* 8, 101237, <https://doi.org/10.1016/j.oneear.2025.101237>, 2025.

868 Manninen, H., Bäck, J., Sihto-Nissilä, S.-L., Huffman, J., Pessi, A.-M., Hiltunen, V., Aalto,
869 P. P., Hidalgo, P., Hari, P., Saarto, A., Kulmala, M., and Petäjä, T.: Patterns in airborne
870 pollen and other primary biological aerosol particles (PBAP), and their contribution to
871 aerosol mass and number in a boreal forest, *Boreal Env. Res.*, 19 (suppl. B), 383-405, 2014.
872

873 Makkonen, U., Virkkula, A., Hellén, H., Hemmilä, M., Sund, J., Äijälä, M., Ehn, M.,
874 Junninen, H., Keronen, P., Petäjä, T., Worsnop, D. R., Kulmala, M., and Hakola, H.: Semi-
875 continuous gas and inorganic aerosol measurements at a boreal forest site: seasonal and
876 diurnal cycles of NH₃, HONO and HNO₃, *Boreal Env. Res.*, 19 (suppl. B), 311-328, 2014.
877

878 Maynard, A. D., and Kuempel, E. D. Airborne Nanostructured Particles and Occupational
879 Health. *J Nanopart Res* 7, 587–614, <https://doi.org/10.1007/s11051-005-6770-9>, 2005.
880

881 Neefjes I., Laapas M., Liu Y., Médus E., Miettunen E., Ahonen L., Quéléver L., Aalto J.,
882 Bäck J., Kerminen V-M., Lamplähti J., Luoma K., Mäki M., Mammarella I., Petäjä T., Rätty
883 M., Sarnela N., Ylivinkka I., Hakala S., Kulmala M., Nieminen T., and Lintunen A.: 25 years
884 of atmospheric and ecosystem measurements in a boreal forest — Seasonal variation and
885 responses to warm and dry years, *Boreal Env. Res.* 27, 1–31, 2022.
886

887 Neusüß, C., Weise, D., Birmili, W., Wex, H., Wiedensohler, A., and Covert, D. S.: Size-
888 segregated chemical, gravimetric and number distribution-derived mass closure of the aerosol
889 in Sagres, Portugal during ACE-2, *Tellus B: Chem. Phys. Meteorol.*, 52, 169-184,
890 DOI:10.1034/j.1600-0889.2000.00039.x, 2000.
891

892 Niemi, J.V., Saarikoski, S., Aurela, M., Tervahattu, H., Hillamo, R., Westphal, D.L., Aarnio,
893 P., Koskentalo, T., Makkonen, U., Vehkamäki, H., and Kulmala, M.: Long-range transport
894 episodes of fine particles in southern Finland during 1999–2007, *Atmospheric Environment*,
895 43, 1255–1264, <https://doi.org/10.1016/j.atmosenv.2008.11.022>, 2009.
896

897 Occhipinti, L. G., and Oluwasanya, P. W.: Particulate Matter Monitoring: Past, Present and
898 Future, *Int. J. Earth Environ. Sci.*, 2, 144, doi: <https://doi.org/10.15344/2456-351X/2017/144>,
899 2017.
900

Pandolfi, M., Alastuey, A., Pérez, N., Reche, C., Castro, I., Shatalov, V., and Querol, X.: Trends analysis of PM source contributions and chemical tracers in NE Spain during 2004–2014: a multi-exponential approach, *Atmos. Chem. Phys.*, 16, 11787–11805, <https://doi.org/10.5194/acp-16-11787-2016>, 2016.

Peters, T., Ott, D., and O'Shaughnessy, P.: Comparison of the GRIMM 1.108 and 1.109 Portable Aerosol Spectrometer to the TSI 3321 Aerodynamic Particle Sizer for Dry Particles, *Ann. Occup. Hyg.*, 50, 843–850, DOI:10.1093/annhyg/mel067, 2006.

Pope, C., Burnett, R. T., and Thun, M. J.: Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *J. Amer. Med. Assoc.*, 287, 1132–1141, doi: 10.1001/jama.287.9.1132, 2003.

Pysarenko, L., Krakovska, S., Savenets, M., Ezhova, E., Lintunen, A., Petäjä, T., Bäck, J. and Kulmala, M.: Two-decade variability of climatic factors and its effect on the link between photosynthesis and meteorological parameters: example of Finland's boreal forest, *Boreal Environment Research*, 27, 131–144, 2022.

Pöschl, U.: Atmospheric Aerosols: Composition, Transformation, Climate and Health Effects, *Angewandte Chemie International Edition*, 44, 7520–7540, <https://doi.org/10.1002/anie.200501122>, 2005.

Rantala, P., Taipale, R., Aalto, J., Kajos, M. K., Patokoski, J., Ruuskanen, T. M., and Rinne, J.: Continuous flux measurements of VOCs using PTR-MS — reliability and feasibility of disjunct-eddy-covariance, surface-layer-gradient, and surface-layer-profile methods. *Boreal Env. Res.*, 19 (suppl. B), 87–107, 2014.

Rinne, J., Ruuskanen, T., Reissell, A., Taipale, R., Hakola, H., and Kulmala, M.: On-line PTR-MS measurements of atmospheric concentrations of volatile organic compounds in a European boreal forest ecosystem, *Boreal Environ. Res.*, 10, 425–436, <http://www.borenv.net/BER/archive/pdfs/ber10/ber10-425.pdf>, 2005.

Riuttanen, L., Hulkkonen, M., Dal Maso, M., Junninen, H., and Kulmala, M.: Trajectory analysis of atmospheric transport of fine particles, SO₂, NO_x and O₃ to the SMEAR II station

935 in Finland in 1996–2008, *Atmos. Chem. Phys.*, 13, 2153–2164, [https://doi.org/10.5194/acp-](https://doi.org/10.5194/acp-13-2153-2013)
936 [13-2153-2013](https://doi.org/10.5194/acp-13-2153-2013), 2013.

937

938 Saarikoski, S., Mäkelä, T., Hillamo, R., Aalto, P., Kerminen, V.-M., and Kulmala, M.:
939 Physicochemical characterization and mass closure of size-segregated atmospheric aerosols
940 in Hyytiälä, Finland, *Boreal Environ. Res.*, 10, 386–400, 2005.

941

942 Schraufnagel, D.E.: The health effects of ultrafine particles, *Exp. Mol. Med.*, 52, 311–317,
943 <https://doi.org/10.1038/s12276-020-0403-3>, 2020.

944

945 Seinfeld, J. and Pandis, S.: *Atmospheric chemistry and physics: from air pollution to climate*
946 *change*. Second edition. John Wiley & Sons, New Jersey, USA, 2006.

947

948 Shiraiwa, M., Ueda, K., Pozzer, A., Lammel, G., Kampf, C. J., Fushimi, A., Enami, S.,
949 Arangio, A. M., Fröhlich-Nowoisky, J., Fujitani, Y., Furuyama, A., Lakey, P. S. J., Lelieveld,
950 J., Lucas, K., Morino, Y., Pöschl, U., Takahama, S., Takami, A., Tong, H., Weber, B.,
951 Yoshino, A., and Sato, K.: Aerosol Health Effects from Molecular to Global Scales,
952 *Environmental Science & Technology*, 51(23), 13545–13567, DOI: 10.1021/acs.est.7b04417,
953 2017.

954

955 Shukla, K., and Aggarwal, S.G.: A Technical Overview on Beta-Attenuation Method for the
956 Monitoring of Particulate Matter in Ambient Air, *Aerosol Air Qual. Res.*, 22, 220195,
957 <https://doi.org/10.4209/aaqr.220195>, 2022.

958

959 Sicard, P., Agathokleous, E., De Marco, A., Paoletti, E., and Calatayud, V.: Urban population
960 exposure to air pollution in Europe over the last decades, *Environ. Sci. Eur.*, 33, 28,
961 <https://doi.org/10.1186/s12302-020-00450-2>, 2021.

962

963 Sinclair, V. A., Ritvanen, J., Urbancic, G., Statnaia, I., Batrak, Y., Moisseev, D., and Kurppa,
964 M.: Boundary-layer height and surface stability at Hyytiälä, Finland, in ERA5 and
965 observations, *Atmos. Meas. Tech.*, 15, 3075–3103, [https://doi.org/10.5194/amt-15-3075-](https://doi.org/10.5194/amt-15-3075-2022)
966 2022, 2022.

967

968 Spindler, G., Müller, K., Brüggemann, E., Gnauk, T., and Herrmann, H.: Long-term size-
 969 segregated characterization of PM₁₀, PM_{2.5}, and PM₁ at the IfT research station Melpitz
 970 downwind of Leipzig (Germany) using high and low-volume filter samplers, *Atmos.*
 971 *Environ.*, 38, 5333–5347, DOI:10.1016/j.atmosenv.2003.12.047, 2004.
 972
 973 Stull, R. B.: *An Introduction to Boundary Layer Meteorology*, Springer Dordrecht, 1988.
 974
 975 Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.:
 976 NOAA's Hysplit Atmospheric Transport and Dispersion Modeling System, *B. Am. Meteorol.*
 977 *Soc.*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
 978
 979 Sui, C., Yu, L., and Vihma, T.: Occurrence and drivers of wintertime temperature extremes in
 980 Northern Europe during 1979–2016. *Tellus A: Dynamic Meteorology and Oceanography*,
 981 72(1), 1–19, <https://doi.org/10.1080/16000870.2020.1788368>, 2020.
 982
 983 Taipale, R., Ruuskanen, T. M., Rinne, J., Kajos, M. K., Hakola, H., Pohja, T., and Kulmala,
 984 M.: Technical Note: Quantitative long-term measurements of VOC concentrations by PTR-
 985 MS – measurement, calibration, and volume mixing ratio calculation methods, *Atmos. Chem.*
 986 *Phys.*, 8, 6681–6698, <https://doi.org/10.5194/acp-8-6681-2008>, 2008.
 987
 988 Tesche, M., Glantz, P., Johansson, C., Norman, M., Hiebsch, A., Ansmann, A., Althausen,
 989 D., Engelmann, R., and Seifert, P.: Volcanic ash over Scandinavia originating from the
 990 Grímsvötn eruptions in May 2011, *J. Geophys. Res.*, 117, D09201,
 991 doi:10.1029/2011JD017090, 2012.
 992
 993 Thomas, H. E., and Prata, A. J.: Sulphur dioxide as a volcanic ash proxy during the April–
 994 May 2010 eruption of Eyjafjallajökull Volcano, Iceland, *Atmos. Chem. Phys.*, 11, 6871–
 995 6880, <https://doi.org/10.5194/acp-11-6871-2011>, 2011.
 996
 997 Van Dingenen, R., Raes, F., Putaud, J.-P., Baltensperger, U., Charron, A., Facchini, M.,
 998 Decesari, S., Sandro, F., Gehrig, R., Hansson, H.-C., Harrison, R., Hüglin, C., Jones, A., Laj,
 999 P., Lorbeer, G., Maenhaut, W., Palmgren, F., Querol, X., Rodriguez, S., and Wåhlin, P.: A
 1000 European aerosol phenomenology - 1: Physical characteristics of particulate matter at

1001 kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2561-2577,
 1002 DOI:10.1016/j.atmosenv.2004.01.040, 2004.
 1003
 1004 Viana, M., Kuhlbusch, T.A.J., Querol, X., Alastuey, A., Harrison, R.M., Hopke, P.K.,
 1005 Winiwarter, W., Vallius, M., Szidat, S., Prévôt, A.S.H., Hueglin, C., Bloemen, H., Wåhlin,
 1006 P., Vecchi, R., Miranda, A.I., Kasper-Giebl, A., Maenhaut, W., and Hitzenberger, R.: Source
 1007 apportionment of particulate matter in Europe: A review of methods and results, *Journal of*
 1008 *Aerosol Science*, 39, 10, 827-849, <https://doi.org/10.1016/j.jaerosci.2008.05.007>, 2008.
 1009
 1010 Virkkula, A., Mäkelä, T., Hillamo, R., Yli-Tuomi, T., Hirsikko, A., Hämeri, K., and
 1011 Koponen, I. K.: A Simple Procedure for Correcting Loading Effects of Aethalometer
 1012 Data, *Journal of the Air & Waste Management Association*, 57, 10, 1214–1222,
 1013 <https://doi.org/10.3155/1047-3289.57.10.1214>, 2007.
 1014
 1015 Waldén, J., Hillamo, R., Aurela, M., Mäkelä, T., and Laurila, S.: Demonstration of the
 1016 equivalence of PM_{2.5} and PM₁₀ measurement methods in Helsinki 2007-2008, *Studies No.*
 1017 *3*, Finnish Meteorological Institute, Helsinki, Finland, ISSN 1796-1203, 2010.
 1018
 1019 Wang, W., Chen, Y., Becker, S., and Liu, B.: Linear trend detection in serially dependent
 1020 hydrometeorological data based on a variance correction Spearman rho method, *Water*, 7(12),
 1021 7045–7065, DOI:10.3390/w7126673, 2015.
 1022
 1023 Wang, J., Xing, J., Mathur, R., Pleim, J. E., Wang, S., Hogrefe, C., Gan, J.-M., Wong, D. C.,
 1024 and Hao, J.: Historical Trends in PM_{2.5}-Related Premature Mortality during 1990–2010
 1025 across the Northern Hemisphere, *Environmental Health Perspectives* 125, 3,
 1026 <https://doi.org/10.1289/EHP298>, 2017.
 1027
 1028 WHO global air quality guidelines: Particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen
 1029 dioxide, sulfur dioxide and carbon monoxide, Geneva: World Health Organization, 2021.
 1030
 1031 Yli-Juuti, T., Mielonen, T., Heikkinen, L., Arola, A., Ehn, M., Isokääntä, S., Keskinen, H.-
 1032 M., Kulmala, M., Laakso, A., Lipponen, A., Luoma, K., Mikkonen, S., Nieminen, T.,
 1033 Paasonen, P., Petäjä, T., Romakkaniemi, S., Tonttila, J., Kokkola, H., and Virtanen, A.:

1034 Significance of the organic aerosol driven climate feedback in the boreal area, *Nature*
 1035 *Communications*, 12(1), 1–9, 2021.

1036

1037 Yttri, K. E., Canonaco, F., Eckhardt, S., Evangeliou, N., Fiebig, M., Gundersen, H.,
 1038 Hjellbrekke, A.-G., Lund Myhre, C., Platt, S. M., Prévôt, A. S. H., Simpson, D., Solberg, S.,
 1039 Surratt, J., Tørseth, K., Uggerud, H., Vadset, M., Wan, X., and Aas, W.: Trends, composition,
 1040 and sources of carbonaceous aerosol at the Birkenes Observatory, northern Europe, 2001–
 1041 2018, *Atmos. Chem. Phys.*, 21, 7149–7170, <https://doi.org/10.5194/acp-21-7149-2021>, 2021.

1042

1043 Yue, S., Pilon, P., Phinney, B., and Cavadias, G.: The influence of autocorrelation on the
 1044 ability to detect trend in hydrological series, *Hydrol. Process.*, 16(9), 1807–1829,
 1045 <https://doi.org/10.1002/hyp.1095>, 2002.