

1 UFP-, BC-, and PM_{2.5}- measurements and effect of dispersion 2 conditions on concentration levels in a residential area affected by 3 wood-smoke pollution from domestic heating during the winter 4 months

5 Peter Bächler¹, Frederik Weis², Sebastian Kohler², Achim Dittler¹

6 ¹Institute of Mechanical Process Engineering and Mechanics, Karlsruhe Institute of Technology, Karlsruhe, 76131, Germany

7 ²Palas GmbH, Karlsruhe, 76187, Germany

8 *Correspondence to:* Peter Bächler (peter.baechler@kit.edu)

9 **Abstract.** The monitoring of ultrafine particle concentrations in ambient air is gaining relevance within the revision of the EU
10 Ambient Air Quality Directive. A prominent source of ultrafine particles (UFP) are combustion processes (e.g. within the
11 scope of wood-fired domestic heating) where the particle emission is typically led unfiltered into the environment contributing
12 significantly to local air pollution. In this study, ultrafine particle concentrations were measured in a residential area affected
13 by wood-smoke pollution during the winter months (Nov. 20, 2024 – Mar. 30, 2025) using a diffusion charge based UFP-
14 monitor (AQ Guard Smart 2000 from Palas®). The measurements show a diurnal trend, where concentrations are significantly
15 increased (e.g. $> 10\,000\text{ cm}^{-3}$) above the background level (approx. $5\,000\text{ cm}^{-3}$) during the morning (approx. 08:00) and evening
16 hours (approx. 19:00 – 22:00), whereby the source is wood-smoke from the surrounding neighbourhood. The dispersion
17 conditions significantly affect the measured concentrations, as only in case of low (or zero) wind speeds, increased UFP
18 concentrations are obtained demonstrating the relevance of local sources (wood-stove operation) on air quality. In the context
19 of “good practice statements” offered by the World Health Organization’s Air Quality guidelines, the maximum daily 1-hour
20 mean concentration of $20\,000\text{ cm}^{-3}$ is exceeded on approx. 33.6% of days during the measurement period. This significant
21 peak exposure on smaller timescales requires monitoring on a high temporal resolution, as longer averaging periods (e.g. daily
22 or annual mean concentrations) do not reflect temporal peak concentrations that can be especially dangerous for high-risk
23 groups. There is no direct link between legally relevant particulate matter (e.g. PM_{2.5}) and ultrafine particle concentrations, as
24 the size distribution of the wood-smoke emission is in the nanometer region and does not significantly contribute to mass-
25 based particulate matter concentrations.

26

27 1 Introduction

28 While there has been significant progress to improve air quality and implement air quality monitoring across the globe, air
29 pollution is still a major factor for health issues and premature deaths (WHO, 2021). Especially in developing countries, smog
30 in cities and traffic hotspots is a regular occurrence significantly impacting the life of the corresponding population (Amin et
31 al., 2024; Le Huong et al., 2024; Mohan et al., 2024; Moreno et al., 2025). When it comes to the European Union, measures
32 to improve air quality have shown a positive effect over the last decades, however there is still room for improvement and
33 recommendations regarding particulate matter concentrations proposed by the World Health Organization (WHO) in 2021 are
34 not necessarily met universally across the European Union. Only recently, the Ambient Air Quality Directive has been revised
35 in order to approach past recommendations set by the World Health Organization in 2005 (European Union, 2024; WHO,
36 2005). In the context of particulate pollutants, the limits for PM₁₀ and PM_{2.5} mean annual concentrations, as well as maximum
37 mean daily concentrations and the number of exceedance days are lowered towards the year 2030 according to Table 1.
38 Furthermore, a new focus is set on “pollutants of emerging concern”, which are only rarely monitored by government official
39 monitoring stations, such as ultrafine particles (UFP) and black carbon (BC). Ultrafine particles are suspected to be especially
40 hazardous due to their ability to penetrate deep into the human body, yet the direct evidence of many studies remains
41 inconclusive (Ohlwein et al., 2019). These pollutants (among others e.g. size distribution of UFP, oxidative potential, etc.)
42 now have to be monitored by so called “supersites” to gather more data and better evaluate health effects. While monitoring
43 of UFP and BC is becoming mandatory, there are no specified limits for these pollutants in the directive.
44 A universal limit for UFP would be difficult to implement, as there is no wide-spread measurement (yet) and the natural
45 formation of UFP due to atmosphere chemistry can vary for different locations and result in different background levels and
46 temporal behavior (e.g. particle growth; new particle formation) that would have to be considered in the context of quantitative
47 UFP-limits [e.g. (Bianchi et al., 2016; Dunne et al., 2016; Marten et al., 2022; Pierce and Adams, 2007; Shrivastava et al.,
48 2024). Nonetheless, the World Health Organization does offer some “good practice statements” when it comes to ultrafine
49 particle concentrations (Table 1). It can be assumed, that the specified concentrations relate to hazardous substances (e.g. such
50 as black-carbon) due to the possibility of non-hazardous high “background” concentrations due to the specified atmospheric
51 processes.

52 **Table 1 – Limits and recommendations for particulate pollutants by the WHO (2021) and the European Union (2024)**

Pollutant	Type of Limit	Revised Ambient Air Quality Directive <i>Dec. 11, 2026 / Jan. 1, 2030</i>	WHO Air Quality Guidelines (2021)
PM_{2.5}	Annual mean	<u>$25 \mu\text{g m}^{-3}$</u> $10 \mu\text{g m}^{-3}$	$5 \mu\text{g m}^{-3}$
	Daily mean	<u>Not considered</u> $25 \mu\text{g m}^{-3}$ (18 exceedance days/year)	$15 \mu\text{g m}^{-3}$ (approx. 3-4 exceedance days/year)
PM₁₀	Annual mean	<u>$40 \mu\text{g m}^{-3}$</u> $20 \mu\text{g m}^{-3}$	$15 \mu\text{g m}^{-3}$
	Daily mean	<u>$50 \mu\text{g m}^{-3}$ (35 exceedance days/year)</u> $45 \mu\text{g m}^{-3}$ (18 exceedance days/year)	$45 \mu\text{g m}^{-3}$ (approx. 3-4 exceedance days/year)
UFP	Annual mean	No limit considered (Monitoring obligation)	Not considered
	Daily mean	No limit considered (Monitoring obligation)	Good practice statement: Distinction between high and low concentration low: $C_n < 1000 \text{ cm}^{-3}$ high: $C_n > 10\,000 \text{ cm}^{-3}$
	Hourly mean	No limit considered (Monitoring obligation)	Good practice statement: $20\,000 \text{ cm}^{-3}$

53
54 Two different 24-hour mean concentration levels are specified, whereby concentrations up to $1\,000 \text{ cm}^{-3}$ are considered “low”
55 and concentration levels exceeding $10\,000 \text{ cm}^{-3}$ are considered “high” and should therefore be avoided to prevent long-term
56 health effects. In addition to these daily means, a 1-hour maximum mean concentration of $20\,000 \text{ cm}^{-3}$ is specified in the good
57 practice statements that reflects short-term exposure to UFP. This averaging period (1-hour) is rather unconventional for
58 particulate pollutants where the shortest averaging period (PM_x fine-dust limit) is 24 hours. For short averaging periods
59 concentration spikes that occur within limited timeframes are much more significant and require measurement technologies
60 and data evaluation with sufficient temporal resolution (Thieringer et al., 2022). This is especially relevant due to the
61 fluctuating nature of ultrafine particles, that are easily carried by wind and can occur due to secondary organic aerosol
62 formation and coagulation.

63 In general (though unique regulation e.g. for small member states may apply), each member state has to provide 1 supersite at
64 an urban / urban background location for each 1 000 000 inhabitants of the country and 1 supersite at a rural background
65 location for each 100 000 km² area. The spatial resolution of these measurements is therefore limited and only several important
66 sources of ultrafine particles and black carbon on ambient air quality (e.g. traffic, industry, etc.) may be covered by the
67 supersites. A supersite should be located downwind the main wind direction of a relevant pollution source. There are many
68 prominent anthropogenic sources for ultrafine particles that are considerably more harmful compared to the natural formation
69 of UFP due to atmosphere chemistry. In literature, many studies focus on air pollution with UFPs in urban environments and
70 traffic (e.g. Dall’Osto et al., 2013; Garcia-Marlès et al., 2024a; Garcia-Marlès et al., 2024b; Kumar et al., 2023; Samad et al.,
71 2022; Trechera et al., 2023; Wang et al., 2011; Zhu et al., 2002).

One major source are combustion processes, where solid or liquid fuels can cause the release of large amounts of soot, ash and other toxic substances (e.g. secondary organic aerosols) into the atmosphere (Nyarku et al., 2021). Indoor sources linked to human activity are commonly studied in exposure assessments (Jeong et al., 2019; Turner et al., 2024). The penetration of particles from outdoor into indoor spaces can dictate the indoor background level and have a major impact on long-term exposure to pollutants (Matson, 2005). In many industrial sites (e.g. during metalworking processes) worker protection from UFP is an important aspect (Elihn et al., 2009; Brouwer et al., 2004). In the automotive and industrial sector, waste gas cleaning [e.g. (fabric) filters, electrostatic precipitators, diesel and gasoline particle filters] can be applied to significantly lower the particle emission of these processes so that they do not contribute to air pollution as significantly (Bächler et al., 2024b; Hammer et al., 2024). Due to the efficiency of waste-gas cleaning technologies and the introduction of EURO norms for vehicles, the impact of traffic-related exhaust emissions on ambient air quality has decreased (Garcia-Marlès et al., 2024b; Leopoldina, 2019). As a consequence, non-exhaust emissions like brake and tire wear are gaining relevance in the automotive sector (Beddows et al., 2023). Air traffic and airports are additional prominent sources / hotspots for ultrafine particles typically featured in literature, whereby the transport of the pollutants plays a major role (Chen et al., 2010; Dröge et al., 2024; Ridolfo et al., 2024; Stacey, 2019; Trebs et al., 2023; Tremper et al., 2022). The role of aerosol transport has also been demonstrated by Junkermann et al. (2022) during aerial measurements where plumes emitted from industrial firing plants at large heights were transported over long distances.

Areas that are comparably rarely featured in official measurements are residential areas in rural or suburban background locations (Dada et al., 2025). The role of wood-fired domestic heating on ambient air quality was shown in a past study for the heating period 2022 / 2023 in a German town (Bächler et al., 2024a). Bari et al. (2009) investigated the relevance of wood-smoke pollution for PM₁₀ in a residential area and Thieringer et al. (2022) highlighted the relevance of short-term concentration peaks in the context of wood-smoke pollution from domestic heating. Especially during the evening hours, wood stoves are used for heating in individual homes and the resulting exhaust negatively impacts ambient air quality. The exposure relevance of residential areas is especially high, since high risk groups (e.g. children, elderly) are often staying in close proximity to their homes and wood-stove operation and the corresponding pollution falls into the evening hours where people are spending time at home. Furthermore, ultrafine particles / aerosols from domestic heating are suspected to have a higher toxicity compared to e.g. traffic sources (Utinger et al., 2025), though this of course depends on the exact origin of the particles (e.g. exhaust vs. non-exhaust emissions of cars). As an example, brake wear can cause the emission of and exposure to heavy metals (Neukirchen et al., 2025) that are very harmful to human health. In any case, the exposure to wood-smoke can show damaging health effects due to a large variety of potential gaseous and solid organic substances (Dilger et al., 2023).


The significance of domestic heating on air quality is also addressed in the revision of the EU ambient air quality directive, as e.g. if certain fine-dust limits cannot be reached within the specified deadline by 2030 without replacing significant fractions of existing (wood-fired / fossil fuel related) domestic heating systems – it is possible to postpone the deadline until compliance with the new limits becomes mandatory. While this procedure is reasonable since the replacement of heating systems can be a large financial investment for individual people, the impact of domestic heating on air quality is still an important issue and

106 studies are not featured as visibly compared to more prominent pollution sources (e.g. traffic or air traffic). Gravimetric fine-
 107 dust limits (compare Table 1) are known to not properly reflect air pollution with ultrafine particles which are predominantly
 108 emitted by wood fires and the corresponding wood-stoves (Trojanowski and Fthenakis, 2019; Wang et al., 2020). Thus, there
 109 is a dissonance of the particle emission (as well as particle emission limits for wood stoves that also relate to gravimetric
 110 concentrations) and particle size of the air pollutants. For smaller firing plants, e.g. for domestic heating, waste gas cleaning is
 111 not mandatory and the exhaust aerosol is let unfiltered into the environment, as the cost of gas-cleaning technologies (here:
 112 primarily electrostatic precipitators) and the lack of regulation is not promoting wide-spread application. The “Blauer Engel /
 113 blue angle – the german ecolabel” proclaims “clean” wood stoves and sets voluntary limits for mass (15 mg/m³) and number
 114 based (3 000 000 #/cm³) particle emissions (Blauer Engel, 2025). Within German regulation, such number based limits are not
 115 considered and only gravimetric concentrations are relevant, whereby the concentrations are higher compared to the ecolabel
 116 [e.g. 40 mg/m³ for most types of wood stoves; 20 mg/m³ for pellet stoves with water compartments, 30 mg/m³ for pellet stoves
 117 without water compartments; values obtained from (Bundesministerium für Justiz und für Verbraucherschutz, 2010) for
 118 installations past the year 2014].
 119 While the step towards number-based concentrations as limits for the emissions of combustion processes is addressing the
 120 dissonance between the emission limit and the actual emitted size distribution, the number concentration limit to be awarded
 121 the “Blauer Engel” is still high considering the mitigation potential that would be achievable with sophisticated waste-gas
 122 cleaning technology such as proper filtration systems (Mylläri et al., 2025; Bächler et al., 2024b; Schiller & Schmid, 2015;
 123 Matthes et al., 2016). Efficiencies of electrostatic precipitators are comparably low considering the high-particle number
 124 concentrations from the combustion process. Cornette et al., (2024) reported number based collection efficiencies of an
 125 electrostatic precipitator for wood-stoves of 83 – 92 %. Mukherjee et al., (2024) reported mass based removal efficiencies of
 126 approx. 70% that were linked to an increase in ultrafine particle emissions. Bürger and Riebel (2022) reported separation
 127 efficiencies of electrostatic precipitators in similar ranges for high temperature processes.
 128 As such the effect on ambient air quality from wood-smoke emissions can be significant and is subject to many external factors
 129 such as the dispersion conditions (e.g. wind speed and direction) as well as the geometry of the surrounding buildings in the
 130 residential area (Blocken et al., 2011, Teutscher et al., 2025). The relevance of strong local sources on ambient air quality was
 131 demonstrated by Norra et al. (2023) during (indicative) air quality measurements using a bicycle in an urban environment that
 132 was heavily influenced by local sources outside the scope of official ambient air quality monitoring. A similar situation exists
 133 in residential areas, where many temporal local sources (in case of this study: wood-smoke) can contribute to air pollution.
 134 This publication presents measurements of the concentration of ultrafine particles in a residential area for a large fraction of
 135 the heating period of the years 2024 / 2025, where wood-smoke from domestic heating significantly affects ambient air quality.
 136 A diffusion-charge based measurement device is used; this type of devices has shown great potential for indicative stationary
 137 and mobile measurements (Asbach et al., 2024). The data is evaluated with high temporal resolution and discussed in the
 138 context of the good practice statements offered by the World Health Organization. In addition to UFP concentration
 139 measurements, the effect of pollutant dispersion on the measured particle concentration is characterized.

140 **2 Materials and Methods**

141 **2.1 Description of residential area and measurement setup**

142 The measurements were performed in the rural German town “Stutensee” (approx. 25 000 inhabitants split among several
143 districts; 45.68 km² area) close to the city of Karlsruhe. Figure 1 gives an overview of the residential area as well as the
144 measurement setup, which was expanded (e.g. addition of BC and wind measurement) compared to a previous publication
145 featuring exploratory measurements for the heating period of 2022 / 2023 (Bächler et al., 2024a). The measurement devices
146 were installed on the 1st floor balcony on a garage of one of the buildings in a height of approx. 4 m. There was some obstruction
147 (e.g. wall on the east side) that could influence the wind measurement to some degree, however no explicit wind-shielding was
148 employed and there was no significant surrounding vegetation.



149
150 **Figure 1: Measurement location in a residential area in “Stutensee” close to the German city of Karlsruhe, reference wind**
151 **measurement at KIT Campus North and picture of the measurement setup including the aerosol measurement technology.**
152 **© OpenStreetMap contributors 2025. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.**

153 The measurement devices shown in Figure 1 are the AQ Guard Smart 2000 (diffusion charge based UFP monitor;
154 manufacturer: Palas® – compare subsection 2.2), a Fidas Smart 100 (aerosol spectrometer; manufacturer: Palas®) and an
155 MA350 (BC measurement; manufacturer: Aethlabs). The wind measurement serves only as an indication on the local wind
156 speed and wind direction and is not necessarily representative of the meteorological wind situation for the entire area. In the
157 map in Figure 1, the position of KITmast, a 200 m high tower for accurate meteorological wind measurements is shown that
158 serves as a reference to the locally measured wind speed and direction (compare chapter 2.3). This publication focuses on UFP
159 measurements in the context of pollutant dispersion during the majority of the heating season of 2024 / 2025 (data from Nov.
160 20, 2024 up to Mar. 30, 2025). Explorative Black carbon measurements (MA350) will be presented in sect. 3.1 to offer further
161 context regarding source apportionment. A follow-up publication for further analysis of the temporal behavior and the
162 correlation between UFP, PM_x and BC will analyze the MA350 data in detail. Further data on relative humidity, ambient
163 temperature and pressure from the Fidas Smart 100 can be found in the supplementary information.

164 **2.2 Description of diffusion-charge based UFP-monitor**

165 The UFP particle number concentration was measured using an AQ Guard Smart 2000 from the manufacturer Palas®. The
166 specifications of the measurement device are summarized in Table 2.

167 **Table 2 – Specifications of AQ Guard Smart 2000 according to the manufacturer (Palas, 2025)**

Measurement principle	Diffusion charging
Measurements	Particle number concentration, Median particle diameter, Lung deposited surface area, pressure, temperature, relative humidity
Particle number concentration range	1 000 – 10 ⁸ cm ⁻³
Lower particle size detection limit	10 nm
Temperature range	-20°C - +40 °C
Dimensions	530 mm · 270 mm · 208 mm
Weight	Approx. 6 kg
Miscellaneous information	Heated inlet, cloud data storage on “MyAtmosphere” platform


168
169 The working principle is based on unipolar diffusion charging of the sample aerosol. At the inlet the sample aerosol is heated
170 to avoid any humidity or condensation effects. Ionized sheath air (filtered by HEPA and activated carbon filters) is generated
171 through unipolar corona discharge. The ambient aerosol is mixed with the ionized sheath air in an ejector / dilutor system at a
172 fixed ratio. Thus, the aerosol is charged by coagulation and excess ions are removed in an ion trap before the charged aerosol
173 (known charge distribution) passes a faraday cup electrometer and the generated current is measured. A mobility analyzer with
174 varying voltage serves as a classifier and generates information on particle size.

175 The data is collected on the cloud database “MyAtmosphere” (my-atmosphere.net) provided by the manufacturer and can be
176 accessed in varying temporal resolutions (1-minute; 15-minute; 1-hour; 24-hours). The wind measurement was linked to the
177 cloud as well enabling a direct allocation of UFP concentration, wind speed and wind direction with high temporal resolution.
178 The data was post-processed regarding incorrect concentration measurements (e.g. unplausibly high or negative concentration
179 readings). This affected only a negligible fraction of the data. The (raw-)dataset is freely available and published under an
180 open-access license according to data availability statement at the end of the article (Bächler et al., 2025).

181

182 **2.3 Characterization of the dispersion conditions**

183 Since the wind measurement was located together with the aerosol measurement devices on the 1st floor balcony of a garage
184 of one of the buildings in the residential area, the corresponding wind data is influenced by the building geometry of the
185 surrounding neighborhood, as e.g. the flow of wind may be obstructed by the buildings. Nonetheless, the measurement serves
186 as an indicator for pollutant dispersion, where high wind speeds reflect a high degree of pollutant dispersion / dilution and low
187 (or non-measurable) wind speeds are an indication for a low degree of pollutant dispersion.
188 In addition to this local wind measurement, data from a meteorological measurement station (compare Figure 1) was provided
189 (Kohler et al., 2018). Figure 2 shows a wind-rose comparing the local wind speed and direction at the measurement setup in
190 “Stutensee” with the meteorological reference measurement (compare figure 1) from KITmast (40 m height) as well as the
191 average diurnal wind behavior that was calculated for the entire measurement period. Here, in case of the reference
192 measurement from KITmast, data from 40 m height and 2 m height is evaluated, whereby wind direction is only measured at
193 40 m height. The measurement at 40 m height is unobstructed by the vegetation zone of the nearby forest and offers
194 representative data “close” to the ground, while the 2 m measurement is located on a clearing where vegetation has a
195 comparably low impact but still influences the corresponding measurements. Both datasets are presented to offer a holistic
196 comparison to the wind measurement in the residential area. For the data obtained at the residential area, only wind direction
197 data with wind speeds > 0 km/h are evaluated, as the device reports 0° wind direction for wind speeds of zero. The wind rose
198 shows the probability / fraction of the dataset of different wind speeds (above zero) coming from a corresponding wind
199 direction in %.



200 **Figure 2: Wind rose and the average average daily behavior of the wind speed for the residential area and the reference**
201 **measurement (KITmast) at 2 m and 40 m height (Kohler et al., 2018)**
202

203 Main wind directions for the reference (NE, ENE and SW) are different from the residential area (SE). Note that the building
204 geometry in the residential area can have a significant impact on the measured wind speed and direction. Thus, the absolute
205 wind speeds obtained at the residential area and from the reference at 2 m height are differing significantly. At an increased
206 height of 40 m where there is no impact of the vegetation zone, the measured wind speeds are significantly higher as is
207 expected. Considering the qualitative behavior of the wind speed, all measurement positions show a similar trend, where
208 average wind speeds are relatively stable ranging from 18:00 in the evening up to approx. 09:00 in the morning. In between
209 09:00 and 18:00 there is a temporal increase in wind speed (e.g. from 0.25 km/h up to 0.6 km/h for the residential area). This
210 diurnal behavior has to be considered in the context of aerosol measurements. A similar profile was reported by Kuhlbusch et
211 al. (2001) for another location in Germany. Due to the same trend between the reference and the wind measurement in the
212 residential area, the locally obtained wind speeds are a valid indication for the corresponding pollutant dispersion conditions.
213 Therefore, only the wind speeds at the measurement position in the residential area will be considered for further evaluation.
214 A Heat Map showing 15-minute average wind speeds for the entire measurement period is provided in the supplementary
215 information.


216

217 **3 Results and Discussion**

218 **3.1 Temporal evolution of daily UFP concentration for the entire measurement period**

219 Wood-stoves are commonly operated during the evening hours within the heating period, causing a characteristic diurnal
220 pattern whereby increased concentrations above the background level are only detected during periods of source activity.
221 Figure 3 shows a Heat Map of 15-minute average UFP concentrations for the entire measurement period ranging from Nov.
222 20, 2024 up to Mar. 30, 2025, resulting in 131 days of complete measurement with 24 hour data availability. Concentrations
223 below the background level of 5 000 cm⁻³ are in blue color. This lower threshold concentration corresponds to the common
224 “natural” background level of the residential area – note however that the “low” concentration level according to the WHO
225 good practice statement (Table 1) is lower by comparison and the color scale aims to highlight exceedances of the background
226 concentration level. The color scale shifts towards yellow color that represents the “high” concentration level of 10 000 cm⁻³
227 according to the WHO good practice statements for exposure within a 24 hour mean concentration and to red color for values
228 exceeding a concentration of 20 000 cm⁻³ which is classified as a daily maximum hourly mean UFP concentration according
229 to the WHO good-practice statements. Note that the peak concentrations can significantly exceed this maximum threshold of
230 20 000 cm⁻³ in the Heat-Map. A UFP Heat Map with a more detailed scale considering the peak concentrations for Dec. 2024
231 – Feb. 2025 with higher temporal resolution is shown in the supplementary information.

232




233

234

Figure 3 – Heat Map of 15-minute mean UFP-concentration for the entire duration of the measurements

235 The Heat Map illustrates, that a significant fraction of the measurement period is subject to air pollution in the morning and
 236 evening hours what is likely linked to human activity.
 237 Further evaluating the data shown in the heatmap in Figure 3, an “average” diurnal particle concentration dynamic was
 238 calculated in Figure 4 and set into the context of the diurnal wind pattern (Figure 2). Here, the overall average 15-minute mean
 239 concentration in a fixed time interval e.g. from 09:00 to 09:15 is calculated for either the entire measurement period (left) or
 240 only the days where the 1-hour average concentration of 20 000 cm³ was exceeded (right). E.g. for Fig. 4 left: the average
 241 concentration for an entire row of the Heat Map is calculated for each time interval to gain a “mean” diurnal pattern for the
 242 entire measurement period. The black curve represents the average 15-minute mean UFP-concentration for the entire
 243 measurement duration. Here, days with and without increased UFP concentrations during the evening hours are taken into
 244 account. Despite days with low concentrations impacting the calculated mean concentrations, the determined concentration
 245 profile shows a pronounced trend, where particle concentrations above the background level (reaching up to around
 246 10 000 cm⁻³) are measured during the morning hours (approx. 8:00 – 10:00) as well as the evening hours
 247 (approx. 18:00 – 22:00).



248
 249 **Figure 4 –Diurnal UFP and wind pattern calculated from 15- minute mean concentrations at the corresponding local time for the**
 250 **entire measurement period and measurement days where 1-hour mean concentration of 20 000 cm is exceeded (compare Figure 9)**

251 During daytime, slightly increased wind speeds are measured (compare Figure 2) causing a larger degree of pollutant
 252 dispersion. So not only are the major sources likely not active during the lunch hours, pollutant dispersion has positive impacts
 253 on the particle concentration level. Identifying pollutant sources is not possible based on concentration data alone, however
 254 section 3.1 offers some insights regarding the corresponding source apportionment for the conclusions drawn from Fig. 3 and
 255 Fig. 4. Considering only measurement days with significant concentration spikes, where 1-hour mean concentrations of 20
 256 000 cm⁻³ were exceeded (compare Figure 9 and sect. 3.4), the diurnal pattern is even more pronounced. The wind speed during
 257 these days with increased air pollution was in a similar region compared to the entire measurement period and also showed the
 258 typical diurnal pattern with no significant deviation regarding the higher wind speeds during the daytime. A similar diurnal
 259 pattern with corresponding concentration increases was reported by Dada et al., (2025) in rural Switzerland, where especially

260 during winter and autumn increased concentrations were measured during morning (approx. 8 000 cm⁻³) and evening hours (6
261 000 cm⁻³) compared to the background level (4 000 cm⁻³). The source was identified as domestic heating and aerosol transport
262 from rush-hours based on intercomparisons of various different measurements (e.g. SMPS, CPC, Catalytic Stripper, NAIS,
263 Trace-gas analysis, Aethalometer, etc.). Another similar diurnal pattern for moderately traffic influenced urban and regional
264 background sites regarding black carbon and aerosol particles in the range of 30 – 200 nm was reported by Sun et al. (2019)
265 where peak concentrations occurred in the morning and evening hours that were allocated to traffic, cooking and heating.
266 Kuhlbusch et al. (2001) also showed a similar diurnal pattern linked to traffic emissions in terms of PM-concentrations.
267 Summarizing, daily UFP concentration evolution is strongly influenced by local sources (wood-stoves) and pollutant
268 dispersion is a significant factor affecting the local concentration level and the measurements.

269 **3.1 Source apportionment in the context of the dispersion conditions**

270
271 Many different natural (e.g. atmosphere chemistry) and anthropogenic (e.g. wood-smoke, abrasion, etc.) sources can contribute
272 to increased UFP concentrations. Previous investigations proposed wood-stoves as the main source for increased UFP
273 concentration levels in this residential area. Of course elevated UFP concentrations alone are not sufficient to verify wood-
274 smoke from domestic heating as the source – however there is sufficient evidence to unambiguously identify the wood-smoke
275 pollution in the residential area:

- 276 - Smell and visual identification of plumes from chimneys of the surrounding neighborhood show the emission of
277 wood-stove exhaust and are direct evidence, even though this is no quantifiable data.
- 278 - Additional black carbon measurements also show qualitatively similar concentration spikes in the evening so that the
279 ultrafine particles likely contain soot from (incomplete) combustion. Unfortunately, the data availability for these
280 measurements only covers several days of the measurement period. The focus of this article remains UFP-
281 concentrations and the Black-Carbon data and the link between PM, UFP and BC on a temporal basis will be analyzed
282 in detail in a follow-up publication. Figure 5 shows the Black-Carbon concentration obtained from the MA350 device
283 for days with 24 hour data availability to serve as further indicative proof regarding the source of increased UFP and
284 BC-concentration levels. Especially during the evening hours, Black-Carbon concentrations of approx. 5 µg m⁻³ are
285 reached. For the inversion period including Feb. 8 and Feb. 9 (compare sect. 3.2.2.) concentrations up to
286 10 µg m⁻³ were measured regarding 15-minute average concentrations (potentially higher concentration spikes –
287 maximum 15-minute average concentration of approx. 17 µg/m³).




Figure 5 – Heat Map of 15-minute mean BC-concentration for several exemplary days (24-hour data availability) during the measurement period

- The increased concentrations in the evening are occurring outside typical rush hours and often reach their peak after 18:00. While there indeed is a highway (A5 in Figure 1) located in the main wind direction (according to the indicative wind measurement which may be influenced by building geometry) that could be a potential source of ultrafine particles from traffic – increased concentrations only occur for low wind speeds close to or below the detection limit of the wind measurement. As such, the transport of pollutants from the highway to the residential area can be ruled out or is at least unlikely to cause concentration spikes in that order of magnitude. As a consequence, increased concentrations in the residential area have to originate from local sources.
- New-particle formation events would have to coincide with other sources and do not occur in larger numbers. Such events should also be mostly independent of wind-speed. Song et al. (2024) did not report significant occurrence of new-particle formation events at another location at the outskirts of Karlsruhe within reasonable proximity to the measurement site. However these investigations were performed during the summer months and are not comparable.
- Other prominent UFP sources (e.g. air traffic) can be ruled out.


3.2 Effect of pollutant dispersion on short timeframes for several exemplary measurement periods

To further demonstrate the impact of pollutant dispersion on ultrafine particle concentrations, several exemplary days or periods are discussed in this section. Correlations for other ambient conditions such as temperature and pressure can be found in the supplementary information – though the impact of these parameters shows no distinctive trend compared to wind speed / pollutant dispersion.

3.2.1. New-years fireworks

A prominent example during the heating period where air pollution is prevalent is the New-Year's celebration, where fireworks are launched in large amounts at midnight (00:00). The explosion of fireworks contributes to air pollution and, while only being a temporal occurrence once a year, can demonstrate the dynamic behavior of UFP concentrations. Other authors also

312 featured the impact of firework celebrations on air pollution (Joshi et al., 2019). Figure 6 shows the UFP concentration in the
 313 context of wind speed / dispersion conditions for the residential area featured in this publication for the year 2024/2025.




314
 315 **Figure 6 – UFP number concentration and dispersion conditions during New-Year’s fireworks.**

316 Over midday and up to the late afternoon at 4 PM, UFP concentrations were around a typical background level and the wind
 317 speed was comparably high at the measurement position (compare Figure 2). In the evening hours from approx. 17:00 until
 318 midnight, concentrations were fluctuating between 5 000 and (mostly) 10 000 cm⁻³, whereby several short-term peaks reached
 319 concentrations exceeding 20 000 cm⁻³. This noise in the data is an indication for source activity, even though average
 320 concentrations are not noticeably high. Reaching 00:00 and the start of the fireworks, concentrations quickly increase, where
 321 a peak concentration of 90 000 cm⁻³ is reached at 00:30. The peak concentration disperses rather quickly until approx. 1:30
 322 during a period, where elevated wind speeds were measured. Results from (Drewnick et al., 2006) showed larger peak
 323 concentrations (up to 150 000 cm⁻³ – 5 minute average concentration) for a German city, however the measurement position
 324 was situated close to the center of firework activity. Drutschke et al. (2011) also report higher peak concentration for firework
 325 festivities up to 1 600 000 cm⁻³. After the decay of the main concentration peak, the concentration level remains relatively
 326 constant at approx. 21 000 cm⁻³. From 2:30 up to approx. 5:00, concentrations fluctuate during a period of pollution dispersion.
 327 Finally, from 5:00 up to the morning hours of 10:00, no wind speed is detected and the natural dispersion of the pollutants is
 328 comparably slow so that concentrations decrease from 20 000 down to 10 000 cm⁻³ over a longer period of time before
 329 increased wind speeds during lunch bring the UFP concentration to the background level.

332 **3.2.2. Stationary temperature inversions / High-pressure-area**

333 An example for a period of the measurement campaign, where the dispersion conditions for UFP and other air pollutants was
334 significantly impaired was during the prevalence of the high-pressure-area “Elvira” in February 2025. The high-pressure-area
335 / stationary temperature inversion prevented air exchange that led to the accumulation of air pollutants. This can have a
336 significant impact on air quality, especially for cities located in basins with unfavourable dispersion conditions (Zhang et al.,
337 2024). This effect on air pollution can also be observed on a temporally resolved basis in the residential area regarding UFP
338 concentrations (Figure 7) and was prominently featured across different news platforms in Germany. It is also clearly visible
339 within the PM_{2.5} Heat Map shown in the supplementary information.




340

341 **Figure 7 – UFP number concentration and dispersion conditions during prevalence of a high pressure area / stationary**
342 **temperature inversions in February 2025**

343 Increased wood-stove operation led to an increase in UFP concentration during the evening hours of Feb. 8. Due to poor
344 pollutant dispersion and the beginning of the impact of the high-pressure-area, this level remained constant up to the lunch
345 hours of the next day. In the evening of Feb. 9, concentrations again increased up to three times the background concentration
346 and remained at a level of $10\,000\text{ cm}^{-3}$ (high concentration according to WHO good practice statement) for the entirety of Feb.
347 10. On Feb. 11, the concentration slowly decreased to the background level up to the evening hours, where wood-stove
348 operation again contributed to air pollution. Towards the end of the effective duration of the high-pressure-area on Feb. 12 and
349 13, the concentration decreased and, especially on Feb. 13, the concentration was at or below the background level and
350 significant wind speeds were measured for the majority of the day. Noise in the data of Feb. 13 is again an indication for source
351 activity and simultaneous dispersion / dilution.
352

353 **3.3 UFP concentration levels in the context of pollutant dispersion during the entire measurement period**

354 To give a direct correlation between ultrafine particles and wind speed, a scatterplot is drawn for the 15-minute average UFP
355 concentrations and the corresponding average wind speed in Figure 8. Correlations for different ambient conditions can be
356 found in the supplementary information – although no clear trends could be derived from this evaluation. In Figure 8, different
357 sectors can be identified that illustrate the main conclusions drawn from the previously discussed measurements of UFP in the
358 context of wind speeds / pollutant dispersion.



359

360 **Figure 8 – Scatterplot of 15- minute mean UFP-concentrations and wind speeds for the entire measurement period**

361 The highest UFP concentrations occur under conditions where pollutant dispersion is very low (red and yellow sector). The
362 threshold wind speed was selected at 0.5 km/h as almost no concentrations above $10\,000\text{ cm}^{-3}$ were measured at higher wind

363 speeds. Data above the background level at low wind speeds amounts to 44.7% of the total data and was likely caused by
364 source activity and wood-smoke pollution.

365 For wind speeds above 0.5 km/h and concentrations above 5 000 cm⁻³ only a very limited amount of data can be found (4 %
366 total) by comparison. With increasing pollutant dispersion, even under conditions with high source activity the effect on air
367 quality is not as significant.

368 In the sector including wind speeds above 0.5 km/h and concentrations below 5 000 cm⁻³ a total of 18.5 % of the measurement
369 data can be found. Pollutant dispersion dominates the overall particle concentration level so that even in case of source activity,
370 the effect on ambient air quality is not measurable, impeding the generation of emission inventories (e.g. climate-damaging
371 black carbon from incomplete combustion from wood-stoves). This data is predominantly obtained during the lunch hours
372 with increased wind speeds.

373 Only for data below the background concentration and at low wind speeds source activity can be ruled out with high certainty.
374 The amount of data in this quadrant amounts to 32.8%.

375 Summarizing, the dispersion conditions have a major impact on the measured air quality in the residential area. In case of low
376 pollutant dispersion, increased concentration levels can persevere across multiple days. For higher wind speeds, pollutant
377 dispersion can dominate the ambient air quality and low concentration can be measured, despite the possibility of wood-smoke
378 emissions.

379 **3.4 Evaluation of average UFP and particulate matter concentrations regarding ambient air quality limits and WHO** 380 **recommendations**

381 In the previous sections, the measured UFP concentrations were already discussed in the context of particle concentration
382 levels introduced by the World Health Organizations' "good practice statements". Particle concentrations above 10 000 cm⁻³
383 (24-hour mean) are considered a high concentration. Furthermore, a daily maximum 1-hour mean concentration of
384 20 000 cm⁻³ should also not be exceeded. In Figure 9, the daily mean UFP concentration as well as the maximum 1-hour mean
385 concentration for the entire measurement period ranging from Nov. 20, 2024 up to Mar. 30, 2025 are calculated and set into
386 the context of the WHO "good practice statements".

387




Figure 9 – Mean daily and maximum 1-hour UFP number concentration for the entire measurement period

Considering the 24-hour mean concentrations, only a total of 16 % of measurement days (21 / 131 days) exceed the 24-hour mean UFP concentration of 10 000 cm⁻³ (black data-points within yellow area). This is mainly caused by the diurnal pattern, where increased concentrations are only detected during the morning and evening hours so that these concentration spikes are not as significant in the context of 24-hour mean concentrations.

However, considering increased short-term exposure to high concentrations above 20 000 cm⁻³, a total of 33.6 % of measurement days (44 / 131 days) exceed the daily maximum 1-hour mean concentration at least once during the day. Considering a total of 24 hours for each measurement day, the number of hours exceeding the 20 000 cm⁻³ threshold is 154 / 3144 which corresponds to 4.9% of hours during the measurement period. These increased concentration levels are very likely hazardous due to their origin from wood combustion and can contribute to long-term health effects and endanger high risk groups such as children and the elderly regarding short-term health effects.

3.5 UFP concentrations in the context of PM_{2.5} measurements

To add further context to the increased UFP peak concentrations (Figure 9), the mean daily PM_{2.5} concentrations obtained from scattered-light based measurements using the Fidas Smart 100 (compare Figure 1) are displayed in Figure 10. These measurements were indicative measurements and not government-official. For further information on the temporally resolved PM_{2.5} behavior, a Heat Map similar to Figure 4 can be accessed in the supplementary information.




Figure 10 – Mean daily PM_{2.5} concentration for the entire measurement period within the scope of the limits specified in the revised Ambient Air Quality Directive (indicative measurement – no government-official data)

The average PM_{2.5} concentration of the measurement period (11.9 µg/m³) is close to the future annual limit values for the year 2030. Therefore, the current mean annual concentration limit of 25 µg/m³ is easily kept. As the measurement period was performed during the heating season, where higher PM concentrations are expected, there will likely be no (or no significant) exceedance of the future 10 µg/m³ limit in case of the year 2025. When it comes to the future number of 18 exceedance days of a concentration of 25 µg/m³, which will be relevant starting in 2030, there were a total of 13 exceedance days during the measurement period (all of which occurred in 2025). Several of these exceedance days were obtained during the impact of the high-pressure-area discussed in sect. 3.2.2. so that the concentration itself is not directly linked to the emission of pollutants but rather the dispersion conditions that may dictate whether or not the number of exceedance days can be complied with.




Figure 11 – 15- minute mean diurnal UFP (compare Figure 4 regarding UFP concentrations) and PM_{2.5} pattern for the entire measurement period and scatterplot of UFP and PM_{2.5} concentrations (compare Figure 9 and Figure 10)

420 In this case, the actual pollution with hazardous nanoparticles from wood combustion is not properly represented by
421 “conventional” air quality monitoring (Figure 11). There is no direct correlation between the PM_{2.5} and UFP measurements,
422 as UFP are below the detectable size range of the Fidas Smart 100 device and remain undetected (additionally, their
423 contribution to mass concentrations is very small – see also scatterplot of 15-minute average UFP and PM_{2.5} concentration in
424 the supplementary information). While there is also a slight diurnal pattern considering average PM_{2.5} concentrations –
425 compared to the large differences during peak exposure times in the morning and evening hours for UFP concentrations (e.g.
426 factor of 2 compared to background level), the absolute difference in PM is comparably low (factor 1.3 – 1.4). The (also mass
427 based but non-size-resolved) BC-concentrations often increased from a close to zero-background level up to 5 µg m⁻³ during
428 the evening hours by comparison (Figure 5).

429 Short-term exposures (e.g. covered by good practice statement for UFP in the form of 1-hour maximum concentration) are not
430 considered in the limits of the Ambient Air Quality Directive due to the daily and annual averaging periods of particle
431 concentrations.

432 These results demonstrate that UFP monitoring is especially relevant for regions where domestic heating using wood as
433 combustion resource is used. While significant UFP-concentrations were measured in the residential area, the PM
434 concentrations (mostly) comply with (or do not show a significant exceedance of) current and future PM-concentration limits.
435 For areas where measurement stations do indeed register an exceedance of PM-concentration limits due to the emissions from
436 wood-combustion for domestic heating as a main source, it has to be assumed that ambient air quality is likely affected by
437 especially hazardous (black-carbon) nanoparticles.

438 **4 Summary and Outlook**

439 Measurements of ultrafine particle concentrations applying a diffusion charge based UFP-monitor were performed in a
440 residential area in a small German town during a large fraction of the heating period of 2024 / 2025. The major source for air
441 pollution in the residential area during this time was wood-stove exhaust for domestic heating. Especially during the morning
442 and evening hours, significant UFP concentration spikes were measured, causing a characteristic diurnal pattern. Here, average
443 concentrations of approx. 10 000 cm⁻³ (corresponds to twice the typical background level) were measured averaging the entire
444 measurement period. The dispersion conditions have a large impact on the measured concentrations, as during daytime
445 increased wind speeds occur and were measured at a reference wind measurement site and in the residential area. Higher wind
446 speeds are linked to lower / decreasing UFP concentrations and only for low (or non-detectable) wind speeds large UFP-
447 concentrations (e.g. exceeding 10 000 cm⁻³ were measured. On approx. 33.6% of days during the measurement period a
448 maximum daily 1-hour mean concentration of 20 000 cm⁻³ was exceeded according to the WHO “good practice statements”
449 on ultrafine particles. During these periods with increased wood-smoke pollution (morning and evening hours) people are
450 more likely to spend time at their homes (indoors). Depending on ventilation / air exchange between interior and exterior, they
451 may or may not be exposed to this outdoor air pollution. How much individual homes are affected can thus vary significantly.

452 Due to the small particle size, these UFP from wood-smoke are not contributing significantly to legally-relevant PM-
453 concentrations, where a less pronounced diurnal pattern was shown with a scattered-light based Fidas Smart 100. This research
454 highlights the dissonance between the size distribution of wood-smoke particle emissions from domestic heating (nanometer
455 region) and conventional air quality measurements (mass based PM-concentrations). A follow-up publication will further
456 discuss explorative (temporally resolved) Black-Carbon measurements and the link to UFP and PM concentration dynamics
457 during this measurement period. Further research will aim to expand the setup to perform further characterization of the air
458 pollutants measured in the residential area.

459 **Data availability**

460 The data that support the findings of this study are openly available in the KITOpen Repository under the
461 doi:10.35097/2aab6yjdyfg4nr69.

462 **Author contribution**

463 **P.B. :** Conceptualization, Methodology, Investigation, Resources, Writing – Original Draft, Writing – Review & Editing
464 **F. W.:** Conceptualization, Validation, Resources, Data Curation, Writing – Review & Editing
465 **S. K.:** Validation, Data Curation, Writing – Review & Editing
466 **A. D.:** Conceptualization, Methodology, Validation, Resources, Supervision, Writing – Review & Editing.

467 **Competing interests**

468 The authors declare the following financial interests/personal relationships which may be considered as potential competing
469 interests: Frederik Weis and Sebastian Kohler are currently employed by Palas GmbH and worked on the development of the
470 measurement device.

471 **Acknowledgements**

472 We kindly acknowledge the collaboration of Martin Kohler from the Institute of Meteorology and Climate Research at
473 Karlsruhe Institute of Technology for providing wind data from KITMast.
474 We kindly acknowledge the collaboration of Jeffrey Blair from Aethlabs for providing the MA350 measurement device.

475 **References**

- 476 Amin, M., Aun, S., Hang, L., Hata, M., Phairuang, W. and Masami, F.: Investigation into personal exposure to ultrafine
477 particle (UFP) in Phnom Penh, Cambodia: A pilot study, *Atmospheric Pollution Research*, 12, 102307, doi:
478 10.1016/j.apr.2024.102307, 2024.
- 479 Asbach, C., Todea, A. Maria and Kaminski, H.: Evaluation of a Partector Pro for atmospheric particle number size
480 distribution and number concentration measurements at an urban background site, *Aerosol Research*, 1, 1–12, doi:
481 10.5194/ar-2-1-2024, 2024.
- 482 Bari, M. Aynul, Baumbach, G., Kuch, B. and Scheffknecht, G.: Wood smoke as a source of particle-phase organic
483 compounds in residential areas, *Atmospheric Environment*, 31, 4722–4732, doi: 10.1016/j.atmosenv.2008.09.006, 2009.
- 484 Bächler, P., Weis, F., Kohler, S., Dittler, A.: Measurement data of UFP-concentrations and PM2.5-concentrations during the
485 winter months (Nov. 20, 2024 - Mar. 30, 2025) in a residential area near Karlsruhe, KITOpen Repository [dataset], doi:
486 10.35097/2aab6yjdyfg4nr69, 2025.
- 487 Bächler, P., Weis, F., Kohler, S. and Dittler, A.: Exploratory measurements of ambient air quality in a residential area
488 applying a diffusion charge based UFP monitor/Orientierende Messungen der Luftqualität in einem Wohngebiet mit einem
489 auf Diffusionsaufladung basierten UFP-Monitor, *GrdL*, 01-02, 15–22, doi: 10.37544/0949-8036-2024-01-02-17, 2024.
- 490 Bächler, P., Meyer, J., Ligotski, R., Krug, P. and Dittler, A.: Measurement of transient nanoparticle emissions of a municipal
491 biomass incineration plant equipped with pulse-jet cleaned filters, *Process Safety and Environmental Protection*, 601–614,
492 doi: 10.1016/j.psep.2024.02.013, 2024.
- 493 Beddows, D. C. S., Harrison, R. M., Gonet, T., Maher, B. A. and Odling, N.: Measurement of road traffic brake and tyre dust
494 emissions using both particle composition and size distribution data, *Environ Pollut*, Pt 1, 121830, doi:
495 10.1016/j.envpol.2023.121830, 2023.
- 496 Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Hermann, E., Adamov, A.,
497 Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H.
498 E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala,
499 M., Dommen, J. and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing,
500 *Science*, 6289, 1109–1112, doi: 10.1126/science.aad5456, 2016.
- 501 Blauer Engel: Kaminöfen für Holz: <https://www.blauer-engel.de/de/zertifizierung/vergabekriterien#UZ212-2020>, last access:
502 07 August 2025.
- 503 Blocken, B., Stathopoulos, T., Carmeliet, J. and Hensen, J. L.M.: Application of computational fluid dynamics in building
504 performance simulation for the outdoor environment: an overview, *Journal of Building Performance Simulation*, 2, 157–184,
505 doi: 10.1080/19401493.2010.513740, 2011.
- 506 Brouwer, D. H., Gijsbers, J. H. J. and Lurvink, M. W. M.: Personal exposure to ultrafine particles in the workplace:
507 exploring sampling techniques and strategies, *Ann Occup Hyg*, 5, 439–453, doi: 10.1093/annhyg/meh040, 2004.
- 508 Bundesministerium für Justiz und für Verbraucherschutz, Bundesamt für Justiz: Erste Verordnung zur Durchführung des
509 Bundes-Immissionsschutzgesetzes, https://www.gesetze-im-internet.de/bimschv_1_2010/, last access: 07 August 2025
510 (2010).
- 511 Bürger, P. and Riebel, U.: Feasibility of high-temperature electrostatic precipitation for the removal of nanoparticles: A case
512 study on iron oxide separation at up to 800 °C, *Journal of Electrostatics*, 103754, doi: 10.1016/j.elstat.2022.103754, 2022.
- 513 Chen, S.-C., Tsai, C.-J., Chou, C. C.-K., Roam, G.-D., Cheng, S.-S. and Wang, Y.-N.: Ultrafine particles at three different
514 sampling locations in Taiwan, *Atmospheric Environment*, 4, 533–540, doi: 10.1016/j.atmosenv.2009.10.044, 2010.

515 Cornette, J. F.P., Dyakov, I. V., Plissart, P., Bram, S. and Blondeau, J.: In-situ evaluation of a commercial electrostatic
516 precipitator integrated in a small-scale wood chip boiler, *Journal of Electrostatics*, 103897, doi:
517 10.1016/j.elstat.2024.103897, 2024.

518 Dada, L., Brem, B. T., Amarandi-Netedu, L.-M., Collaud Coen, M., Evangeliou, N., Hueglin, C., Nowak, N., Modini, R.,
519 Steinbacher, M. and Gysel-Beer, M.: Sources of ultrafine particles at a rural midland site in Switzerland, *Aerosol Research*,
520 1, 315–336, doi: 10.5194/ar-3-315-2025, 2025.

521 Dall'Osto, M., Querol, X., Alastuey, A., O'Dowd, C., Harrison, R. M., Wenger, J. and Gómez-Moreno, F. J.: On the spatial
522 distribution and evolution of ultrafine particles in Barcelona, *Atmos. Chem. Phys.*, 2, 741–759, doi: 10.5194/acp-13-741-
523 2013, 2013.

524 Dilger, M., Armant, O., Ramme, L., Mülhopt, S., Sapcaru, S. C., Schlager, C., Dilger, E., Reda, A., Orasche, J., Schnelle-
525 Kreis, J., Conlon, T. M., Yildirim, A. Önder, Hartwig, A., Zimmermann, R., Hiller, K., Diabaté, S., Paur, H.-R. and Weiss,
526 C.: Systems toxicology of complex wood combustion aerosol reveals gaseous carbonyl compounds as critical constituents,
527 *Environ Int*, 108169, doi: 10.1016/j.envint.2023.108169, 2023.

528 Drewnick, F., Hings, S. S., Curtius, J., Eerdekens, G. and Williams, J.: Measurement of fine particulate and gas-phase
529 species during the New Year's fireworks 2005 in Mainz, Germany, *Atmospheric Environment*, 23, 4316–4327, doi:
530 10.1016/j.atmosenv.2006.03.040, 2006.

531 Dröge, J., Klingelhöfer, D., Braun, M. and Groneberg, D. A.: Influence of a large commercial airport on the ultrafine particle
532 number concentration in a distant residential area under different wind conditions and the impact of the COVID-19
533 pandemic, *Environ Pollut*, 123390, doi: 10.1016/j.envpol.2024.123390, 2024.

534 Dunne, E. M., Gordon, H., Kürten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle, K. J., Adamov, A.,
535 Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., Curtius, J., Dommen, J., Donahue,
536 N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J.,
537 Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J., Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J.,
538 Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C. L. S., Riccobono, F., Richards, N. A. D., Rissanen, M. P.,
539 Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M., Sipilä, M., Smith, J. N., Stozkhov, Y., Tomé, A., Tröstl,
540 J., Wagner, P. E., Wimmer, D., Winkler, P. M., Worsnop, D. R. and Carslaw, K. S.: Global atmospheric particle formation
541 from CERN CLOUD measurements, *Science*, 6316, 1119–1124, doi: 10.1126/science.aaf2649, 2016.

542 Dutschke, A., Lohrer, C., Kurth, L., Seeger, S., Barthel, M. and Panne, U.: Aerosol Emissions from Outdoor Firework
543 Displays, *Chem Eng & Technol*, 12, 2044–2050, doi: 10.1002/ceat.201100080, 2011.

544 Elihn, K. and Berg, P.: Ultrafine particle characteristics in seven industrial plants, *Ann Occup Hyg*, 5, 475–484, doi:
545 10.1093/annhyg/mep033, 2009.

546 European Union: Directive (EU) 2024/2881 of the European Parliament and of the Council of 23 October 2024 on ambient
547 air quality and cleaner air for Europe (recast), <https://eur-lex.europa.eu/eli/dir/2024/2881/oj>, last access: 07 August 2025
548 (2024).

549 Garcia-Marlès, M., Lara, R., Reche, C., Pérez, N., Tobías, A., Savadkoohi, M., Beddows, D., Salma, I., Vörösmarty, M.,
550 Weidinger, T., Hueglin, C., Mihalopoulos, N., Grivas, G., Kalkavouras, P., Ondráček, J., Zíková, N., Niemi, J. V.,
551 Manninen, H. E., Green, D. C., Tremper, A. H., Norman, M., Vratolis, S., Eleftheriadis, K., Gómez-Moreno, F. J., Alonso-
552 Blanco, E., Wiedensohler, A., Weinhold, K., Merkel, M., Bastian, S., Hoffmann, B., Altug, H., Petit, J.-E., Favez, O., Dos
553 Santos, S. Martins, Putaud, J.-P., Dinoi, A., Contini, D., Timonen, H., Lampilahti, J., Petäjä, T., Pandolfi, M., Hopke, P. K.,
554 Harrison, R. M., Alastuey, A. and Querol, X.: Inter-annual trends of ultrafine particles in urban Europe, *Environ Int*, 108510,
555 doi: 10.1016/j.envint.2024.108510, 2024.

556 Garcia-Marlès, M., Lara, R., Reche, C., Pérez, N., Tobías, A., Savadkoohi, M., Beddows, D., Salma, I., Vörösmarty, M.,
557 Weidinger, T., Hueglin, C., Mihalopoulos, N., Grivas, G., Kalkavouras, P., Ondracek, J., Zikova, N., Niemi, J. V.,
558 Manninen, H. E., Green, D. C., Tremper, A. H., Norman, M., Vratolis, S., Diapouli, E., Eleftheriadis, K., Gómez-Moreno, F.
559 J., Alonso-Blanco, E., Wiedensohler, A., Weinhold, K., Merkel, M., Bastian, S., Hoffmann, B., Altug, H., Petit, J.-E.,
560 Acharja, P., Favez, O., Santos, S. Martins Dos, Putaud, J.-P., Dinoi, A., Contini, D., Casans, A., Casquero-Vera, J. Andrés,
561 Crumeyrolle, S., Bourrianne, E., van Poppel, M., Dreesen, F. E., Harni, S., Timonen, H., Lampilahti, J., Petäjä, T., Pandolfi,
562 M., Hopke, P. K., Harrison, R. M., Alastuey, A. and Querol, X.: Source apportionment of ultrafine particles in urban Europe,
563 *Environ Int*, 109149, doi: 10.1016/j.envint.2024.109149, 2024.

564 Hammer, T., Roos, D., Giechaskiel, B., Melas, A. and Vasilatou, K.: Influence of soot aerosol properties on the counting
565 efficiency of instruments used for the periodic technical inspection of diesel vehicles, *Aerosol Research*, 2, 261–270, doi:
566 10.5194/ar-2-261-2024, 2024.

567 Jeong, C.-H., Salehi, S., Wu, J., North, M. L., Kim, J. Sung, Chow, C.-W. and Evans, G. J.: Indoor measurements of air
568 pollutants in residential houses in urban and suburban areas: Indoor versus ambient concentrations, *Sci Total Environ*,
569 133446, doi: 10.1016/j.scitotenv.2019.07.252, 2019.

570 Joshi, M., Khan, A., Anand, S. and Sapra, B. K.: Size evolution of ultrafine particles: Differential signatures of normal and
571 episodic events, *Environ Pollut*, Pt B, 354–360, doi: 10.1016/j.envpol.2015.10.001, 2016.

572 Junkermann, W. and Vogel, B.: Ultrafine particles over Germany – an aerial survey, *Tellus B: Chemical and Physical*
573 *Meteorology*, 1, 29250, doi: 10.3402/tellusb.v68.29250, 2022.

574 Kohler, M., Metzger, J. and Kalthoff, N.: Trends in temperature and wind speed from 40 years of observations at a 200-m
575 high meteorological tower in Southwest Germany, *Intl Journal of Climatology*, 1, 23–34, doi: 10.1002/joc.5157, 2018.

576 Kuhlbusch, T.: Diurnal variations of aerosol characteristics at a rural measuring site close to the Ruhr-Area, Germany,
577 *Atmospheric Environment*, 13–21, doi: 10.1016/S1352-2310(00)00489-1, 2001.

578 Kumar, P., Zavala-Reyes, J. C., Kalaiarasan, G., Abubakar-Waziri, H., Young, G., Mudway, I., Dilliway, C., Lakhdar, R.,
579 Mumby, S., Kłosowski, M. M., Pain, C. C., Adcock, I. M., Watson, J. S., Sephton, M. A., Chung, K. Fan and Porter, A. E.:
580 Characteristics of fine and ultrafine aerosols in the London underground, *Sci Total Environ*, Pt 1, 159315, doi:
581 10.1016/j.scitotenv.2022.159315, 2023.

582 Le Huong, T., Tuan, V. Anh and Vinh Man, N. Dinh: Commuter exposure to ultrafine particles (UFPs), lung deposited
583 surface area (LDSA), and noise in Ho Chi Minh City, Vietnam, *Journal of Transport & Health*, 101896, doi:
584 10.1016/j.jth.2024.101896, 2024.

585 German National Academy of Sciences, Leopoldina: Clean air – Nitrogen oxides and particulate matter in ambient air: Basic
586 principles and recommendations, 1st edition, Deutsche Akademie der Naturforscher Leopoldina, Halle (Saale), ISBN (eng)
587 978-3-8047-4013-6 , ISBN (ger) 978-3-8047-4012-9, 2019

588 Marten, R., Xiao, M., Rörup, B., Wang, M., Kong, W., He, X.-C., Stolzenburg, D., Pfeifer, J., Marie, G., Wang, D. S., Scholz,
589 W., Baccarini, A., Lee, C. Ping, Amorim, A., Baalbaki, R., Bell, D. M., Bertozzi, B., Caudillo, L., Chu, B., Dada, L., Duplissy,
590 J., Finkenzeller, H., Carracedo, L. Gonzalez, Granzin, M., Hansel, A., Heinritzi, M., Hofbauer, V., Kemppainen, D., Kürten,
591 A., Lampimäki, M., Lehtipalo, K., Makhmutov, V., Manninen, H. E., Mentler, B., Petäjä, T., Philippov, M., Shen, J., Simon,
592 M., Stozhkov, Y., Tomé, A., Wagner, A. C., Wang, Y., Weber, S. K., Wu, Y., Zauner-Wieczorek, M., Curtius, J., Kulmala,
593 M., Möhler, O., Volkamer, R., Winkler, P. M., Worsnop, D. R., Dommen, J., Flagan, R. C., Kirkby, J., Donahue, N. M.,
594 Lamkaddam, H., Baltensperger, U. and El Haddad, I.: Survival of newly formed particles in haze conditions, *Environ Sci*
595 *Atmos*, 3, 491–499, doi: 10.1039/d2ea00007e, 2022.

596 Matthes, M., Hartmann, I., Groll, A. and Riebel, U.: Investigation on application and performance of emission reduction
597 measures at a pellet boiler, *Biomass Conv. Bioref.*, 3, 301–313, doi: 10.1007/s13399-015-0187-1, 2016.

598 Matson, U.: Indoor and outdoor concentrations of ultrafine particles in some Scandinavian rural and urban areas, *Sci Total*
599 *Environ*, 1-3, 169–176, doi: 10.1016/j.scitotenv.2004.10.002, 2005.

600 Mohan, V., Kumar Soni, V. and Kumar Mishra, R.: Analysing the impact of day-night road traffic variation on ultrafine
601 particle number size distribution and concentration at an urban site in the megacity Delhi, *Atmospheric Pollution Research*,
602 4, 102065, doi: 10.1016/j.apr.2024.102065, 2024.

603 Moreno Camacho, F. L., Bustos Quevedo, D., Archila-Peña, D., Pachón, J. E., Rojas, N. Y., Mateus-Fontecha and Blanco,
604 K.: Number Concentration, Size Distribution, and Lung-Deposited Surface Area of Airborne Particles in Three Urban Areas
605 of Colombia, *Atmosphere*, 5, 558, doi: 10.3390/atmos16050558, 2025.

606 Mukherjee, A., Hartikainen, A., Joutsensaari, J., Basnet, S., Mesceriakovas, A., Ihalainen, M., Yli-Pirilä, P., Leskinen, J.,
607 Somero, M., Louhisalmi, J., Fang, Z., Kalberer, M., Rudich, Y., Tissari, J., Czech, H., Zimmermann, R. and Sippula, O.:
608 Black carbon and particle lung-deposited surface area in residential wood combustion emissions: Effects of an electrostatic
609 precipitator and photochemical aging, *Sci Total Environ*, 175840, doi: 10.1016/j.scitotenv.2024.175840, 2024.

610 Mylläri, F., Kuittinen, N., Aurela, M., Lepistö, T., Heikkilä, P., Salo, L., Markkula, L., Karjalainen, P., Kuula, J., Harni, S.,
611 Kyllönen, K., Similä, S., Jalkanen, K., Autio, J., Palonen, M., Valtatie, J., Häyrinen, A., Timonen, H., and Rönkkö, T.:
612 Primary particle emissions and atmospheric secondary aerosol formation potential from a large-scale wood pellet-fired
613 heating plant, *Aerosol Research Discuss.* [preprint], <https://doi.org/10.5194/ar-2025-14>, in review, 2025.

614 Neukirchen, C., Saraji-Bozorgzad, M. Reza, Mäder, M., Mudan, A. Paul, Czasch, P., Becker, J., Di Bucchianico, S., Trapp,
615 C., Zimmermann, R. and Adam, T.: Comprehensive elemental and physical characterization of vehicle brake wear emissions
616 from two different brake pads following the Global Technical Regulation methodology, *J Hazard Mater*, 136609, doi:
617 10.1016/j.jhazmat.2024.136609, 2025.

618 Norra, S., Song, J., Gebhardt, R., Bauer, J., Broß, M., Fuchs, M., Huber, T., Mack, D. and Saathoff, H.: Spatio-temporal
619 dynamics of aerosol distribution in an urban environment recorded in situ by means of a bike based monitoring system,
620 *Front. Environ. Sci.*, doi: 10.3389/fenvs.2023.749477, 2023.

621 Nyarku, M., Buonanno, G., Ofosu, F., Jayaratne, R., Mazaheri, M. and Morawska, L.: Ultrafine particles in key
622 microenvironments in rural and urban areas of Ghana, *Atmospheric Pollution Research*, 11, 101212, doi:
623 10.1016/j.apr.2021.101212, 2021.

624 Ohlwein, S., Kappeler, R., Kutlar Joss, M., Künzli, N. and Hoffmann, B.: Health effects of ultrafine particles: a systematic
625 literature review update of epidemiological evidence, *Int J Public Health*, 4, 547–559, doi: 10.1007/s00038-019-01202-7,
626 2019.

627 Palas GmbH, Product information of AQGuard Smart 2000, <https://www.palas.de/en/product/aq-guard-smart2000>, last
628 access: 25 November 2025

629 Pierce, J. R. and Adams, P. J.: Efficiency of cloud condensation nuclei formation from ultrafine particles, *Atmos. Chem.*
630 *Phys.*, 5, 1367–1379, doi: 10.5194/acp-7-1367-2007, 2007.

631 Ridolfo, S., Querol, X., Karanasiou, A., Rodríguez-Luque, A., Pérez, N., Alastuey, A., Jaén, C., van Drooge, B. L., Pandolfi,
632 M., Pedrero, M. and Amato, F.: Size distribution, sources and chemistry of ultrafine particles at Barcelona-El Prat Airport,
633 Spain, *Environ Int*, 109057, doi: 10.1016/j.envint.2024.109057, 2024.

634 Samad, A., Arango, K., Alvarez Florez, D., Chourdakis, I. and Vogt, U.: Assessment of Coarse, Fine, and Ultrafine Particles
635 in S-Bahn Trains and Underground Stations in Stuttgart, *Atmosphere*, 11, 1875, doi: 10.3390/atmos13111875, 2022.

636 Schiller, S. and Schmid, H.-J.: Highly efficient filtration of ultrafine dust in baghouse filters using precoat materials, *Powder*
637 *Technology*, 96–105, doi: 10.1016/j.powtec.2015.03.048, 2015.

638 Shrivastava, M., Fan, J., Zhang, Y., Rasool, Q. Z., Zhao, B., Shen, J., Pierce, J. R., Jathar, S. H., Akherati, A., Zhang, J.,
639 Zaveri, R. A., Gaudet, B., Liu, Y., Andreae, M. O., Pöhlker, M. L., Donahue, N. M., Wang, Y. and Seinfeld, J. H.: Intense
640 formation of secondary ultrafine particles from Amazonian vegetation fires and their invigoration of deep clouds and
641 precipitation, *One Earth*, 6, 1029–1043, doi: 10.1016/j.oneear.2024.05.015, 2024.

642 Song, J., Saathoff, H., Jiang, F., Gao, L., Zhang, H. and Leisner, T.: Sources of organic gases and aerosol particles and their
643 roles in nighttime particle growth at a rural forested site in southwest Germany, *Atmos. Chem. Phys.*, 11, 6699–6717, doi:
644 10.5194/acp-24-6699-2024, 2024.

645 Stacey, B.: Measurement of ultrafine particles at airports: A review, *Atmospheric Environment*, 463–477, doi:
646 10.1016/j.atmosenv.2018.10.041, 2019.

647 Sun, J., Birmili, W., Hermann, M., Tuch, T., Weinhold, K., Spindler, G., Schladitz, A., Bastian, S., Löschau, G., Cyrys, J.,
648 Gu, J., Flentje, H., Briel, B., Asbach, C., Kaminski, H., Ries, L., Sohmer, R., Gerwig, H., Wirtz, K., Meinhardt, F.,
649 Schwerin, A., Bath, O., Ma, N. and Wiedensohler, A.: Variability of black carbon mass concentrations, sub-micrometer
650 particle number concentrations and size distributions: results of the German Ultrafine Aerosol Network ranging from city
651 street to High Alpine locations, *Atmospheric Environment*, 256–268, doi: 10.1016/j.atmosenv.2018.12.029, 2019.

652 Teutscher, D., Bukreev, F., Kummerländer, A., Simonis, S., Bächler, P., Rezaee, A., Hermansdorfer, M. and Krause, M. J.:
653 A digital urban twin enabling interactive pollution predictions and enhanced planning, *Building and Environment*, 113093,
654 doi: 10.1016/j.buildenv.2025.113093, 2025.

655 Thieringer, J. R. D., Szabadi, J., Meyer, J. and Dittler, A.: Impact of Residential Real-World Wood Stove Operation on Air
656 Quality concerning PM_{2.5} Immission, *Processes*, 3, 545, doi: 10.3390/pr10030545, 2022.

657 Trebs, I., Lett, C., Krein, A. and Junk, J.: Air quality impacts of aviation activities at a mid-sized airport in central Europe,
658 *Atmospheric Pollution Research*, 3, 101696, doi: 10.1016/j.apr.2023.101696, 2023.

659 Trechera, P., Garcia-Marlès, M., Liu, X., Reche, C., Pérez, N., Savadkoohi, M., Beddows, D., Salma, I., Vörösmarty, M.,
660 Casans, A., Casquero-Vera, J. Andrés, Hueglin, C., Marchand, N., Chazeau, B., Gille, G., Kalkavouras, P., Mihalopoulos,
661 N., Ondracek, J., Zikova, N., Niemi, J. V., Manninen, H. E., Green, D. C., Tremper, A. H., Norman, M., Vratolis, S.,
662 Eleftheriadis, K., Gómez-Moreno, F. J., Alonso-Blanco, E., Gerwig, H., Wiedensohler, A., Weinhold, K., Merkel, M.,
663 Bastian, S., Petit, J.-E., Favez, O., Crumeyrolle, S., Ferlay, N., Martins Dos Santos, S., Putaud, J.-P., Timonen, H.,
664 Lampilahti, J., Asbach, C., Wolf, C., Kaminski, H., Altug, H., Hoffmann, B., Rich, D. Q., Pandolfi, M., Harrison, R. M.,
665 Hopke, P. K., Petäjä, T., Alastuey, A. and Querol, X.: Phenomenology of ultrafine particle concentrations and size
666 distribution across urban Europe, *Environ Int*, 107744, doi: 10.1016/j.envint.2023.107744, 2023.

667 Tremper, A. H., Jephcote, C., Gulliver, J., Hibbs, L., Green, D. C., Font, A., Priestman, M., Hansell, A. L. and Fuller, G. W.:
668 Sources of particle number concentration and noise near London Gatwick Airport, *Environ Int*, 107092, doi:
669 10.1016/j.envint.2022.107092, 2022.

670 Trojanowski, R. and Fthenakis, V.: Nanoparticle emissions from residential wood combustion: A critical literature review,
671 characterization, and recommendations, *Renewable and Sustainable Energy Reviews*, 515–528, doi:
672 10.1016/j.rser.2019.01.007, 2019.

673 Turner, A., Wolfe, C. and Ryan, P. H.: Personal exposure to ultrafine particles in multiple microenvironments among
674 adolescents, *J Expo Sci Environ Epidemiol*, 5, 878–885, doi: 10.1038/s41370-023-00638-7, 2024.

675 Utinger, B., Barth, A., Paul, A., Mukherjee, A., Campbell, S. John, Müller, C.-M., Ihalainen, M., Yli-Pirilä, P., Kortelainen,
676 M., Fang, Z., Martens, P., Somero, M., Louhisalmi, J., Hohaus, T., Czech, H., Sippula, O., Rudich, Y., Zimmermann, R. and

677 Kalberer, M.: Emission dynamics of reactive oxygen species and oxidative potential in particles from a petrol car and wood
678 stove, *Aerosol Research*, 1, 205–218, doi: 10.5194/ar-3-205-2025, 2025.

679 Wang, D., Li, Q., Shen, G., Deng, J., Zhou, W., Hao, J. and Jiang, J.: Significant ultrafine particle emissions from residential
680 solid fuel combustion, *Sci Total Environ*, 136992, doi: 10.1016/j.scitotenv.2020.136992, 2020.

681 Wang, Y., Hopke, P. K., Chalupa, D. C. and Utell, M. J.: Long-term study of urban ultrafine particles and other pollutants,
682 *Atmospheric Environment*, 40, 7672–7680, doi: 10.1016/j.atmosenv.2010.08.022, 2011.

683 World Health Organization: WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen
684 dioxide, sulfur dioxide and carbon monoxide, <https://www.who.int/publications/i/item/9789240034228>, last access: 07.
685 August 2025 (2021).

686 World Health Organization: Air quality guidelines global update 2005, [https://www.who.int/publications/i/item/WHO-SDE-](https://www.who.int/publications/i/item/WHO-SDE-PHE-OEH-06.02)
687 [PHE-OEH-06.02](https://www.who.int/publications/i/item/WHO-SDE-PHE-OEH-06.02), last access: 07. August 2025 (2005).

688 Yin, S., Lu, Z., Zhang, Y., Song, L., Bi, S., Luo, X., Yao, L., Bi, X., Bo, H. and Feng, Y.: Characteristics of number
689 concentration, size distribution and components of particulate matter emitted from a typical large civil airport, *Sci Total*
690 *Environ*, 172040, doi: 10.1016/j.scitotenv.2024.172040, 2024.

691 Zhang, H., Huang, W., Shen, X., Ramisetty, R., Song, J., Kiseleva, O., Holst, C. Claus, Khan, B., Leisner, T. and Saathoff,
692 H.: Aerosol composition, air quality, and boundary layer dynamics in the urban background of Stuttgart in winter, *Atmos.*
693 *Chem. Phys.*, 18, 10617–10637, doi: 10.5194/acp-24-10617-2024, 2024.

694 Zhu, Y., Hinds, W. C., Kim, S. and Sioutas, C.: Concentration and size distribution of ultrafine particles near a major
695 highway, *J Air Waste Manag Assoc*, 9, 1032–1042, doi: 10.1080/10473289.2002.10470842, 2002.

696